

Immunocytochemical Study on Microtubule Reorganization in HL-60 Leukemia Cells Undergoing Apoptosis

Shahin Ahmadian*, Yaghub Pazhang and Ahmad Sharifabrizi

Dept. of Biochemistry, Institute of Biochemistry and Biophysics, University of Tehran, Tehran, Iran

Received 11 December 2006; revised 13 May 2007; accepted 20 May 2007

ABSTRACT

Background: Microtubules (MT) are important components of cell cytoskeleton and play key roles in cell motility, mitosis and meiosis. They are also the targets of several anticancer agents which indicating their importance in maintaining cell viability. Microtubular reorganization contributing to apoptotic morphology occurs in normal and neoplastic cells undergoing apoptosis induced by cytotoxic drugs. The aim of this study was to correlate the changes in the MT with behavior of the γ -tubulin in apoptotic cell, and to see if apoptotic MT showed biochemical characteristics of stable MT. **Methods:** Apoptosis was induced in the human leukemia cells (HL-60) by treatment with 1 μ M of all-trans retinoic acid over a 5-day period. The time course of changes was assessed using flow cytometry, DNA fragmentation and immunocytochemistry in cells labeled for α -tubulins, acetylated α -tubulin and γ -tubulin. **Results:** The results indicated that γ -tubulin content is increased after cells have gone through the apoptosis with a diffuse cytoplasmic pattern. α -tubulin did not reveal any specific pattern of polymerization in apoptotic cells and acetylated α -tubulin content was also decreased in comparison with non-apoptotic cells. **Conclusion:** Our results support the idea that microtubule reorganization is an important factor of the mammalian cells response to apoptosis, and the altered properties of the MT did not reflect changes in function as apoptosis progresses. *Iran. Biomed. J. 11 (4): 209-214, 2007*

Keywords: Apoptosis, Microtubule, γ -tubulin, Immunocytochemistry

INTRODUCTION

Apoptosis is a term used to describe the terminal morphological and biochemical events seen in programmed cell death [1]. Apoptosis has recently been the subject of great interest because it has been clearly demonstrated to mediate cell death, not only during development but also in neoplasia in response to cancer chemotherapy and radiation [2, 3]. Studies on leukemia and lymphoma cells have been significant in determining the importance of apoptosis in the cancer process. Considerable morphological change occurs in apoptosis and it has been proposed that the cytoskeleton is involved. Some studies have directly implicated actin in the apoptotic process, particularly in the production of apoptotic bodies [4, 5]. Our studies have focused on microtubules (MT), which we have shown is reorganized, during apoptosis.

MT perform diverse functions in mammalian cells, including transport of chromosomes and

membrane enclosed organelles. This functional diversity is controlled at the transcriptional level, by several post-translational modifications, and also by differential binding of microtubule-associated proteins (MAP) [6-8]. MT exist in a state of dynamic instability *in vivo* [9-11] and *in vitro* [12]. In mammalian cells the MT nucleate from the microtubule organizing center (MTOC) or centrosome which stabilizes the minus end. Most cytoplasmic MT have rapid cycles of polymerization-depolymerization but there is a small proportion of relatively stable, slow turn over MT [13, 14]. Stable MT are biochemically distinct from newly formed MT. Certain post-translational modifications of α -tubulin, namely acetylation of Lys 40 and detyrosination of the c-terminal tyrosyl residue, are frequently found associated with stable MT [15-18]. Evidence suggests that such post-translational modifications occur preferentially in the microtubule and in a time-dependent manner [14]. Acetylation of α -tubulin is also associated with

*Corresponding Author; Tel. (+98-21) 6695 6983; Fax: (+98-21) 6640 4680; E-mail: ahmadian@ibb.ut.ac.ir

MT resistant to depolymerization by anti-mitotic drugs and cold shock [15, 17], and is regarded as marker of stable MT.

In the present work, we have used immunocytochemical staining and flow cytometry to examine changes in microtubule network and cell content of acetylated and non-acetylated α -tubulin in HL-60 cells undergoing all-trans retinoic acid (ATRA) induced apoptosis.

MATERIALS AND METHODS

Cell culture and induction of apoptosis. An established HL-60 human myeloid leukemia cell line was used for experiments (Iranian National Cell Bank, Pasteur Institute of Iran, Tehran, NCBI code C427). Cells were maintained in suspension culture at 37°C under 5% CO₂/air in RPMI 1640 (Gibco Inc. UK) supplemented with 10% FCS and antibiotics: 100 U/ml of penicillin and 100 µg/ml of streptomycin. The Cells were sub-cultured three times weekly. ATRA (Sigma Chemical Co., St. Louis, MO, USA), was used as an inductor of apoptosis. Immediately after inoculation, a solution of ATRA was added at a final concentration of 10⁻⁶ M. An equivalent volume of ethanol was added to control cells during the apoptosis process; the medium was removed at 48-h intervals and replaced with a fresh one containing ATRA. Simultaneously, fresh medium free of ATRA was also changed in the control flasks. Cell membrane integrity was confirmed using the trypan blue staining method.

Immunocytochemistry. Immunofluorescence microscopy was performed according to the reference [19] with some modifications. Cells were first centrifuged on poly-L-lysine coated coverslips, then the coverslips bearing cells were immersed in -20°C methanol for 5 minutes. Coverslips were then rehydrated with PBS, and were washed three times with Ab buffer (PBS, 3% BSA, 0.1% Triton X-100, 0.02% azide). The medium was supplemented with DAPI fluorescent dye (Sigma-Aldrich, Inc. USA) at a concentration of 1 µg/ml to stain DNA. Primary antibodies anti α -tubulin (Cat no. T-9026), α -acetylated tubulin (Cat. no. T-6793) and γ -tubulin (Cat. no. T-3559), all from Sigma-Aldrich Inc. (USA) chemicals were diluted in Ab buffer and added to the coverslips and incubated for 30 minutes in the room temperature. The samples were washed three times with PBS and incubated in secondary fluorescent conjugated antibody for 30 minutes. The

coverslips were mounted on slides using glycerol, observed in a Leitz fluorescence microscope and fluorescence micrographs were recorded using CCD camera (Hitachi, kp-D20BP, Japan). The apoptosis process was detected in individual cells on the basis of characteristic changes in nuclear morphology (chromatin collapse and nucleus breakdown). In the apoptosis cells, MT and γ -tubulin distribution were assessed and compared with the cytoskeleton of interphase and mitotic cells.

Hoechst assay. The cells were washed with PBS, and then were fixed for 5 minutes in ice cold 3% paraphormaldehyde solution in PBS. Cells were centrifuged and washed with PBS, and then were resuspended in 50 µl Hoechst 333492 solution as described before [20]. The cells were mounted on a glass slide using glycerol. Cells were visualized with a DAPI filter (Ex = 330 A, Emission = 480) under an immunofluorescence microscope.

Immunoblotting. Western-blot analysis was performed as described before [20]. Briefly, equal numbers of cells were lysed using a 30-s sonication in the lysis buffer [DTT 1 mM, EDTA 1 M, PMSF (10 µM) and the protein concentration in the lysate was measured using Bradford assay. Equal amounts of proteins (10 µg) were loaded into the mini gel (10% Lamelli acrylamide gel) electrophoresed at constant voltage of 100 V for 3 hours, and then transferred overnight to nitrocellulose membranes. Membranes were incubated for 2 hours in the blocking solution containing 5% skim milk, then incubated in primary antibodies against α , α -acetylated and γ -tubulin. The Membranes were washed in washing buffer and were further developed in the secondary antibodies for 1 h (ALK-P conjugated for γ -tubulin HRP-conjugated and for α - and α -acetylated detection) and finally was incubated in the corresponding substrate for secondary antibodies after a three-time washing in the wash buffer until the bands were developed.

DNA laddering. DNA fragmentation was performed according to Porter and Lee [21] with some modifications. Cells were lysed with digestion buffer (Tris-HCl, 10 mM; NaCl, 100 mM; EDTA, 25 mM; SDS 1%, proteinase k, 0.2 mg/m) and were incubated at 50°C for 2 hours. The lysate was extracted with phenol/chloroform/isoamyl alcohol and twice with phenol/chloroform/isoamyl alcohol (25/24/1) and once with chloroform/isoamyl alcohol (24/1). DNA was precipitated by adding ammonium

acetate to final concentration of 2.5 M, mixed and 2 volumes of pure ethanol was added, and then incubated at -20°C overnight. DNA was pelleted by centrifugation at $12,000 \times g$ for 30 minutes, washed with 70% ethanol, air dried and dissolved finally in the Tris-EDTA buffer (Tris-HCl (pH = 8), 10 mM; EDTA, 1 mM) and visualized using ethidium bromide containing agarose gels.

Annexin-5 assay. Annexin-5 assay was performed as described before [21] using commercial kit (from IQ Products Company, The Netherlands). Briefly, cells were washed twice in calcium buffer, incubated at 4°C for 20 minutes at dark in Annexin-5 FITC washed with calcium buffer and incubated for 10 minutes in propidium iodide and analyzed using an FAC scan (Becton Dickinson) flow cytometer.

RESULTS

Induction of apoptosis by ATRA. The human HL-60 leukemia cell line was treated with $1 \mu\text{M}$ ATRA for various periods of time, and cells were harvested for morphological, flow cytometric and DNA analysis. The morphological changes typical of apoptosis were visible 4-5 days after treatment. DNA fragmentation was observed at five days post-treatment (Fig. 1). Flow cytometric analysis of ATRA-treated cells by Annexin V is shown in Figure 2. The percentage of cells positive for Annexin 5 staining was 10.87% four days after treatment and significantly increased (24.43%) five days after treatment. For non-treatment cells, the percentage of positive cells was h.54% (data not shown).



Fig. 1. Detection of apoptosis by DNA laddering. Apoptosis was documented after 5 days of ATRA treatment.

Relative content of γ , α and α -acetylated tubulin. The content of γ -tubulin was significantly increased following apoptosis. Acetylated- α tubulin and α -tubulin were decreased following apoptosis induction in comparison with non-apoptotic control cells as identified by Western-blotting (Fig. 3).

Changes in dynamic of MT following apoptosis. A dual staining both for the nucleus and microtubule network allowed identifying individual cells by the morphology of nucleus. Apoptosis was characterized by fragmented nuclei as identified by Hoechst assay, so those cells were at the latest stage of apoptosis. Examination of α -tubulin-stained cytocentrifuge preparations showed that all cells both treated and untreated showed some tubulin positivities. In apoptotic cells, two patterns of MT were observed: lack of staining and faint, disperse pattern of staining. No obvious MTOC was observe (Fig. 4 a and b). In normal cells, half displayed polymerized tubulin structures in cytoplasm and a




Fig. 2. Indications of apoptosis as analyzed by Annexin-5 assay. Percentage of cells positive both for propidium iodide and annexin-5 staining was significantly increased after 5 days of ATRA treatment (**right**) in comparison to day 0 treatment (**left**).




Fig. 3. Immunoblot analysis of α -tubulin (a), acetylated- α -tubulin (b) and γ -tubulin (c) in normal non-apoptosis and apoptotic HL-60. N, non-apoptotic; A, apoptotic.

Faint microtubular network were seen on or around the nucleus (Fig. 4 c and d). Immunofluorescence staining for the α -acetylated MT showed changes comparable to that of α -tubulin, but there was no consistent staining of α -acetylated cell in apoptotic cells (Fig. 5 a and b). In non-apoptotic cells, a fine rim of acetylated α -tubulin was seen around the nucleus (Fig. 5 c and d).

γ -tubulin distribution in different stages of apoptosis. In normal non-apoptotic cells, the staining was mostly confined to the centrosomal areas and a faint, localized and very weak staining for γ -tubulin was also observed in cytoplasm (Fig. 6 a and b). In the apoptotic cells, there was no centrosomal staining but areas of cytoplasmic staining were more intense than non-apoptotic cells (Fig. 6 c and d).




Fig. 4. Immunofluorescent staining of α -tubulin in (a and b) apoptotic and (c and d) non-apoptotic HL-60 cells. Cells were fixed and stained with α -tubulin monoclonal antibody and fluorescein-conjugated secondary antibodies, as described in material and methods. Nuclear morphology in the same subset which signifies the stage of apoptosis is clarified using Hoechst staining on the same cells (c and d). A, apoptotic; N, non-apoptotic; D, differentiated.

DISCUSSION

Pattern of MT in a typical tissue culture cell is usually in the form of MT distributed to their minus ends clustered together and their plus ends extending out into the cytoplasm. The canonical model for how such arrays are generated is that MT, nucleated at centrosomes, grow through the cytoplasm. Dynamic instability describes the stochastic behavior of MT in which individual MT transit between phases of growth and fast shrinkage. Upon contact with capture sites, they would be stabilized, forming patterns.

Tubulin self-assembles *in vitro* when a critical concentration is exceeded, indicating that MTOC are not absolutely required for MT nucleation. However, even in the normal cytoplasm of living cells, concentration of tubulin is not high enough to account for a free nucleation of MT, independent of an MTOC [22]. On the other hand, over-expressed γ -tubulin is shown to be associated with ectopic microtubule nucleation outside the centrosome in mammalian cell cytoplasm. This can be considered as an explanation of our observed increase in γ -tubulin content after apoptosis. It is possible that γ -tubulin might be increased in response to a decreased tubulin content for the formation of multiple MTOC that might form in apoptosis as suggested by other investigators [23], possibly as a defense mechanism against progression of apoptosis. Increased cytoplasmic staining of γ -tubulin observed in our study is also supportive for this hypothesis. Induction of apoptosis by cytostatic drug taxol in HL-60 cells [5] and VP-16 in CCRF-CEM cells also




Fig. 5. Immunofluorescent staining of acetylated α -tubulin in (a) apoptotic and (c) non-apoptotic HL-60 cells which shows a fine labeling of acetylated α -tubulin around the nucleus. Nuclear morphology in the same subset is clarified using Hoechst staining on the same cells (b and d). A, apoptotic; N, non-apoptotic.

induces multiple MTOC [24]. On the other hand, the increased γ -tubulin content might only reflect that its high level in differentiated cells have not returned to normal yet. ATRA-induced apoptosis in HL-60 cells is claimed to be a process independent of differentiation [25, 26] which argues against this assumption, although a recent proteomics analysis [27] have suggested that differentiation is a prerequisite of apoptosis induced by ATRA.

Dynamic MT rapidly grows using free unmodified tubulin, and frequently shrinks, never existing long enough to be significantly (detectably) modified. Stable MT, being older, simultaneously accumulate significant concentrations of post-translationally modified tubulin forms (acetylated and tyrosinated) [13]. Selective stabilization of MT has been discussed as a general model for morphogenesis and acetylation and tyrosination of tubulin is shown to be a major event in differentiated related phenomena [28]. In this model, a cell in the absence of an outside morphogenetic signal randomly polymerizes MT in all directions with the rapid turn-over predicted by the property of dynamic instability *in vitro* as well as the behavior observed in living cells.

According to our observations, apoptotic cells

would have less abundant α and α -acetylated tubulin and will have disperse cytoplasmic γ -tubulin, α and α -acetylated tubulin without having an organized structure of any of these components. This is comparable to the previous report showing microtubule disruption in ATRA-induced apoptosis of HL-60 cells [29]. We believe that these findings about the role of γ -tubulin and microtubule cytoskeleton in APL (Acute promyelocytic leukemia) apoptosis would be enough justification for performing further studies on the function of γ -tubulin in ATRA resistant subsets of leukemia cases.

ACKNOWLEDGEMENTS

We wish to thank M. Shafieezadeh for her help to prepare the manuscript, K. Bolouri for assistance in preparation of photographic material. This work was supported by Research Council of the University of Tehran (graduate program support grant No: 6401007/6/01).




Fig. 6. Immunofluorescent staining of γ -tubulin in (a) non-apoptotic and (c) apoptotic HL-60 cells. In non-apoptotic cells, staining is mostly confined to the centrosomal area. In apoptotic cells, there is no centrosomal staining, but the cytoplasmic staining is more intense than non-apoptotic cells. Nuclear morphology in the same subset is clarified using Hoechst staining on the same cells (b and d). N, non-apoptotic; A, apoptotic.

REFERENCES

1. Wyllie, A.H. (1992) Apoptosis and the regulation of cell numbers in normal and neoplastic tissues: an overview. *Cancer Metastasis Rev.* 11 (2): 95-103.
2. Hickman, J.A. (1992) Apoptosis induced by anticancer drugs. *Cancer Metastasis Rev.* 11 (2): 121-139.
3. Eastman, A. (1990) Activation of programmed cell death by anticancer agents: cisplatin as a model system. *Cancer Cells* 2 (8-9): 275-280.
4. Cotter, T.G., Lennon, S.V., Glynn, J.M. and Green, D.R. (1992) Microfilament-disrupting agents prevent the formation of apoptotic bodies in tumor cells undergoing apoptosis. *Cancer Res.* 52 (4): 997-1005.
5. Grzanka, A., Grzanka, D. and Orlikowska, M. (2003) Cytoskeletal reorganization during process of apoptosis induced by cytostatic drugs in K-562 and HL-60 leukemia cell lines. *Biochem. Pharmacol.* 66 (8): 1611-1617.
6. MacRae, T.H. (1992) Towards an understanding of microtubule function and cell organization: an overview. *Biochem. Cell Biol.* 70 (10-11): 835-841.
7. Joshi, H.C. and Cleveland, D.W. (1990) Diversity among tubulin subunits: toward what functional end? *Cell Motil. Cytoskeleton* 16 (3): 159-163.
8. Mandelkow, E. and Mandelkow, E.M. (1990) Microtubular structure and tubulin polymerization. *Curr. Opin. Cell Biol.* 2 (1): 3-9.
9. Sammak, P.J. and Borisy, G.G. (1988) Direct observation of microtubule dynamics in living cells. *Nature* 332 (6166): 724-726.
10. Bulinski, J.C. and Gunderson, G.G. (1991) Stabilization of post-translational modification of microtubules during cellular morphogenesis. *Bioassays* 13 (6): 285-293.
11. Mitcheson, T. and Kirschner, M. (1984) Dynamic instability of microtubule growth. *Nature* 312 (5991): 237-242.
12. Sheldon Wadsworth, P. (1993) Observation and quantification of individual microtubule behavior *in vivo*: Microtubule dynamics are cell type specific. *J. Cell Biol.* 120: 935-945.
13. Schulze, E., Asai, D.J., Bulinski, J.C. and Kirschner, M. (1987) Posttranslational modification and microtubule stability. *J. Cell Biol.* 105 (5): 2167-2177.
14. Webster, D.R. and Borisy, G.G. (1989) Microtubules are acetylated in domains that turn over slowly. *J. Cell Sci.* 92 (Pt 1): 57-65.
15. Khawaja, S., Gunderson, G.G. and Bulinski, J.C. (1988) Enhanced stability of microtubules enriched in deetyrosinated tubulin is not a direct function of deetyrosination level. *J. Cell Biol.* 106 (1): 141-149.
16. Idriss, H., Stammers, D.K., Ross, C.K. and Burns, R.G. (1991) The dynamic instability of microtubules is not modulated by alpha-tubulin tyrosinylation. *Cell Motil. Cytoskeleton* 20 (1): 30-37.
17. Piperno, G., LeDizet, M. and Chang, X. (1987) Microtubules containing acetylated alpha-tubulin in mammalian cells in culture. *J. Cell Biol.* 104 (2): 289-302.
18. Huang, B. (1990) Genetics and biochemistry of centrosomes and spindle poles. *Curr Opin. Cell Biol.* 2 (1): 28-32.
19. Bobiniec, Y., Fukuda, M. and Nishida, E. (2000) Identification and characterization of *Caenorhabditis elegans* γ -tubulin in dividing cells and differentiated tissues. *J. Cell Science* 113: 3747-3759.
20. Roy, M.K., Thalang, V.N., Trakoontivakorn, G. and Nakahara, K. (2004) Mechanism of mahanine-induced apoptosis in human leukemia cells (HL-60). *Biomed. Pharmacol.* 67:41-51.
21. Porter, L.A. and Lee, J.M. (2001) α -, β - and γ -tubulin polymerization in response to DNA damage. *Exp. Cell Res.* 270:151-158.
22. Hyman, A. and Karsenti, E. (1998) The role of nucleation in patterning microtubule networks. *J. Cell Sci.* 111 (Pt 15): 2077-2083.
23. Shu, H. and Joshi, H.C. (1995) Gamma-tubulin can both nucleate microtubule assembly and self-assemble into novel tubular structures in mammalian cells. *J. Cell Biol.* 30 (5): 1137-1147.
24. Pittman, S., Geyp, M., Fraser, M., Ellem, K., Peaston, A. and Ireland, C. (1997) Multiple centrosomal microtubule organizing centers and increased microtubule stability are early features of VP-16-induced apoptosis in CCRF-CEM cells. *Leuk. Res.* 21 (6): 491-499.
25. Olins, A.L., Herrmann, H., Lichter, P. and Olins, D.E. (2000) Retinoic acid differentiation of HL-60 cells promotes cytoskeletal polarization. *Exp. Cell Res.* 254 (1): 130-142.
26. Garattini, E. and Terao, M. (2006) Granulocytic maturation in cultures of acute myeloid leukemia is not always accompanied by increased apoptosis. *Leuk. Res.* 30 (5): 519-520.
27. Guo, X., Ying, W., Wan, J., Hu, Z., Qian, X., Zhang, H. He, F. (2001) Proteomic characterization of early-stage differentiation of mouse embryonic stem cells into neural cells induced by all-trans retinoic acid *in vitro*. *Electrophoresis* 22 (14): 3067-3075.
28. Gunderson, G.G., Khawaja, S. and Bulinsky, J.C. (1989) Generation of a stable, posttranslationally modified microtubule array is an early event in myogenic differentiation. *J. Cell Biol.* 109 (5): 2275-2288.
29. Veselska, R., Zitterbart, K. and Jelinkova, S. (2003) Specific cytoskeleton changes during apoptosis accompanying induced differentiation of HL-60 myeloid leukemia cells. *Oncol. Rep.* 10 (4):1049-1058.