

# Anti-Proliferative Properties, Biocompatibility, and Chemical Composition of Different Extracts of *Plantago major* Medicinal Plant

Samaneh Rahamouz-Haghighi<sup>1</sup>, Khadijeh Bagheri<sup>1</sup>, Hossein Danafar<sup>2</sup> and Ali Sharafi<sup>2,3\*</sup>

<sup>1</sup>Department of Plant Production and Genetics, Faculty of Agriculture, University of Zanjan, Zanjan, Iran; <sup>2</sup>Zanjan Pharmaceutical Biotechnology Research Center, Zanjan University of Medical Sciences, Zanjan, Iran; <sup>3</sup>Department of Pharmaceutical Biotechnology, School of Pharmacy, Zanjan University of Medical Sciences, Zanjan, Iran

Received 18 March 2020; accepted 27 July 2020; published online 20 January 2021

## ABSTRACT

**Background:** To study the anticancer activity of *Plantago major*, we assessed the effect of ethanolic, methanolic and acetonetic extracts of this plant on HCT-116, SW-480, and HEK-293 cell lines as control. **Methods:** The cytotoxic activity, biocompatibility, and toxicity were evaluated by MTT assay, hemolysis, and *Artemia salina*-LD<sub>50</sub> (on mice) tests, respectively. The analysis of the extracts was performed by GC-MS analysis. **Results:** The results showed that all the extracts had the most antiproliferative properties on the HCT-116 cell line. The *P. major* root extract was more effective than the aerial parts, and IC<sub>50</sub> values for ethanolic, methanolic and acetonetic root extracts were 405.59, 470.16, and 82.26 µg/mL, respectively on HCT-116 cell line at 72 h. Hemolysis degree of the ethanolic extract of aerial and root parts were approximately 1% at 400 µg/mL. Using the ethanolic extracts, the *Artemia* survived every concentration, and no toxicity was observed. One week after the oral administration of different parts of *P. major* extracts, none of the mice died, even those were administered 2000 mg/kg. The results of GC/MS analysis showed that *P. major* extracts contain potential anticancer compounds, such as stearic acid (8.61%) in aerial parts of methanolic extract and 1,2- Benzenedicarboxylic acid, mono(2-ethylhexyl)ester (88.07% and 40.63%) in aerial and root parts of acetonetic extract of *P. major*. **Conclusions:** Our findings suggest that the *P. major* is a source of potential compounds with antiproliferative properties. **DOI: 10.29252/ibj.25.2.106**

**Keywords:** Gas chromatography-mass spectrometry, HCT-116 cells, Hemolysis, Lethal dose 50

**Corresponding Author:** Ali Sharafi

<sup>2</sup>Zanjan Pharmaceutical Biotechnology Research Center, Zanjan University of Medical sciences, Zanjan, Iran; Postal Code: 45139-56184; Tel.: (+98-24) 33473635; E-mail: alisharafi@zums.ac.ir

## INTRODUCTION

Since the most anticancer drugs are associated with severe side effects, new drugs with higher efficacy and fewer adverse effects are needed<sup>[1]</sup>. Natural-based active compounds, such as curcumin, artemisinin, and taxol, have been used for cancer treatment<sup>[2]</sup>. Plant secondary metabolites, owing to their antiproliferative features, have also been utilized as drugs for the treatment of cancer<sup>[3]</sup>.

*Plantago major* is a medicinal herb from the *Plantaginaceae*<sup>[4]</sup>. Based on traditional medicine, it has various medicinal uses without significant side effects. Most therapeutic effects of *P. major* described in

traditional medicine have not fully been investigated; thus, complementary studies are required to reveal more medicinal characteristics of this plant. Having a broad range of medical application in many countries, *P. major* can be exploited as a source of cost-effective drug candidate<sup>[5]</sup>. *P. major* contains important compounds, such as polysaccharides, flavonoids<sup>[6,7]</sup>, phenolic compounds<sup>[8]</sup>, monoterpenoids, and triterpenoids<sup>[9]</sup>, lipids, and caffeic acid derivatives<sup>[10]</sup>. Moreover, caffeoyl phenylethanoid glycosides and iridoids glycosides are the most distinctive category of compounds that show important correspondence in their chemotaxonomy significance<sup>[11-13]</sup>.

Terpenoids are relatively non-toxic substances; these

## List of Abbreviations:

**GC-MS**, gas chromatography-mass spectrometry; **LD<sub>50</sub>**, lethal dose 50; **OECD**, Organisation for Economic Cooperation and Development

compounds have great potential to be applied as chemotherapeutic agents in battling cancer<sup>[14]</sup>. As the main constituents of *Plantago* extract, triterpenoids have exhibited superb antiproliferative effects and apoptosis on cancer cells<sup>[15-17]</sup>. Fatty acids are one of the most important medicinal sources with multi-biological (antimicrobial and antifungal) activities<sup>[18]</sup>.

Till now, various therapeutic effects of *P. major* have been revealed<sup>[5]</sup>. The antibacterial activity of *P. major* has been suggested in Holetz *et al.*'s study<sup>[19]</sup>. The aquatic extract of *P. major* has been indicated to have an effective antileukemia, anticarcinoma, and antiviral activities, along with modulate cell-mediated immunity<sup>[20]</sup>. Moreover, the aerial parts of this plant have been used for the treatment of anemia and hematopoietic disorders; however, its possessions on hematopoietic cells, mainly on totipotential stem cells, remain unidentified<sup>[21]</sup>. It has been reported that *P. major* L. has an uterotonic action in the guinea pig, a prophylactic influence on mammary cancer in mice, and a protective effect against systemic *Streptococcus pneumoniae* infection in mice<sup>[22-24]</sup>. *P. major* extract has a significant inhibitive effect on Ehrlich ascites tumor<sup>[25]</sup>.

To the best of our knowledge there is no previous research, associated with the aerial and root parts of *P. major* on colon cancer cells (HCT-116 and SW-480 cell lines). Hence, we aimed to investigate the effect of the whole aerial parts and roots of this plant on two cell lines of colorectal cancer by MTT assay, hemolysis, toxicity activity on *Artemia salina* and LD<sub>50</sub> determination, along with the analysis of the volatile compounds of this plant species.

## MATERIALS AND METHODS

### Herbal materials

The plants were collected from Zanjan city of Iran (geographical coordinates of the collection sites: 36°41'15.5"N 48°24'02.2"E) and authenticated at the Department of Botany, University of Zanjan, Iran. All the plant sections were cut into small pieces and dried in the shade at room temperature for one week.

### Plant extraction

Approximately 20 grams of the dried aerial (stem and leaf) and root parts of *P. major* was ground to a coarse powder and extracted by the reflux method using 200 mL of ethanol or methanol for 8 h and acetone for 3 hours. The extracts were then filtered and concentrated in an evaporator under pressure at 35-45 °C for 75 min. The extracts were kept at 4 °C<sup>[26]</sup>.

### Cell line culture

HCT-116, SW-480, and HEK-293 cell lines were obtained from the Pasteur Institute of Iran, Tehran and were cultured in DMEM and RPMI-1640 media supplemented with penicillin-streptomycin (1%) and 10% FBS in 5% CO<sub>2</sub> incubator at 37 °C.

### Viability assay

The inhibitory effect of the ethanolic, methanolic and acetic extracts of *P. major* on the cell lines was determined by MTT assay. The cells were seeded onto a 96-well plate at a density of  $7 \times 10^3$  cells/well. The cells were attached and grown for 24 h to reach 70-80% confluency. Subsequently, 10 mg of concentrated extracts was dissolved in 100 µL of DMSO and dissolved in 900 µL of the culture medium (DMEM or RPMI-1640) for the preparation of 25, 50, 100, 200, and 400 µg/mL of the extracts using dilution method. The extracts were filtered by 0.45-µm membrane filters. Medium and DMSO were considered as + and - control, respectively. Cells were treated with the prepared extracts and incubated for 1-3 days. Thereafter, 20 µL of MTT (5 mg/mL) was added and kept at an incubator for four hours. The media was then removed by aspiration. DMSO (200 µL) was added to each well to dissolve the obtained formazan. The absorbance was read by an ELISA plate reader (Tecan Infinite M200, Austria) at 570 and 690 nm, and the OD was documented<sup>[27]</sup>. The inhibitory rate of the cell growth was considered by: % Growth inhibition =  $(1 - \text{OD extract-treated}) / \text{OD negative control} \times 100$

### Hemolytic toxicity

To examine the biocompatibility of the ethanolic extract of aerial and root parts, the hemolysis assay was performed<sup>[28]</sup>. Freshly prepared human RBCs collected in ethylenediaminetetraacetic acid-containing tube were washed with isotonic PBS (pH 7.4) by centrifugation at 1663 ×g for 5 min. Next, the tube containing the erythrocytes was resuspended in the same medium at a final hematocrit of 5%. Then aerial and root extracts with the concentration of 25, 50, 100, 200, and 400 µg/mL were added to 0.4 mL of diluted human RBC suspension. All of the samples were prepared in triplicate, and the suspension was shaken before incubation at 37 °C for 4 h. The sample was then centrifuged (Eppendorf Centrifuge 5417R) at 5400 ×g for 5 min to remove non-lysed human RBCs. Afterwards, 100 µL of the supernatant from the sample tube was moved to a 96-well plate. The supernatant was used, and hemoglobin release was assessed at 545 nm. Sodium dodecyl sulfate (0.1%) and PBS were used to establish 100% and 0% hemolysis as the

positive and negative samples, respectively. The percentage of hemolysis was described by the following equation where + and - controls are the absorbance of the solution at 100 and 0% hemolysis. Hemolysis % = [(sample absorbance - negative control) / (positive control - negative control)] × 100

### Toxicity assay on *A. salina*

The general toxicity of the ethanolic extracts on *A. salina* was assessed<sup>[29]</sup>. *A. salina* eggs were achieved from Urmia University, the West Azerbaijan Province, Iran. The cysts were seeded in a flask containing 35 g of NaCl in 1 L of distilled water. After incubation for 36-48 h at 28 °C, the larvae hatched within 48 h. The test was performed on the larvae of brine shrimp (*A. salina* Leach.). At first, a stock solution of 10 mg of ethanolic extract of aerial and root parts was dissolved in 100 µL of DMSO and then in 900 µL of the medium to prepare the stock (10 mg/ml). It was diluted to make the concentrations ranging from 0.78125 to 10 mg/ml. Ethanolic extracts (20 µL) was added to each well of the 96-well microtiter plates containing 180 µL of RPMI-1640 to form the extract concentration ranging from 1000 µg/mL to 7.8125 µg/mL. After that, 10 nauplii per well were added to the 96-well plates and incubated at 25 °C for 24 h. Afterwards, the numbers of surviving nauplii in each well were calculated under a binocular microscope after 24 h. All experimental settings for each concentration were in triplicates. Additionally, the negative control contained only 10 nauplii and artificial sea water. The percentages of nauplii deaths were calculated by considering the number of survivors in the test and control wells. The lethality was determined by Abbott's formula: Lethality (%) = [(Test - Control) / Control] × 100.

### Oral acute toxicity

An oral acute toxicity was performed to calculate the LD<sub>50</sub>. Ten Swiss Albino mice, obtained from the Pasteur Institute of Iran (25-35 g), were selected equally from both sexes. For the adaptation of the mice with laboratory conditions, every five mice were kept in a cage for seven days prior to testing and had free access to food and water according to OECD Guidelines with some modifications<sup>[30]</sup>. Different doses (control, 250, 500, 1000, and 2000 mg/kg) of the alcoholic extracts of *P. major* (various parts) were orally administered to each animal. If all animals were survived after 24 h, two additional mice were selected and treated at the highest dose (2000 mg/kg). If these two mice survived, then the LD<sub>50</sub> was more than the limited dose, and the test was stopped. All tested animals were weighed before treatment and 24 h and one week after that.

### GC-MS analysis

GC-MS of the methanolic extracts (due to the better solubility of the compounds in methanol than ethanol) and acetonic extracts of *P. major* aerial and root parts was used for analysis. GC-MS analysis was carried out by Agilent technologies 5975c, USA. Next, 1 µL of the methanolic extract was subjected to the GC-MS system equipped with a capillary column (30 m × 250 µm × 0.25 µm, Agilent). Helium was used at the flow rate of 1.0 ml/min. The injector and the interface temperature were kept at 350 °C. The column temperature was attuned as follows: the initial temperature was 50 °C (2 min) then increased at a rate of 4 °C/min up to 230 °C (2 min). The identification of the components was determined by comparing mass spectral fragmentation patterns in MS data libraries (NIST08.L)<sup>[31]</sup>.

### Statistical analysis

The experiments were directed in triplicate, and group-wise comparison and statistical analysis of the results were performed by ANOVA and Duncan's new multiple range test. SPSS v21 was used for statistical analyses. *p* < 0.05 was considered as statistically significant. The IC<sub>50</sub> values were analyzed with ED50plus v1.0 software.

### Ethical statement

The above-mentioned sampling/treatment protocols were approved by the Research Ethics Committee of University of Zanjan, Zanjan, Iran (ethical code: 21699).

## RESULTS

### Cytotoxicity activity

#### *Cell proliferation inhibition activity of different extracts of the P. major aerial parts*

Based on the results, the alcoholic extracts of *P. major* represented more antiproliferative properties on the HCT-116 in comparison with SW-480 cell line (Fig. 1). Also, the acetonic extract of aerial parts had the most inhibition effect (59%, 43%, and 27% and 37%, 29%, and 23%, respectively) on the viability of HCT-116 cells and normal cell lines at the highest concentration (400 µg/mL) in 24, 48, and 72 h, (Fig. 2A, 2B, and 2C).


Despite the lower cytotoxicity of alcoholic and acetonic extracts on SW-480 cells, compared to HCT-116 cells, the alcoholic extracts showed 92-94% viability at the concentration of 400 µg/mL in 24 h, while acetonic extract had no effect at the same time and at 48 and 72 h. Besides, the acetonic extract had the same activity with alcoholic extracts on SW-480

cells. In both cell lines, the ethanolic extract had a better cytotoxicity effect than methanolic extract (Fig. D-2F and 2G-2I). This proliferation inhibition activity of the cells was not only time-dependent but also dose-dependent. The cytotoxic effect of the alcoholic extracts in comparison to acetic extract on HEK-293 normal cells at 72 h showed that the alcoholic extracts had low inhibitory effects (between 2% to 13%) only at the concentrations of 200 and 400  $\mu\text{g/mL}$ , whereas the acetic extract revealed a significant cytotoxicity effect on the HEK-293 cells in all tested concentrations (25-400  $\mu\text{g/mL}$ ). Indeed, these results indicated that the alcoholic extracts of aerial parts of *P. major* had significant cytotoxic activity on cancer cell lines, whereas at high concentration, it showed a partial cell proliferation inhibition activity on normal cells. However, the acetic extract possessed high cytotoxicity activity on both cancer and normal cell lines; therefore, based on the results, the use of acetic extract is not recommended for the treatment of colorectal cancer.

$\text{IC}_{50}$  values of ethanolic, methanolic and acetic extracts were calculated as follows: 475.20, 655.09, and 221.64  $\mu\text{g/mL}$  for HCT-116 cells, 646.06, 756.38, and 715.28  $\mu\text{g/mL}$  for SW-480 cells, and 904.98, 1016.55, and 107.85  $\mu\text{g/mL}$  for HEK-293 cell line at 72 h, respectively (Table 1). The lowest  $\text{IC}_{50}$  was attributed to the effect of the acetic extract on HCT-116 cells (221.64  $\mu\text{g/mL}$ ) and HEK-293 cells (107.85  $\mu\text{g/mL}$ ).


#### Cell proliferation inhibition activity of root part extracts of the *P. major*

The *P. major* (ethanolic, methanolic and acetic) root extracts had more cytotoxic activity on the HCT-116 than SW-480 cells, similar to the aerial parts extracts (Fig. 3A-3C); however, the *P. major* root extracts had more antiproliferative activity than aerial parts extract on SW-480 cells (Fig. 3D-3F). Although the methanolic extract was found to have less antiproliferative properties than the ethanolic extract of *P. major* root part, the root methanolic extract displayed a more cytotoxic effect on HCT-116 than aerial parts extracts (Fig. 3G-3I). The inhibitory effect of acetic extracts *P. major* root on the viability of HEK-293 normal cells was considerable (65%). This effect on root extracts was less than the extracts obtained from aerial parts (76%) at 72 h, despite its greater cytotoxic effect on cancer cell lines.  $\text{IC}_{50}$  values of ethanolic, methanolic and acetic extracts were 405, 470, and 82  $\mu\text{g/mL}$  for HCT-116 cells, 513, 687, and 698  $\mu\text{g/mL}$  for SW-480 cells, and 948, 1563, and 125  $\mu\text{g/mL}$  for HEK-293 cells at 72 h, respectively (Table 1). The lowest  $\text{IC}_{50}$  was related to the effect of the acetic extract on HCT-116 cells and HEK-293 cells. Since the  $\text{IC}_{50}$  values of the ethanolic extract on HCT-116 and SW-480 cell lines (405 and 513  $\mu\text{g/mL}$ ) were less than a normal cell (948  $\mu\text{g/mL}$ ), this extract possessed valuable characteristics. The  $\text{IC}_{50}$  values of the alcoholic and acetic extracts of *P. major* root on cancer cell lines were lower than the aerial parts extracts.



**Fig. 1.** The growth inhibitory effect of acetic extract of *P. major* aerial parts on the HCT-116 cancer cell line. (A) Cells were seeded at a density of  $7 \times 10^3$  cells per well in a 96-well plate. The morphology of cells after treatment with aerial parts acetic extracts (B) one, (C) two, and (D and E) three days after treatment.





**Fig. 2.** The result of MTT assay, one, two and three days after treatments. The percentage of the viability of the colorectal carcinoma cell lines (HCT-116 and SW-480) and embryonic kidney normal cell line (HEK-293) were treated with different extracts of aerial parts of *P. major* at 24, 48, and 72 h, respectively. The acetonic (A, B, and C), ethanolic (D, E, and F), and ethanolic (G, H, and I) extracts. Values represent the mean of three replications  $\pm$  standard deviation.

### The biocompatibility assay of the ethanolic extracts (hemolysis toxicity)

To verify the biocompatibility of the ethanolic extracts of *P. major* aerial and root parts, we examined the hemolytic activity. Hemolysis degree for both the aerial and root parts of *P. major* was determined to be lower than 1% at the highest concentration (400  $\mu\text{g/mL}$ ) for 4 h (Fig. 4).

### Toxicity assay of ethanolic extracts on *A. salina*

The general toxicity of aerial and root parts of *P. major* ethanolic extracts was assessed against *A. salina*. The percentage of lethality was used as a bioassay indicator for the toxicity of ethanolic extracts. At all concentrations of ethanolic extracts ranging from 7.8125 to 1000  $\mu\text{g/mL}$ , all of the nauplii were alive, and no toxicity was observed (Fig. 5).

### Oral acute toxicity assay of the alcoholic extract ( $\text{LD}_{50}$ )

Mice were treated with a range of oral doses. Acute toxicity is known as an *in vivo* biocompatibility test.

One week after the oral administration of *P. major* aerial and root parts extracts, none of the animals were died. According to OCED and Hodge and Sterner Scale, we can approve that both the aerial and root extracts of *P. major* were practically safe and non-toxic. Furthermore, the change in the body weight is an important factor and an indicator of the potential toxicity (Table 2). The weight of mice after 24 and 168 hours (one week) faintly increased, indicating that all the mice had a natural activity with normal behavior.


### Determination of the composition of alcoholic and acetonic extracts by GC-MS

The composition of different extracts of *P. major* was evaluated by GC-MS, and its components were identified by the NIST08.L library. The extracts of *P. major* showed the presence of fatty acids, phenols, terpenoids, amines, amides, siloxanes, esters, alkanes, aldehydes, benzene derivatives etc. The presence of the volatile components were carried out by GC-MS, which detected acetol (5.03%), elaidic acid (5.48%),

**Table 1.** IC<sub>50</sub> estimations of various extracts on HCT-116, SW-80, and HEK-293 cell lines at different exposure times

Cell/part	24 h			48 h			72 h		
	Methanolic	Ethanollic	Acetonic	Methanolic	Ethanollic	Acetonic	Methanolic	Ethanollic	Acetonic
HCT-116									
Aerial	1391.12 ± 3.1	930.87 ± 2.3	436.76 ± 2.5	1170.76 ± 1.8	698.60 ± 3.1	307.34 ± 4.5	655.09 ± 1.9	475.20 ± 3.0	221.64 ± 2.2
Root	1142.78 ± 1.7	897.14 ± 4.1	229.91 ± 2.4	655.09 ± 4.2	897.46 ± 3.6	220.12 ± 2.8	470.16 ± 2.5	405.59 ± 3.4	82.26 ± 2.7
SW-480									
Aerial	4065.77 ± 2.3	2877.36 ± 2.4	3876.00 ± 4.8	2822.86 ± 2.2	990.24 ± 3.0	1355.71 ± 3.4	756.38 ± 2.6	646.06 ± 3.5	715.283 ± 3.4
Root	3152.87 ± 2.4	1173.53 ± 3.1	1054.53 ± 1.4	1275.02 ± 5.1	713.23 ± 1.3	901.53 ± 1.2	687.12 ± 3.3	513.08 ± 3.4	698.19 ± 1.8
HEK-293									
Aerial	3877.36 ± 9.2	3353.17 ± 2.3	204.70 ± 3.7	2366.70 ± 4.1	2381.27 ± 1.4	107.82 ± 2.5	1016.55 ± 3.8	904.98 ± 2.6	107.85 ± 2.7
Root	8824.49 ± 3.6	4026.77 ± 5.8	246.42 ± 1.9	4202.72 ± 4.2	1443.96 ± 2.4	184.76 ± 4.3	1563.04 ± 1.3	948.15 ± 3.9	125.89 ± 1.5


ED<sub>50</sub> plus V1.0 gives an estimated value of IC<sub>50</sub>; Mean of three replications ± SD at 24, 48, and 72 h.



**Fig. 3.** The result of MTT assay, one, two and three days after treatments. The percent of the viability of the colorectal carcinoma cell lines (HCT-116 and SW-480) and embryonic kidney normal cell line (HEK-293) were treated with different extracts of the root part of *P. major* at 24, 48, and 72 h, respectively. The acetonic (A, B, and C), ethanolic (D, E, and F), and methanolic (G, H, I) extracts. Values represent the mean of three replications  $\pm$  standard deviation.

octacosane (6.12%), octadecanoic acid (8.61%), and palmitic acid (15.18%) compounds in the analysis of the methanolic extract of the *P. major* aerial part. However, in the acetonic extracts, it found Bis(2-ethylhexyl) phthalate (3.67%), 1,2-Benzenedicarboxylic acid, and mono(2-ethylhexyl) ester (88.07%). GC-MS also detected octasiloxane, 1,1,3,3,5,5,7,7,9,9,11,11,13,13, 15,15-hexadecamethyl- (4.97%), cyclohexasiloxane, dodecamethyl- (6.35%), verbenone (6.96%), isoborneol (8.68%), tetradecamethylcycloheptasiloxane (9.74%), and n-hexadecanoic acid (13.8%) in the methanolic extract of *P. major* root part and heneicosane (5.23%), cis-9-hexadecenoic acid (7.99%), n-hexadecanoic acid (9.88%), 13-Docosen-1-ol, (Z)- (15.91%) 1,2-benzenedicarboxylic acid, and mono(2-ethylhexyl) ester (40.63%) in the acetonic extracts (as the dominant constituents). A list of compounds identified by GC-MS analysis is represented in Supplementary Table 3. In the present study, the common compounds in the methanolic extracts of both parts of the plant were benzaldehyde, 2-nitro-,diaminomethylidenehydrazone (0.25% and 1.40%), pentadecanoic acid, 13-methyl-,methyl ester (0.36%

and 1.76%), hexadecamethylcyclo-octasiloxane (0.32% and 4.51%), and hexadecanoic acid (15.17% and 13.79%); however, in the acetonic extracts, the most frequent compounds were 1,2-benzenedicarboxylic acid, mono(2-ethylhexyl) ester (88.07% and 40.63%), 2-pentanone, 4-hydroxy-4-methyl- (0.12%, and 1.29%), and Oxirane,heptadecyl- (1.01% and 1.98%), respectively. Both cyclohexasiloxane



**Fig. 4.** Hemolytic assay to verify the biocompatibility of ethanolic extracts of *P. major* aerial and root parts.

dodecamethyl-/CAS number: 540-97-6 (1.71%, 6.35%, 0.89%, and 0.47%) and cycloheptasiloxane, tetradecamethyl-/CAS number: 107-50-6 (1.16%, 9.74%, 1.02%, and 1.68%) compounds were cyclic methyl siloxanes and found in both the methanolic and acetonic extracts of aerial and root parts of the plant with different percentage, respectively.

## DISCUSSION

*P. major* is a medicinal plant with various therapeutic benefits. *P. major* extract can inhibit the cell proliferation of melanoma, renal and mammae<sup>[32]</sup>, skin<sup>[10]</sup>, leukemia<sup>[33]</sup>, breast adenocarcinoma, and melanoma (UACC-62) cell lines<sup>[16]</sup>. The *in vitro* cytotoxic activity of the methanolic extract of *P. major* has been evaluated on HCT-15, SQC-UIISO, OVCAR, and KB<sup>[21]</sup>, which demonstrated that *P. major* has hematopoietic activity. A comprehensive literature review on *P. major* extract, as an antineoplastic agent, revealed that this plant is an efficient chemoprophylactic and antimetastatic agent against several malignancies, such as breast cancer, hepatoma, and Ehrlich ascites tumor<sup>[23,34-36]</sup>. Based on the research conducted in 2013, *P. major* extract displayed a proapoptotic impact on the hypergastrinemia rat (*Rattus norvegicus*) model. This effect has been attributed to the increased level of caspase 3<sup>[37]</sup>. Cytotoxicity potential of methanolic extracts from leaves of *P. major* were evaluated against three human cancer cell lines. The results indicated that *P. major* extracts have cell proliferation inhibition activity on breast adenocarcinoma and melanoma cell lines recommended by the National Cancer Institute (USA)<sup>[16]</sup>.

Considering the aforementioned properties of *P. major*, in the first phase of the current study, the cytotoxic

effects of the different extracts of *P. major* was examined by MTT test on HCT-116, SW-480, and HEK-293. The ethanolic, methanolic and acetonic extracts of *P. major* exhibited a significant cytotoxic activity on colorectal carcinoma cell lines. Based on the obtained results, the ethanolic extract was more cytotoxic compared to the methanolic extract and did not show any significant cytotoxicity effect on the normal cell line, except for in the concentrations of 200 and 400 µg/mL. In this regard, the alcoholic extracts derived from the root parts of *P. major* might be deliberated as a valuable source of metabolites with potential uses as antitumor drug precursors. The US National Cancer Institute argues that the IC<sub>50</sub> value should be below 30 µg/mL so that a crude extract can serve as an appropriate agent for further refinement<sup>[2]</sup>. The root extracts of *P. major* showed more cytotoxicity than the aerial parts (leaf and stem) extracts and had a lower and more valuable IC<sub>50</sub> index at 72 h. Indeed, these results indicated that the alcoholic extracts of aerial parts of *P. major* had significant cytotoxic activity on cancer cell lines, whereas at high concentration, it showed a partial cell proliferation inhibition activity on normal cells. However, the acetonic extract possessed high cytotoxicity activity on both cancer and normal cell lines; therefore, using acetonic extract is not recommended for the treatment of colorectal cancer. Since the IC<sub>50</sub> value of the ethanolic extract on HCT-116 and SW-480 cell lines (475.20 and 646.06 µg/mL) was less than a normal cell (904.98 µg/mL); hence, it can be considered as valuable and useful extracts for medicinal treatment.

In the second phase of the current study, aerial and root parts of *P. major* extracts were evaluated by hemolysis assay, general toxicity assay on *A. salina* and oral acute toxicity study (LD<sub>50</sub>) on adult Swiss Albino mice. The study of Atta *et al.*<sup>[38]</sup> has reported




Fig. 5. Toxicity of ethanolic extracts of *P. major* on *A. salina*



**Table 2.** Weight change and Mortality rates of mice after 24 hours and one week, submitted to various doses of *P. major*

Groups	Dose (mg/kg)	No. of animals	Mean weight of animals (24 h)	Duration of study (one week)	Dead rate (%)
Control	-	5	30 ± 1.19	34 ± 1.11	0
	175	5	35 ± 1.10	36 ± 1.27	0
	5000	5	29 ± 1.58	31 ± 1.30	0
M. A. M	17.5	5	30 ± 1.21	31 ± 1.27	0
	175	5	35 ± 1.64	34 ± 1.15	0
	5000	5	25 ± 0.70	28 ± 1.33	0
M. A. E	17.5	5	33 ± 2.05	33 ± 1.58	0
	175	5	35 ± 1.42	37 ± 1.58	0
	5000	5	25 ± 1.45	26 ± 0.79	0
M. R.M	17.5	5	35 ± 1.33	34 ± 2.57	0
	175	5	36 ± 2.91	37 ± 0.79	0
	5000	5	24 ± 1.27	24 ± 0.70	0
M. R. E	17.5	5	36 ± 1.27	34 ± 4.69	0
	175	5	36 ± 1.27	34 ± 4.69	0
	5000	5	24 ± 1.27	24 ± 0.70	0

Value is mean ± standard deviation. M.A.M and M.A.E., *P. major* aerial parts methanolic and ethanolic extracts, respectively; M.R.M. and M.R. E., *P. major* root methanolic and ethanolic extracts, respectively.

that the oral administration of *P. major* seed methanol extracts in doses up to 2.5 g/kg body weight did not cause any major signs of acute toxicity, and no deaths were reported up to 72 h after the oral administration<sup>[38]</sup>. In this study, at all the concentrations of extracts, we observed no toxicity. Hence, it confirms the safety of these extracts and shows that these extracts are practically nontoxic. Mirzaei *et al.*<sup>[39]</sup> selected the *P. major* to investigate its toxicity against both *A. salina* and *A. uramiana*. The result showed a positive correlation between the data obtained from the two aforesaid species, and the LC<sub>50</sub> of the *P. major* was 303.7 µg/ml for the methanolic extract, whereas the thymol standard possessed the LC<sub>50</sub> value of 7.2 µg/ml.

Indeed, it can be concluded that this extract did not exhibit high toxicity in comparison with the thymol. Thus, this medicinal plant can be considered as a safe and non-toxic agent<sup>[39]</sup>. This result showed that the high concentration of both aerial and root parts of *P. major* did not affect the membrane integrity of RBCs and both *P. major* aerial and root extracts are biocompatible. It is interesting to note that the hemolytic activity of *P. major* has not been reported so far, and the obtained data indicated that this medicinal plant is practically nonhemolytic since its hemolysis value is below 1%. It also been reported that when the hemolysis percentage does not exceed 10%, it can serve as a non-toxic agent<sup>[40]</sup>. Moreover, we performed GC-MS to investigate the components of the extracts

of aerial and root parts of *P. major*. The GC-MS analysis of extracts from *P. major* revealed the existence of fatty acids, terpenoids, siloxanes, and other medicinal compounds. Therefore, the extracts can be used as anticancer drugs against tumor growth and propagation because *P. major* has effective anticancer compounds. Some of the compounds identified in our study of analysis of the *P. major* extracts include Silanediol, dimethyl-<sup>[41]</sup>, camphor<sup>[42]</sup>, Borneol<sup>[43]</sup>, Isoborneol/Isocamphol<sup>[44]</sup>, 1,2-Benzenedicarboxylic acid, and mono(2-ethylhexyl) ester with antiviral features<sup>[45]</sup>. Two antimalarial compounds, i.e. 2,4-Di-tert-butylphenol<sup>[46]</sup> and n-hexadecanoic acid<sup>[47]</sup>, were also found in the root extracts. Nortriptyline is a commercially available compound that has anti-depressant properties<sup>[48]</sup>; however, most of the compounds reported in the extracts have antibacterial, antifungal, anticancer, antioxidant and anti-inflammatory properties and are fatty acid. Fatty acids, such as oleic acid, linoleic acid, and palmitic acid, are important sources with antimicrobial and antifungal activities<sup>[18]</sup>. The antimicrobial properties of *P. major* are attributed to the presence of camphor, bornyl acetate, and borneol compounds; the biological activities of these compounds have previously been investigated<sup>[49]</sup>.

Our findings support the notion that the extracts of *P. major* might contain a variety of secondary metabolites that represent the multi-biological activities, which can be applied for the development of antitumor drug

precursors. The purification of these bioactive compounds is thought to be useful for the formulation of therapeutic agents against cancer. Some of the compounds identified in this study, including, 2,4-Di-tert-butylphenol, camphor, gentisic acid, isoborneol, linoleic acid, methyl ester/linoleic acid ester, myristic acid, methyl sterate, oleic acid, palmitic acid, p-cymene, stearophanic acid, trans-anethole etc., are therapeutically important.

### ACKNOWLEDGEMENTS

The study was financially supported by University of Zanjan, Zanjan, Iran. The authors would like to thank the authority of Department of Pharmaceutical Biotechnology, School of Pharmacy, Zanjan University of Medical Sciences, Zanjan, Iran for providing laboratory equipment.

**CONFLICT OF INTEREST.** None declared.

### REFERENCES

- Powis G. Dose-dependent metabolism, therapeutic effect, and toxicity of anticancer drugs in man. *Drug metabolism reviews* 1983; **14**(6): 1145-1163.
- Rajkumar V, Gunjan G, Ashok Kumar R, Lazar M. Evaluation of cytotoxic potential of *Acorus calamus* rhizome. *Ethnobotanical leaflets* 2009; **7**: Article 2.
- Gali-Muhtasib H, Bakkar N. Modulating cell cycle: current applications and prospects for future drug development. *Current cancer drug targets* 2002; **2**(4): 309-336.
- Wang H, Zhao C, Huang Y, Wang F, Li Y. Chemical constituents and bioactivities of plantaginis herba. *Hong Kong medicinal journal* 2015; **22**: 29-35.
- Najafian Y, Hamed SS, Farshchi MK, Feyzabadi Z. *Plantago major* in traditional persian medicine and modern phytotherapy: a narrative review. *Electronic physician* 2018; **10**(2): 6390-6399.
- Radu N, Ghita I, Coman O, Rau I. Therapeutic effect of flavonoids derived from *Plantago* species. *Molecular crystals and liquid crystals* 2010; **523**(1): 273-281.
- Radu N, Ghita I, Rau I. Therapeutic effect of polysaccharides from *Plantago* species. *molecular crystals liquid crystals* 2010; **523**(1): 236-246.
- Dalar A, Türker M, Konczak I. Antioxidant capacity and phenolic constituents of *Malva neglecta* Wallr. and *Plantago lanceolata* L. from Eastern Anatolia Region of Turkey. *Journal of herbal medicine* 2012; **2**: 42-51.
- Afifi M, Salama O, Hassan M, Naeim Z, Mohammed K. Phenolic constituents of *Plantago albicans* L. *Journal of pharmaceutical sciences* 2000; **16**(2): 204-212.
- Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. *Journal of ethnopharmacology* 2000; **71**(1-2): 1-21.
- Rønsted N, Göbel E, Franzyk H, Jensen SR, Olsen CE. Chemotaxonomy of *Plantago*. Iridoid glucosides and caffeoyl phenylethanoid glycosides. *Phytochemistry* 2000; **55**(4): 337-348.
- Taskova R, Evstatieva L, Handjieva N, Popov S. Iridoid patterns of genus *Plantago* L. and their systematic significance. *Zeitschrift für naturforschung C*. 2002; **57**(1-2): 42-50.
- Janković T, Menković N, Zdunić G, Beara I, Balog K, et al. Quantitative determination of aucubin in seven *Plantago* species using HPLC, HPTLC, and LC-ESI-MS methods. *Analytical letters* 2010; **43**(16): 2487-2495.
- Akihisa T, Yasukawa K, Tokuda H. Potentially cancer chemopreventive and anti-inflammatory terpenoids from natural sources. *Studies in natural products chemistry* 2003; **29**: 73-126.
- Hertog MG, Hollman PC, Katan MB, Kromhout D. Intake of potentially anticarcinogenic flavonoids and their determinants in adults in The Netherlands. *Nutrition and cancer* 1993; **20**: 21-29.
- Gálvez M, Martín-Cordero C, López-Lazaro M, Cortés F, Ayuso MJ. Cytotoxic effect of *Plantago* spp. on cancer cell lines. *Journal of ethnopharmacology* 2003; **88**(2-3): 125-130.
- Sugiyarto AD, Dwisetyawani A, Pitoyo A. Estimasi kemulim pahan dan distribusi *Plantago major* L. di Gunung Lawu. *Biodiversitas* 2006; **7**(2): 143-146.
- Krishnaveni M, Dhanalakshmi R, Nandhini N. GC-MS analysis of phytochemicals, fatty acid profile, antimicrobial activity of *Gossypium* seeds. *International journal of pharmaceutical sciences review and research* 2014; **27**(1): 273-276.
- Holetz FB, Pessini GL, Sanches NR, Cortez DAG, Nakamura CV, Dias Filho BP. Screening of some plants used in the Brazilian folk medicine for the treatment of infectious diseases. *Memórias do Instituto Oswaldo Cruz* 2002; **97**(7): 1027-1031.
- Chiang LC, Chiang W, Chang MY, Lin CC. In vitro cytotoxic, antiviral and immunomodulatory effects of *Plantago major* and *Plantago asiatica*. *American journal chinese medicine* 2003; **31**(2): 225-234.
- Velasco-Lezama R, Tapia-Aguilar R, Román-Ramos R, Vega-Avila E, Pérez-Gutiérrez MS. Effect of *Plantago major* on cell proliferation *in vitro*. *Journal of ethnopharmacology* 2006; **103**(1): 36-42.
- Shipochliev T. Uterotonic action of extracts from a group of medicinal plants. *Veterinarno meditsinski nauki* 1981; **18**(4): 94-98.
- Lithander A. Intracellular fluid of waybread (*Plantago major*) as a prophylactic for mammary cancer in mice. *Tumor biology* 1992; **13**(3): 138-141.
- Hetland G, Samuelsen AB, Løvik M, Paulsen BS, Aaberge IS, Groeng EC, Michaelsen TE. Protective effect of *Plantago major* L. Pectin polysaccharide against systemic *Streptococcus pneumoniae* infection in mice. *Scandinavian journal immunology* 2000; **52**(4): 348-355.
- Ozaslan M, Didem Karagöz I, Kalender ME, Kilic IH, Sari I, Karagöz A. *In vivo* antitumoral effect of *Plantago*

- major* L. extract on BALB/C mouse with Ehrlich ascites tumor. *American journal of chinese medicine* 2007; **35**(5): 841-851.
26. Rahamooz-Haghighi S, Asadi MH. Anti-proliferative effect of the extracts and essential oil of *Pimpinella anisum* on gastric cancer cells. *Journal of herbmed pharmacology* 2016; **5**(4): 157-161.
  27. Plumb JA. Cell Sensitivity Assays. In: Langdon SP eds. *Cancer Cell Culture. Methods in Molecular Medicine™*. Vol 88. Humana Press. Retrieved from: <https://doi.org/10.1385/1-59259-406-9:165>.
  28. Aghajanzadeh M, Zamani M, Rashidzadeh H, Rostamizadeh K, Sharafi A, Danafar H. Amphiphilic Y shaped miktoarm star copolymer for anticancer hydrophobic and hydrophilic drugs codelivery: Synthesis, characterization, in vitro, and in vivo biocompatibility study. *Journal of biomedical materials research part a* 2018; **106**(11): 2817-2826.
  29. Rajabi S, Ramazani A, Hamidi M, Naji T. *Artemia salina* as a model organism in toxicity assessment of nanoparticles. *Daru* 2015; **23**(1): 20.
  30. OECD guidelines for the testing of chemicals acute oral toxicity- up and down procedure (UDP). Retrieved from: <https://www.oecd-ilibrary.org/docserver/9789264071049-en.pdf?expires=1611038404&id=id&accname=guest&checksum=4BD111B77C5C007B934ABFE2B1B45663>
  31. Jamilah J, Sharifa ASA, Sharifah NR. GC-MS analysis of various extracts from leaf of *Plantago major* used as traditional medicine. *World applied sciences journal* 2012; **17**: 67-70.
  32. Bashford JN, Norwood J, Chapman SR. Why are patients prescribed proton pump inhibitors? Retrospective analysis of link between morbidity and prescribing in the general practice research database. *British medical journal* 1998; **317** (7156): 452-456.
  33. Ikawati Z, Wahyuono S, Maeyama K. Screening of several Indonesian medicinal plants for their inhibitory effect on histamine release from RBL-2H3 cells. *Journal of ethnopharmacology* 2001; **75**(2-3): 249-256.
  34. Yeşilada E, Sezik E, Fujita T, Tanaka S, Tabata M. Screening of some Turkish medicinal plants for their antiulcerogenic activities. *Phytotherapy research* 1993; **7**(3): 263-265.
  35. Chiang LC, Ng LT, Chiang W, Chang MY, Lin CC. Immunomodulatory activities of flavonoids, monoterpenoids, triterpenoids, iridoid glycosides and phenolic compounds of *Plantago* species. *Planta medica* 2003; **69**(07): 600-604.
  36. Kobeasy I, Abdel-Fatah M, El-Salam SMA, Mohamed ZEOM. Biochemical studies on *Plantago major* L. and *Cyamopsis tetragonoloba* L. *International journal of biodiversity and conservation* 2011; **3**(3): 83-91.
  37. Sutrisna E, Maskoen AM, Sujatno M, Sastramihardja HS. Potential apoptotic effect of Plantain extract (*Plantago major* L.) through increasing of caspase-3 level on hypergastrinemic rat model. *International journal of research in pharmaceutical sciences and nanotechnology* 2013; **2**(3): 371-381.
  38. Atta AH, Soad N, Samar M. Potential protective effect of some plant extracts against carbon tetrachloride-induced hepatotoxicity. *African journal of traditional complementary alternative medicines* 2006; **3**(3): 1-9.
  39. Mirzaei M, Mirzaei A. Comparison of the *Artemia salina* and *Artemia uramiana* bioassays for toxicity of 4 Iranian medicinal plants. *International research journal of biological sciences* 2013; **2**(3): 49-54.
  40. Amin K, Dannenfelser RM. In vitro hemolysis: guidance for the pharmaceutical scientist. *Journal of pharmaceutical sciences* 2006; **95**(6): 1173-1176.
  41. Femi-Adepojua AG, Adepojub AO, Durodolac FA, Abiola AGM. Phytochemical, antimicrobial and bio-active component analysis of *Platycerium superbum* (L.) methanolic extract. *International journal of sciences: basic and applied research* 2018; **40**: 98-107.
  42. Chen W, Vermaak I, Viljoen A. Camphor—a fumigant during the black death and a coveted fragrant wood in ancient Egypt and Babylon—a review. *Molecules* 2013; **18**(5): 5434-5454.
  43. Šála M, De Palma AM, Hřebabecský H, Nencka R, Dračinský M, Leyssen P, Neyts J, Holý A. Design, synthesis, and biological evaluation of novel coxsackievirus B3 inhibitors. *Bioorganic and medicinal chemistry* 2010; **18**(12): 4374-4384.
  44. Armaka M, Papanikolaou E, Sivropoulou A, Arsenakis M. Antiviral properties of isoborneol, a potent inhibitor of *Herpes simplex virus* type 1. *Antiviral research* 1999; **43**(2): 79-92.
  45. Govindappa M, Prathap S, Vinay V, Channabasava R. Chemical composition of methanol extract of endophytic fungi, *Alternaria* sp. of *Tebebuia argentea* and their antimicrobial and antioxidant activity. *International journal of biological and pharmaceutical research* 2014; **5**(11): 861-869.
  46. Kusch P, Deininger S, Specht S, Maniako R, Haubrich S, Pommerening T, Lin PKT, Hoerauf A, Kaiser A. In vitro and in vivo antimalarial activity assays of seeds from *Balanites aegyptiaca*: compounds of the extract show growth inhibition and activity against plasmodial aminopeptidase. *Journal of parasitology research* 2011; **2011**: 368692.
  47. Ahmed D. Biological activities of characterized isolates of n-hexane extract of *Azadirachta indica* A. Juss (Neem) leaves. *Nature and science* 2013; **11**(5): 141-147.
  48. Nebes RD, Pollock BG, Houck PR, Butters MA, Mulsant BH, Zmuda MD, Reynolds 3<sup>rd</sup> CF. Persistence of cognitive impairment in geriatric patients following antidepressant treatment: a randomized, double-blind clinical trial with nortriptyline and paroxetine. *Journal of psychiatric research* 2003; **37**(2): 99-108.
  49. Benali T, Habbadi K, Abdelmajid K, Marmouzi I, Zengin G, Bouyahya A, Chamkhi I, Chtibi H, Tarik A, Achbani E, Hammani K. GC-MS analysis, antioxidant and antimicrobial activities of *Achillea Odorata* Subsp. *Pectinata* and *ruta montana* essential oils and their potential use as food preservatives. *Foods* 2020; **9**(5): 668.