

Transient Expression Assay of $\Lambda\gamma$ -588 (A/G) Mutations in the K562 Cell Line

Mohammad Hamid¹, Frouzandeh Mahjoubi², Mohammad Taghi Akbari³, Hossein Khanahmad⁴, Fatemeh Jamshidi¹, Sirous Zeinali¹ and Morteza Karimipour^{*1}

¹Molecular Medicine Division, Biotechnology Research Center, Pasteur Institute of Iran, Tehran;

²Dept. of Clinical Genetics, National Institute of Genetic Engineering and Biotechnology (NIGEB), Tehran;

³Dept. of Medical Genetics, Faculty of Medical Sciences, Tarbiat Modarres University, Tehran;

⁴BCG Research Center, Pasteur Institute of Iran, Karaj, Iran

Received 10 October 2010; revised 6 March 2011; accepted 8 March 2011

ABSTRACT

Background: In the previous study, we have shown that the presence of A allele at position -588 in $\Lambda\gamma$ -globin gene was highly frequent and closely associated with fetal hemoglobin elevation among β -thalassemia intermedia patients. Therefore, we decided to investigate whether this allele (A allele at -588) could result in an increase in $\Lambda\gamma$ -globin gene expression to ameliorate the severity of the disease in thalassemia patients.

Methods: Three constructs containing μ locus control region, $\Lambda\gamma$ -globin and β -globin genes were designed and employed in the transient expression assay. The difference among constructs was in the promoter region of $\Lambda\gamma$ -globin gene (A and G alleles at -588). A construct with T to C base substitution at -175 of $\Lambda\gamma$ -globin, created by site-directed mutagenesis, was selected as positive control. The K562 cell line was transfected with the above constructs. Subsequently, the expression of $\Lambda\gamma$ -globin gene was determined by quantitative real-time reverse transcription-PCR. **Results:** There was not a significant increase in the expression of $\Lambda\gamma$ -globin gene in the construct containing A allele comparing the one with G allele at -588. **Conclusions:** -588 (A>G) mutation does not play a major role in regulation of $\Lambda\gamma$ -globin gene, suggesting that other factors may be involved. *Iran. Biomed. J. 15 (1 & 2): 15-21, 2011*

Keywords: β -thalassemia, $\Lambda\gamma$ -globin, K562 cells

INTRODUCTION

The human β -globin cluster contains five developmentally regulated genes (5'- ϵ -G γ -A γ - δ - β -3') on the short arm of chromosome 11. The G γ and A γ genes are normally expressed at high levels only during fetal development, followed by switching to adult (δ , β) during the perinatal period. Due to switching, fetal hemoglobin (HbF) comprises only 1% to 2% of the total hemoglobin in healthy adults [1, 2].

The genes of β -globin cluster are regulated by locus control region (LCR), which is composed of five major DNase I hypersensitive sites (HS1-HS5) located far upstream of the genes [3]. It seems the LCR activates β -globin gene expression through a direct interaction with the promoter region and make

an open chromosomal region more accessible to *trans*-acting factors [4]. Many studies have investigated the roles of individual HS in various expression assays [5, 6]. HS2 acts as a strong enhancer in both transient expression assays and transgenic mice experiments. HS3 was also shown to enhance the expression of β -globin gene. HS4 does not enhance solely, but can increase expression in cooperation with other HSs [5]. In contrast to normal switching, the individuals with hereditary persistence of fetal hemoglobin (HPFH) have constantly high levels of HbF during adult life. This elevation is mainly due to large deletions within β -globin cluster in HPFH and $\delta\beta$ -thalassemia or in some cases of non-deletional HPFH (nd-HPFH) by mutations in the promoter region of γ -globin genes [7, 8]. Several nd-HPFH mutations have been

*Corresponding Author; Tel. (+98-21) 6648 0780; Fax: (+98-21) 6648 0780; E-mail: mortezakarimi@yahoo.com

Table 1. γ -globin oligonucleotide primers used for PCR and real-time experiments.

Primer	Location	Sequence (5'>3')	Amplicon (bp)
A γ F	-883 to -860	<u>accggt</u> CACAGTACCTGCCAAAGAACATTC ¹	2788
A γ R	+1885 to +1905	<u>gcggccgc</u> AATCAATCCAGCCCCAGGTC ²	
A γ F	-883 to -860	<u>accggt</u> CACAGTACCTGCCAAAGAACATTC ¹	720
R1 site direct	-183 to -160	<u>CCGTTTCAGACAGATGTTTGCATT</u> ³	
R2 site direct	-183 to -160	AATGCAAACATCTGTCTGAAACGG ³	1840
R2 A γ R	+1633 to +1657	<u>atggcgccgc</u> ACACATACACACTTCCCTCAATATA ²	
Real A γ F		ATGTGGATCCTGAGAACTTCAAGCT	136
Real A γ R	+1451 to +1469	GACAGGGCACTGGCCACTG	

Underlined bases are related to ¹*AgeI* site, ²*NotI* site and ³underlined bases are mutated to produce -175(T>C) base substitution

reported in the promoter region of either the γ - or γ -globin genes [9]. Most of these mutations occur in transcription factor binding sites, creating the new factor binding motifs or disrupting the existing ones [9]. The regulatory sequences 5' to the γ -globin gene harbor some polymorphic markers, including -588 A>G, -521 C>G, -500 C>T, -499 T>A, -202 C>T, -198 T>C, -196 C>T, -195 C>G, -175 T>C, -117 G>A, and -114 C>T. The most important mutations are -175 T>C and -117 G>A that their functional roles have been demonstrated in transgenic mice [8, 10-13].

The level of HbF may also be influenced by some other genetic determinants linked to the β -globin gene cluster such as polymorphisms in the regulatory region of γ -globin gene. The presence or absence of the -158 C>T γ -*XmnI* polymorphism has been identified to be associated with the high level of HbF in thalassemia intermedia (TI) patients [3]. We have previously shown that the presence of A allele at -588 was highly frequent and closely associated with HbF elevation in TI patients [3].

Therefore, we decided to investigate whether this allele (A allele at -588) could result in an increase in γ -globin gene expression. For this purpose, three constructs containing μ LCR, γ -globin and β -globin genes were designed for exploiting in the transient expression assay in K562 erythroid cell line.

MATERIALS AND METHODS

Plasmid DNA constructions. The genomic DNA was obtained from TI patients with A and G alleles at -588 of γ -globin [3]. The primers A γ F and A γ R were designed to amplify the sequences from -883 to +1905 of γ -globin gene (Table 1). These primers also were engineered to contain *AgeI* (forward

primer) and *NotI* (reverse primer) at their 5' ends (Table 1) and the fragments cloned into the T/A vector (Fermentase, Hanover, MD, USA). Intact core sequences of HS4 (442 bp), HS3 (2004 bp) and HS2 (1468 bp) designated μ LCR (unpublished data) were cloned into pBGGT plasmid [14]. In order to clone γ -globin fragment, two complementary oligonucleotides containing *NotI* and *Sall* sequences with *XhoI* ends were annealed together and directly cloned into the unique *XhoI* site of the pBGGT construct. The 3.9-kb μ LCR fragment was released by *NheI* and *AgeI* from pBGGT construct and inserted into pCDNA3.1 plasmid (Invitrogen, Paisley, UK). Subsequently, the fragments of γ -globin gene (*AgeI-NotI*) with A allele at -588 and β -globin gene (*NotI-PmeI*) subcloned in pCDNA3.1 μ LCR construct, generating plasmid construct pCDNA3.1 μ LCR^{A γ -588(A)} β . PCR mutagenesis was used to generate a 2560-bp fragment of the PCR products containing the T>C mutation at position -175 in three stages. Stage 3 was modified to include primers A γ F and R2 A γ R (Table 1), engineered to contain an *AgeI* (forward primer) and *NotI* sites (reverse primer) at their 5' ends. The presence of mutation was confirmed by DNA sequencing. Similar strategies were also adopted to create constructs pCDNA3.1 μ LCR^{A γ -588(G)} β and pCDNA3.1 μ LCR^{A γ -175(C)} β (Fig. 1).

Cell growth and transfection. Human erythroleukemia K562 was cultured in DMEM (GIBCO-BRL, Grand Island, NY) supplemented with 10% fetal bovine serum (GIBCO-BRL, Grand Island, NY, USA), penicillin/streptomycin (BioSera, Ringmer, East Sussex, UK) in a humidified 5% CO₂ environment. Transient DNA transfections were performed by electroporation using a Bio-Rad Gene Pulser (Bio-Rad Laboratories, Hercules, CA, USA)




Fig. 1. Schematic representation of the expression constructs and their individual components.

and Lipofectamine™ LTX (Invitrogen Corp., Carlsbad, CA, USA) in triplicate form. Approximately, 2.5×10^6 cells of K562 were electroporated with 50 $\mu\text{g}/\text{ml}$ of the constructs and mock plasmid (pCDNA3.1). Lipofectamine™ LTX was prepared according to the manufacturer's protocol.

Efficiency of electroporation and transfection.

The rate of transfection in K562 cell line was determined by analysis of green fluorescent protein (GFP) expression of transfected pEGFP-N1 plasmid (Clontech, Palo Alto, CA, USA) by flow cytometry (Partec, Germany). Transfection efficiency was determined 48 h after transfection. Transfection and electroporation experiments were optimized based on Table 2. For electroporation, the cells were suspended in DMEM without FBS and transferred to a sterile electroporation cuvette (Bio-Rad Gene

Pulser cuvette, 0.4 cm), in the presence of 50 $\mu\text{g}/\text{ml}$ of pEGFP-N1 plasmid.

Total RNA extraction and cDNA synthesis. Total RNA was extracted from K562 cell line using Trizol reagents (Sigma- Aldrich, Germany) following the manufacturer's instructions and resuspended in 30 μl of DEPC-treated water. One microgram of total RNA was reverse transcribed with Oligo(dT) primers and Superscript III reverse transcriptase (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. The concentration of cDNA was determined by a Nanodrop ND-1000 Spectrophotometer (Nanodrop Technologies, Rockland, DE) at 260/280 nm.

Real-time PCR. Real-time reverse transcription-PCR was performed in final reaction volumes of 25 μl , consisting of 12.5 μl SYBR Green I master mix (Applied Biosystems, Warrington, UK), 500 nm of forward and reverse primers and 5 ng of cDNA. Thermal cycling was performed on the ABI 7300 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) using the following cycling conditions: 10 min at 95°C as first denaturation step, followed by 40 cycles at 95°C for 15 s and 60°C for 1 min. Each complete amplification stage was followed by a dissociation stage; at 95°C for 15 s, 60°C for 30 s and 95°C for 15 s. The samples were analyzed in triplicate for each studied construct. PCR primers (Real $\text{A}\gamma\text{F}$ and Real $\text{A}\gamma\text{R}$) were designed using the primer express software v.3.0 (Applied Biosystems, Foster City, CA, USA) to specifically amplify mRNA from the γ -globin gene (Table 1). To avoid amplification of contaminating genomic DNA, exon-exon junction forward primer was designed.

Table 2. Electroporation and transfection experiments in K562 cell line.

Transfection condition	Lipofectamine™ LTX kit			DNA ($\mu\text{g}/\text{ml}$)	Electroporation voltage (V)
	DNA (μg)	TR (μL)	PR (μL)		
1	0.2	0.50	0.2		
2	0.2	0.70	0.2	50	270
3	0.2	1.10	0.2		
4	0.5	1.25	0.5		
5	0.5	2.00	0.5	50	300
6	0.5	2.75	0.5		
7	1.0	2.50	1.0		
8	1.0	3.50	1.0	50	220
9	1.0	5.50	1.0		
10	2.5	6.25	2.5		
11	2.5	10.50	2.5	50	500
12	2.5	13.75	2.5		

DNA, pEGFP-N1 plasmid; TR, transfection reagent; PR, plus reagent

Table 3. Efficiency of transfection by Lipofectamine™ LTX kit and electroporation in K562 cell line.

Transfection condition	Lipofectamine™ LTX kit (%)	Electroporation (%)		
			Cell viability (%)	
			PBS	DMEM
1-3	5.30 ± 0.80	23.15 ± 2.1	15	68
4-6	8.74 ± 0.45	26.13 ± 2.8	16	75
7-9	5.39 ± 1.30	19.00 ± 1.9	13	78
10-12	0.65 ± 1.50	8.96 ± 3.2	17	65

Gene expression was calculated according to Livak and Schmittgen [15] using β -actin as an internal reference gene. The PCR efficiency of γ -globin gene was calculated according to the slope of the standard curve and the following equation: Efficiency = $[10^{(-1/\text{slope})}] - 1$ [16]. Statistical analysis including mean, standard deviation and correlation coefficients were carried out using SPSS for windows, version 12 (SPSS Inc., Chicago, IL, USA).

Statistical analysis. Quantitative variables were expressed as means \pm S.D. while qualitative variables were expressed as percentages. Fisher's exact test was used for comparing the qualitative variables. *P* values lower than 0.05 were considered as statistically significant.

RESULTS

Constructs. The three constructs included pCDNA3.1 μ LCR γ ^{588(G)} β , pCDNA3.1 μ LCR γ ^{588(A)} β and pCDNA3.1 μ LCR γ ^{175(T>C)} β were created and confirmed by sequencing. These constructs contain about 4 kb μ LCR, 3 kb γ -globin gene and 2 kb β -globin gene. We assumed that the μ LCR would

have a local effect on the promoters of γ - and β -globin genes [1].

Optimization of transfection. Transfection experiments by Lipofectamine™ LTX kit (Invitrogen, USA) were optimized according to the suppliers' instruction with variable amounts of plasmid DNA and the volumes of transfection reagents. For each transfection, twelve different conditions were examined, and every experiment was repeated three times. The results of transfection yield for each set of transfection are shown in Table 3. The highest mean rate of transfection by Lipofectamine™ LTX kit was 8.74 ± 0.45 (Fig. 2 and Table 3). In addition, to optimize DNA electroporation, different electroporation conditions were tested (Table 2), including variations in electric field and capacitance. The best results, represented by 26.13 ± 2.8 GFP-positive cells, were achieved in a voltage of 300 V combined with a capacitance of 950 μ F (Fig. 2). As shown in Table 3, cell viability was greatly reduced in phosphate buffered saline (PBS); therefore, we used DMEM as electroporation buffer which resulted in much higher (75%) cell viability.




Fig. 2. A flow cytometric assay to determine the percentage of GFP-positive cells in K562 cell line. (A) non-transfected negative control cell; (B) transfected cell by electroporation; (C) transfected cell by Lipofectamine™ LTX kit.

Table 4. The mRNA levels of γ -globin gene in different constructs transfected to K562 cells, compared to mock plasmid.

Constructs	Avg. ΔC_T^{γ}	Avg. ΔC_T^{β} -actin	ΔC_T	$\Delta \Delta C_T$	$2^{-\Delta \Delta C_T}$
Transfection by electroporation					
pCDNA3.1 (Mock plasmid)	20.8	21.1	-0.3	0.00 \pm 0.12	1.0 (0.9- 1.1)
pCDNA3.1 μ LCR γ -588(A) β	22.0	24.5	-2.5	-2.20 \pm 1.60	4.6 (1.5-13.9)
pCDNA3.1 μ LCR γ -588(G) β	23.4	25.8	-2.4	-2.10 \pm 0.40	4.3 (3.2-5.6)
pCDNA3.1 μ LCR γ -175(T>C) β	20.5	25.3	-4.8	-4.50 \pm 2.00	22.6 (6.9-111.4)
Transfection by lipofectamine TM LTX kit					
pCDNA3.1 (Mock plasmid)	24.4	25.4	-1.0	0.00 \pm 0.80	1.0 (0.2-1.8)
pCDNA3.1 μ LCR γ -588(A) β	24.3	26.5	-2.2	-1.20 \pm 1.00	2.3 (1.3-4.6)
pCDNA3.1 μ LCR γ -588(G) β	24.0	26.0	-2.0	-1.00 \pm 0.40	2.0 (1.5-2.6)
pCDNA3.1 μ LCR γ -175(T>C) β	20.8	25.0	-4.2	-3.20 \pm 2.20	9.2 (2-42.2)

Expression studies in K562 cells. The K562 cells were transfected with the created constructs by determined LipofectamineTM LTX kit transfection conditions and electroporated with 300 V and 950 μ F in triplicate form. All transfections were performed at least in triplicate and repeated several times. To determine the relative amount of the γ transcripts in transfected K562 cells, qRT-PCR assay was employed using synthesized cDNA from the total RNA isolated from the transfected K562 cells. The data of transient expression experiments are summarized in Table 4. In comparison to the Mock plasmid (pCDNA3.1), the -175 (T>C) mutation increased the expression of the γ -globin gene by 9.2 and 22.6 folds based on Lipofectamine LTX kit and electroporation, respectively. The expression of γ -globin gene in two other constructs, including A and G allele at -588 were (4.6 and 2.3) and (4.3 and 2.0) times, respectively. The variation of expression results might be related to transfection methods. Nevertheless, the expression level of constructs containing A allele was 1.15 fold higher than constructs with G allele. However, the difference between the expression levels was not significant.

DISCUSSION

In order to determine the effect of any mutations in the regulatory regions of the γ -globin gene, the human β -globin gene has to have a normal expression. This depends on many types of regulatory elements residing within the β -globin cluster including the LCR, globin enhancer elements, the individual globin gene promoter and upstream regions. In addition, the promoter of β -globin gene itself may compete with the γ -globin

gene for interaction with the LCR. The LCR also interacts with the minimal γ -gene promoters and sequences of the upstream promoter [2, 5, 13,17]. The linking of the LCR sequences to the γ -globin gene in the expression constructs is critical to detect the impact of any mutations in regulatory elements of the γ -globin gene expression [18]. It has been also presumed that regulation occurs through physical interactions between factors interacting with these elements, which are located at considerable distances from each other [2, 5, 13,17].

In the present study, we have developed a functional assay to compare the possible effect of A γ -588 (A/G) mutations on its gene expression in the K562 cell line. The constructs elements were designed in a way to mimic their *in vivo* transcriptional orientations except the distance between different elements in the construct. The construct included the following elements: the μ LCR, γ -globin gene with mutant and normal promoters and a normal β -globin gene. For the functional study, the rate of transfection in K562 cells was determined by analysis GFP expression using flow cytometry. We found that the yield of transfection was very low, imposing important limitations on functional studies. To overcome these limitations, we optimized and compared different conditions for the transfection and electroporation assays (Table 2). The highest mean rate of transfection by LipofectamineTM LTX kit and electroporation was achieved by 8.74 ± 0.45 and 26.13 ± 2.8 in GFP- positive cells, respectively. The other studies, reported different rate of electroporation, achieved transfection efficiency of about 84.4% and 40% [19, 20]. This rate of transfection is higher than our yield (26.13 ± 2.8), but cell viability in the present study is higher than that previously reported [19, 20]. These main

differences may be due to the use of DMEM as an electroporation buffer in the electroporation assay.

We have previously shown that, the presence of A allele at -588 was highly frequent and closely associated with HbF elevation in β -TI patients, especially those who had IVSII-1/IVSII-1 genotype [3]. Therefore, we aimed to determine the effect of -588 (A/G) mutations in the γ -globin gene expression in the K562 cell line. It has been reported that the γ -175 T>C mutation increases the γ -globin expression and hence, causing nd-HPFH [18]. Thus, we applied the -175 T>C site-directed mutation in the γ -globin promoter in order to determine the ability of our construct model in up-regulating the γ -globin gene *in vitro* system. In the most transient expression assays up-regulation of the -175 (T>C) mutation in *in vitro* system could result in a maximum 20-fold increase in γ -globin promoter strength by reporter gene assay [21-24]. In *in vivo* assay, the -175 mutation also leads to 50 to 100 fold up-regulation in γ -gene in adult erythroid cells [22, 24, 25].

Our data showed that the construct containing the -175 T>C mutation (pCDNA3.1 μ LCR γ ^{-175(C)}) β is able to increase the expression up to 9.2 to 22.6 folds, compared to the Mock plasmid (Table 4). This increase is greater than previously reported *in vitro*. In the pCDNA3.1 μ LCR γ ^{-588(A)} β and pCDNA3.1 μ LCR γ ^{-588(G)} β constructs, the expression of γ -globin gene were (4.6-2.3) and (4.3-2.0) times, respectively when compared to the mock plasmid. However, there was not a significant difference between the expression of γ -globin gene between the construct containing A and the one with G allele at -588. Our result is close to the previously published data indicating that the γ -588 A>G had no effect on γ -globin expression [26]. In addition, transient expression assays showed 2.6 fold decrease in chloramphenicol acetyl transferase protein levels, due to the presence of the γ -588 A>G mutation, compared with the normal γ promoter [11].

In conclusion, our results indicate that -588 A allele does not play a major role in overexpression of γ -globin gene, suggesting that other factors are involved. Therefore, this mutation may not relate to increased promoter function of γ -globin gene. However, it is worth mentioning that the up-regulation of γ -globin gene might be dependent on DNA structure modifications, which is not occurred in this *in vitro* system.

REFERENCES

- Vitale, M., Calzolari, R., Di Marzo, R., Acuto, S., and Maggio, A.A. (2001) A region upstream of the human delta-globin gene shows a stage-specific interaction with globin promoters in erythroid cell lines. *Blood Cells Mol. Dis.* 27: 874-81.
- Langdon, S.D. and Kaufman, R.E. (1998) Gamma-globin gene promoter elements required for interaction with globin enhancers. *Blood* 91: 309-318.
- Hamid, M., Mahjoubi, F., Akbari, M.T., Arab, A., Zeinali, S. and Karimipoor, M. (2009) Molecular analysis of gamma-globin promoters, HS-111 and 3'HS1, in beta-thalassemia intermedia patients associated with high levels of Hb F. *Hemoglobin* 33: 428-438.
- Ragoczy, T., Bender, M.A., Telling, A., Byron, R. and Groudine, M. (2006) The locus control region is required for association of the murine beta-globin locus with engaged transcription factories during erythroid maturation. *Genes Dev.* 20: 1447-1457.
- Molete, J.M., Petrykowska, H., Bouhassira, E.E., Feng, Y.Q., Miller, W. and Hardison, R.C. (2001) Sequences flanking hypersensitive sites of the beta-globin locus control region are required for synergistic enhancement. *Mol. Cell Biol.* 21: 2969-2980.
- Hardison, R., Slightom, J.L., Gumucio, D.L., Goodman, M., Stojanovic, N. and Miller, W. (1997) Locus control regions of mammalian beta-globin gene clusters: combining phylogenetic analyses and experimental results to gain functional insights. *Gene* 205: 73-94.
- de Vooght, K.M., van Wijk, R., Ploos van Amstel, H.K. and van Solinge, W.W. (2007) Characterization of the -16C>G sequence variation in the promoters of both HBG1 and HBG2: convergent evolution of the human gamma-globin genes. *Blood Cells Mol. Dis.* 39: 70-74.
- Liu, L.R., Du, Z.W., Zhao, H.L., Liu, X.L., Huang, X.D., Shen, J., Ju, L.M., Fang, F.D. and Zhang, J.W. (2005) T to C substitution at -175 or -173 of the gamma-globin promoter affects GATA-1 and Oct-1 binding *in vitro* differently but can independently reproduce the hereditary persistence of fetal hemoglobin phenotype in transgenic mice. *J. Biol. Chem.* 280: 7452-7459.
- Hamid, M., Mahjoubi, F., Akbari, M., Zeinali, S. and Karimipoor, M. (2009) The Cretan type of nondeletional hereditary persistence of fetal hemoglobin in an Iranian family. *Ann. Hematol.* 88: 1267-1268.
- Weatherall, D.J., Clegg, J.B., Gibbons, R., Higgs, D.R., Old, J.M., Olivieri, N.F., Thien, S. L. and Wood, W.G. (2001) *The Thalassemia Syndromes. Blackwell Science Publisher Ltd., UK.*
- Patrinos, G.P., Kollia, P., Papapanagiotou, E.,

- Loutradi-Anagnostou, A., Loukopoulos, D. and Papadakis, M.N. (2001) Agamma-haplotypes: a new group of genetic markers for thalassemic mutations inside the 5' regulatory region of the human Agamma-globin gene. *Am. J. Hematol.* 66: 99-104.
12. Schreiber, R., Goncalves, M.S., Junqueira, M.L., Saad, S.T., Krieger, J.E. and Costa, F.F. (2001) The Agamma-195 (C-->G) mutation in hereditary persistence of fetal hemoglobin is not associated with activation of a reporter gene *in vitro*. *Braz. J. Med. Biol. Res.* 34: 489-492.
 13. Chassanidis, C., Kalamaras, A., Phylactides, M., Pourfarzad, F., Likousi, S., Maroulis, V., Papadakis M.N., Vamvakopoulos, N.K., Marinou, V.A., Patrinos, G.P., and Kollia, P. (2009) The Hellenic type of nondeletional hereditary persistence of fetal hemoglobin results from a novel mutation (g-109G>T) in the HBG2 gene promoter. *Ann. Hematol.* 88:549-555.
 14. Khanahmad, H., Noori Dalooi, M.R., Shokrgozar, M.A., Azadmanesh, K., Niavarani, A.R., Karimi, M., Rabbani, B., Khalili, M., Bagheri, R., Maryami, F. and Zeinali, S. (2006) A novel single step double positive double negative selection strategy for beta-globin gene replacement. *Biochem. Biophys. Res. Commun.* 345: 14-20.
 15. Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2 CT method. *Methods* 25:402-408.
 16. Vaerman, J.L., Saussoy, P., and Ingargiola, I. (2004) Evaluation of real-time PCR data. *J. Biol. Regul. Homeost. Agents* 18: 212-214.
 17. Stamatoyannopoulos, G., Josephson, B., Zhang, J. W., and Li, Q. (1993) Developmental regulation of human gamma-globin genes in transgenic mice. *Mol. Cell Biol.* 13:7636-7644.
 18. Balta, G., Brickner, H.E., Takegawa, S., Kazazian, H.H., Jr., Papayannopoulou, T., Forget, B.G., and Atweh, G.F. (1994) Increased expression of the G gamma and A gamma globin genes associated with a mutation in the A gamma enhancer. *Blood* 83:3727-3737.
 19. Delgado-Canedo, A., Santos, D.G., Chies, J.A., Kvitko, K. and Nardi, N.B. (2006) Optimization of an electroporation protocol using the K562 cell line as a model: role of cell cycle phase and cytoplasmic DNases. *Cytotechnology* 51:141-148.
 20. Van Tendeloo, V.F., Ponsaerts, P., Lardon, F., Nijs, G., Lenjou, M., Van Broeckhoven, C., Van Bockstaele, D.R., and Berneman, Z.N. (2001) Highly efficient gene delivery by mRNA electroporation in human hematopoietic cells: superiority to lipofection and passive pulsing of mRNA and to electroporation of plasmid cDNA for tumor antigen loading of dendritic cells. *Blood* 98:49-56.
 21. Nicolis, S., Ronchi, A., Malgarctti, N., Mantovani, R., Giglioni, B. and Ottolenghi, S. (1989) Increased erythroid-specific expression of a mutated HPFH {gamma}-globin promoter requires the erythroid factor NFE-1. *Nucleic Acids Res.* 17:5509-5516.
 22. Gumucio, D. L., Lockwood, W. K., Weber, J. L., Saulino, A. M., Delgrosso, K., Surrey, S., Schwartz, E., Goodman, M., and Collins, F. S. (1990) The -175T----C mutation increases promoter strength in erythroid cells: correlation with evolutionary conservation of binding sites for two trans-acting factors. *Blood* 75:756-761.
 23. Motum, P., Lindeman, R., Harvey, M.P. and Trent, R.J. (1993) Comparative studies of nondeletional HPFH gamma-globin gene promoters. *Exp. Hematol.* 21: 852-858.
 24. McDonagh, K.T., Lin, H.J., Lowrey, C.H., Bodine, D.M. and Nienhuis, A.W. (1991) The upstream region of the human gamma-globin gene promoter. Identification and functional analysis of nuclear protein binding sites. *J. Biol. Chem.* 266:11965-11974.
 25. Liu, L. R., Du, Z. W., Zhao, H. L., Liu, X. L., Huang, X. D., Shen, J., Ju, L. M., Fang, F. D., and Zhang, J. W. (2005) T to C substitution at -175 or -173 of the gamma-globin promoter affects GATA-1 and Oct-1 binding *in vitro* differently but can independently reproduce the hereditary persistence of fetal hemoglobin phenotype in transgenic mice. *J. Biol. Chem.* 280(9): 7452-7459.
 26. Patrinos, G., Loutradi-Anagnostou, A. and Papadakis, M.N. (1995) A novel DNA polymorphism the human A gamma-globin gene (A gamma-588, A-->G) is linked wWith the XMN/polymorphism (G-gamma-158, C-->T). *Hemoglobin* 19:419-423.