



# Can Personality Traits Be Linked to the Development of Alexithymia? Findings from a Cross-Sectional Study

Maedeh Rahimi<sup>1</sup>, Fatemeh Zarimeidani<sup>2</sup>, Hedieh Sadat Bahreini<sup>3</sup>, Atena Kadkhodaei<sup>2</sup>, Rahem Rahmati<sup>2</sup>, Hadi Raeisi Shahraki<sup>4\*</sup>

<sup>1</sup>Students Research Committee, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup>Students Research Committee, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>3</sup>Student Research Committee, YUTBAT, Yeditepe University, Istanbul, Turkey

<sup>4</sup>Department of Epidemiology and Biostatistics, School of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

## OPEN ACCESS

### \*Corresponding Author:

Dept. of Epidemiology and Biostatistics, School of Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

## ABSTRACT

**Introduction:** Scientists have discovered that alexithymia poses a significant challenge for students, surpassing other academic and non-academic issues. Its non-clinical nature implies that individuals afflicted with this condition often fail to realize they are experiencing a problem unless they possess sufficient mental health awareness and understanding. Therefore, this research aimed to explore the correlation between students' personality traits and alexithymia within the student population.

**Methods and Materials:** Following the approval from the ethics committee, for this cross-sectional investigation, a total of 290 students enrolled in the first semester of the 2019-2020 academic year at Shahrekord University of Medical Sciences was chosen using a multi-stage cluster sampling approach. Standardized questionnaires, specifically the NEO-FFI (Five Factor Model) and the TAS-20 (Toronto Alexithymia Scale), were employed to assess alexithymia and personality traits. To compare the scores of various personality traits between students with and without alexithymia, an independent t-test using SPSS 21.0 software was conducted. Additionally, the logistic regression model was utilized to examine the adjusted associations.

**Results:** Among 290 students, the average age of the participants in this study was  $22.6 \pm 3.8$  years. Most students were female (77.9%) and single (88.5%). Based on the TAS-20 assessment, 43.8% exhibited symptoms of alexithymia. Among these students, the neuroticism trait was significantly higher than the non-alexithymic students ( $27.20 \pm 3.90$  versus  $25.48 \pm 4.90$ ;  $p = 0.01$ ), indicating a greater tendency toward emotional instability. Additionally, the average score for the conscientiousness trait was significantly higher in the alexithymic students ( $20.16 \pm 3.84$  versus  $21.58 \pm 5.00$ ;  $p = 0.01$ ), suggesting a greater inclination towards being organized and responsible. The logistic regression analysis results indicated that for each unit increase in the neuroticism trait, there was a 10% increase in the odds of experiencing alexithymia. Conversely, for each unit increase in the score of conscientiousness trait, there was an 8% decrease in the odds of experiencing alexithymia.

**Conclusion and Discussion:** Given the notable correlation between personality traits and alexithymia, improving screening methods and intervention programs focusing on personality traits can help prevent or alleviate the symptoms among university students. Targeted interventions can be developed to support and assist students experiencing this condition by identifying and addressing specific personality traits associated with alexithymia.

### Citation:

Rahimi M, Zarimeidani F, Bahreini HS, Kadkhodaei A, Rahmati R, Raeisi Shahraki H. Can Personality Traits Be Linked to the Development of Alexithymia? Findings from a Cross-Sectional Study. *Iranian biomedical journal* 2024; 28(7): 10.

Keywords: Personality, Psychiatry, Statistical models

