



# Effect of Anzerut (*Astragalus fasciculifolius*) on Spatial Memory Impairment and Its Potential Antioxidant Effects in a Rat Model of Multiple Sclerosis

Mehrdad Shahrani Korrani<sup>1\*</sup>, Negar Khalili Geshnigani<sup>2</sup>, Fatemeh Azizi Farsani<sup>1</sup>, Maryam Anjomshoa<sup>1</sup>, Najmeh Asgharzadeh<sup>1</sup>, Seyed Ali Hosseinian

<sup>1</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>2</sup>Students Research Committee, Shahrekord University of Medical Sciences, Shahrekord, Iran

## ABSTRACT

### OPEN ACCESS

#### \*Corresponding Author:

Medical Plants Research Center,  
Shahrekord University of Medical  
Sciences, Shahrekord, Iran

#### Citation:

Shahrani Korrani M, Khalili Geshnigani N, Azizi Farsani F, Anjomshoa M, Asgharzadeh N, Hosseinian SA. Effect of Anzerut (*Astragalus fasciculifolius*) on Spatial Memory Impairment and Its Potential Antioxidant Effects in a Rat Model of Multiple Sclerosis. *Iranian biomedical journal* 2024; 28(7): 97.

**Introduction:** Multiple Sclerosis (MS) is an inflammatory autoimmune disease in which the myelin sheaths of nerve cells are damaged in the brain and spinal cord. Cognitive changes, including memory impairment, are common in MS patients, and there is still no definitive treatment available. Plants of the genus *Astragalus* from the legume family contain more than 900 species of annual and perennial herbaceous plants in Iran, most of which are endemic to the region. Recent studies have shown that the *Astragalus* has antioxidant, neuroprotective, and anticonvulsant effects. Anzerut is a species on which limited studies have been conducted. Considering the various biological effects of the *Astragalus* genus, we aimed to investigate the impact of Anzerut on spatial memory impairment in a rat model of MS.

**Methods and Materials:** In the present study, 42 male rats weighing between 230-280 g were randomly divided into six groups of seven rats each. Ethidium bromide was used to induce MS. Then, doses of 150, 450, and 800 mg/kg of Anzerut extract were administered intraperitoneally for 14 days. Behavioral testing using the Morris water maze, hippocampal malondialdehyde (MDA) level measurement, and total antioxidant capacity were evaluated. Finally, one-way ANOVA followed by Tukey's test will be used to compare the data.

**Results:** The injection of ethidium bromide caused a significant decrease in the time spent ( $p = 0.01$ ) and distance traveled ( $p = 0.001$ ) in the target quadrant on the test day compared to the control group. Treatment with Anzerut at a dose of 150 mg/kg for 14 days significantly increased the time spent and distance traveled ( $p = 0.05$ ) in the target quadrant on the test day compared to the MS group. Additionally, the injection of ethidium bromide resulted in a significant increase in hippocampal MDA level and a decrease in the total antioxidant capacity of the hippocampus.

**Conclusion and Discussion:** The injection of ethidium bromide into the CA1 region of the hippocampus induces MS, while the administration of Anzerut extract improves learning and memory in the experimental MS model. This effect appears to be mediated through antioxidant pathways in the brain.

**Keywords:** Antioxidant; *Astragalus* plant, Multiple sclerosis

