



Food Insecurity and Breast Cancer Incidence: An Ecological Study in Razavi Khorasan Province, Iran

Iman Mousaei¹, Amin Moradi¹, Ali Taghipour¹, Maryam Salari², Ehsan Mosafarkhani^{3*}

¹Department of Epidemiology, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

²Department of Biostatistics, School of Health, Social Determinants of Health Research Center, Basic Sciences Research Institute, Mashhad University of Medical Sciences, Mashhad, Iran

³Department of Epidemiology, School of Health, Management and Social Determinants of Health Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

OPEN ACCESS

*Corresponding Author:

Dept. of Epidemiology, School of Health, Management and Social Determinants of Health Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Citation:

Mousaei I, Moradi A, Taghipour A, Salari M, Mosafarkhani E. Food Insecurity and Breast Cancer Incidence: An Ecological Study in Razavi Khorasan Province, Iran. *Iranian biomedical journal* 2024; 28(7): 375.

ABSTRACT

Introduction: Food insecurity has emerged as a significant public health concern worldwide, yet its association with breast cancer (BC) incidence remains understudied, particularly in low- and middle-income countries like Iran.

Methods and Materials: This cross-sectional study utilized electronic healthcare data from Mashhad University of Medical Sciences, aggregated at the city level across the population covered by Mashhad University in Iran in 2022. Age-standardized incidence rates of BC in each city served as the primary outcome variable. A multivariable linear regression model was constructed with age-standardized BC incidence rate as the dependent variable and food insecurity, mean age, obesity, diabetes, smoking, and total fertility rate as independent predictor variables.

Results: Higher levels of food insecurity were associated with lower age-standardized BC incidence rates ($\beta = -0.0103$; 95% CI: -0.0147, -0.0059; $p = 0.001$) after adjusting for mean age, obesity, diabetes, smoking prevalence, and total fertility rate. Obesity ($\beta = 0.9460$; 95% CI: 0.7312, 1.1608; $p = 0.001$) and smoking prevalence ($\beta = 0.3899$; 95% CI: 0.2751, 0.5047; $p = 0.001$) were positively associated with BC incidence. However, diabetes prevalence ($\beta = -2.4594$; 95% CI: -3.0742, -1.8446; $p = 0.001$) and total fertility rate ($\beta = -8.5297$; 95% CI: -10.6775, -6.3819; $p = 0.001$) showed inverse associations.

Conclusion and Discussion: This ecological study demonstrated the association of higher levels of food insecurity with lower age-standardized BC incidence rates in Razavi Khorasan province, Iran. These findings highlight the importance of addressing social determinants like food insecurity in cancer prevention and control strategies, particularly in low- and middle-income countries.

Keywords: Food insecurity, Breast neoplasms, Iran

