

## Thyroiditis Incidence and COVID-19 Epidemic in Sanandaj, Iran: A Time-Series Analysis

Maziar Afshar<sup>1</sup>, Abbas Aghaei<sup>2</sup>, Bijan Nuri<sup>2</sup>, Bahar Moasses Ghafari<sup>3\*</sup>

<sup>1</sup>Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran <sup>2</sup>Department of Epidemiology and Biostatistics, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran <sup>3</sup>Department of Nuclear Medicine, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

## ABSTRACT

## **OPEN ACCESS**

Citation:

Incidence

429

\*Corresponding Author: Dept. of Nuclear Medicine, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

Afshar M, Aghaei A, Nuri B,

Moasses Ghafari B. Thyroiditis

and

Epidemic in Sanandaj, Iran: A

Time-Series Analysis. Iranian

biomedical journal 2024; 28(7):

**Introduction:** This study aimed to investigate the potential correlation between reported cases of thyroiditis and the COVID-19 pandemic in Sanandaj, Iran, using time-series analysis techniques. The data was collected from the two nuclear medicine centers in Kurdistan province, which are the only centers and the main referral facilities for nuclear medicine.

**Methods and Materials:** Based on thyroid scan findings from these two centers in 2019 and 2020, individuals diagnosed with thyroiditis were included. Timeseries analysis, cross-correlation analysis, and analysis of lagged effects examined the temporal patterns and potential correlations between thyroiditis cases and COVID-19 data.

**Results:** The time-series analysis employed an autoregressive integrated moving average (ARIMA) model with a lag length of 3. While the cross-correlation analysis showed positive and significant correlations at lags 1 to 5, with the highest cross-correlation coefficient of 0.4674 (p = 0.05) at lag 3, suggesting a potential 3-month optimal lag, this lag was not meaningful. The analysis of lagged effects did not reveal a statistically significant relationship between the two variables (F(6, 27) = 1.18, p = 0.3444), indicating that the trends of thyroiditis and COVID-19 cases are not related and COVID-19 cannot predict subsequent thyroiditis cases.

**Conclusion and Discussion:** Despite clinical and observational reasoning suggesting a potential association between COVID-19 and thyroiditis, this study did not find strong evidence for a significant correlation between these two incidents in the population studied.

Keywords: COVID-19, Nuclear medicine, Thyroiditis

COVID-19

