

Expression of Epithelial-Mesenchymal Transition-Related Genes in Primary Brain Tumors

Shiva Bayat^{1#}, Negar Mousavi^{1#}, Fatemeh Vand-Rajabpour¹,
Ali Mamivand¹, Mina Tabrizi^{1*}, Alireza Khoshnevisan²

¹Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, P.O. Box 14155-6447, Tehran 14176-13151, Iran; ²Neurosurgery Department, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

OPEN ACCESS

Article type: Review Article

Received: July 26, 2025

Revised: November 23, 2025

Accepted: December 5, 2025

Published online: December 9, 2025

How to cite:

Bayat S, Mousavi N, Vand-Rajabpour F, Mamivand A, Tabrizi M, Khoshnevisan A. Expression of Epithelial-Mesenchymal Transition-Related Genes in Primary Brain Tumors. *Iran. Biomed. J.* 2026; 30(1): 59-66



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

ABSTRACT

Background: Gliomas are the most common malignant primary brain tumors in adults. Meningiomas, on the other hand, are benign primary brain tumors treated by surgical resection. Epithelial-to-mesenchymal transition (EMT), stemness, and leader behavior are among the most important complex processes contributing to tumor progression and aggressiveness. Thus, the evaluation of molecular targets involved in the EMT processes in primary brain tumors can help elucidate the molecular mechanisms underlying glioma aggressiveness, ultimately identifying potential areas for further research.

Methods: We conducted expression analysis of zinc finger E-box homeobox 1 (*ZEB1*), Sry-related HMG box (*SOX2*), and *p21* in 31 glioma (15 glioblastoma [GBM] and 16 non-GBM) and 44 meningioma samples using quantitative real-time PCR. We also compared expression levels between glioma and meningioma, as well as between GBM and non-GBM tumor samples.

Results: The expression levels of *SOX2* and *ZEB1* showed a significant increase in gliomas compared to meningiomas ($p = 0.016$ and 0.017 , respectively). However, *p21* did not reveal any significant difference in expression between the two groups. Notably, *p21* was the only gene that exhibited a significant increase in expression in GBM samples compared to non-GBM samples, with a p value of 0.027 .

Conclusion: Our results suggest that *SOX2* and *ZEB1* expression levels have a significant role in glioma progression. Interestingly, based on the elevated *p21* expression level in the GBM, we also suggest that aggressive tumor features arise from distinct molecular mechanisms depending on tumor subtype, not a single universal pathway. **DOI: 10.61882/ibj.5133**

Keywords: Glioma, Meningioma, SOX2, ZEB1

Corresponding Author: Mina Tabrizi

Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, P. O. Box 14155-6447, Tehran 14176-13151, Iran; E-mail: tabrizi@tums.ac.ir; ORCID ID: 0000-0002-1677-214X

[#]These authors contributed equally.

1. INTRODUCTION

Primary brain tumors are heterogeneous neoplasms derived from various cell types of the central nervous system (CNS). Among these tumors, glioma and meningioma are the two most common primary brain tumors in adults, each exhibiting distinct malignant potential.

Glioma primarily originates from glial cells and is mostly found in the CNS. Generally, these tumors are more prevalent in men than in women. Gliomas can be classified into several cell types: astrocytic (astrocytoma), oligodendrocytic (oligodendroglioma), ependymal (ependymoma), or mixed. The World Health Organization classified gliomas into grades I to IV

based on differentiation, anaplasia, aggressiveness, and histopathological features^[1]. Glioblastoma (GBM), the most lethal subtype, has a five-year survival rate ranging from 0.05% to 4.7% after diagnosis and occurs in 0.59 to 3.69 per 100,000 individuals^[2-4]. GBM comprises approximately 82% of all glioma cases, with an age-adjusted incidence rate ranging from 4.67 to 5.73 per 100,000 people^[5,6]. Although GBM tumors are highly invasive and tend to infiltrate the surrounding parenchyma, they are restricted to the CNS and rarely metastasize to regional lymph nodes, lungs, and pleura, and occasionally to the bone and liver.

Meningioma, the most common non-malignant brain tumor, arises from the arachnoid cap cells located at various sites in the CNS, including the spinal cord. More than 80% of these tumors remain benign with silent progression, which limits the availability of comprehensive epidemiological data on their characteristics. The overall five-year survival rate for meningioma patients is 70%, though it decreases with age. Meningioma can often be treated with surgical resection due to its benign nature; however, 15-20% of these tumors may display aggressive behavior, necessitating further adjuvant therapies^[2].

A wide range of mechanisms has been studied and proven to be implicated in glioma aggressiveness. It has been confirmed that glioma stem-like cells have features that allow for self-renewal and production of progenitors that facilitate tumor progression^[7]. Furthermore, mutations in isocitrate dehydrogenase 1 and changes in methylation patterns of O-6-methylguanine-DNA methyl transferase gene are also among the contributing factors in tumor malignancy^[8,9]. A key mechanism contributing to tumor aggressiveness is the ability of glioma cells to invade and migrate into the surrounding normal brain tissue in the brain, leading to incomplete removal and resistance to radiotherapy^[10].

Epithelial-to-mesenchymal transition (EMT) is a complex process through which tumor cells lose their epithelial characteristics, such as E-cadherin expression, and attain mesenchymal features, resulting in decreased adhesion to adjacent cells. EMT leads to increased motility and migration, giving tumor cells more malignant properties. Beyond EMT itself, accompanying epigenetic changes in these cells can drive the formation of cancer stem cells, a feature associated with increased tumor aggressiveness and unfavorable prognosis^[11,12]. These reversible changes are significantly influenced by the tumor microenvironment. A plethora of signaling pathways, including Wnt, play important roles in the EMT process, in which tumor cells acquire metastatic abilities^[11].

Zinc finger E-box homeobox 1 (*ZEB1*) is a key transcription factor involved in EMT and cancer progression^[13-16]. There is increasing evidence for its

role in drug resistance to temozolomide^[17], and its expression is a contributing factor to shorter survival of patients.

p21 is a cell cycle regulator involved in the induction of senescence in response to DNA damage and p53 activation. It has also been proven to play a significant role in the self-renewal of neural stem cells by repressing Sry-related HMG box (*SOX2*) expression, a transcription factor that is also important in cancer stem cell biology. Recently, studies have suggested that p53, p21, and cyclin-dependent kinase (CDK) inhibition, the genes that are developmentally controlled or induced by injury, may significantly influence collective leader-driven cell migration^[18]. p53 is a well-documented key factor in leader-driven cell migration by modulating downstream p21 and CDK activity^[18].

Given the role of *SOX2*, *ZEB1*, and *p21* in stemness, EMT, leader cell behavior, and migration processes, these factors were selected for further analysis in this study. Since glioma is a malignant brain tumor with high potential for metastasis, invasion, and relapse, and knowing that meningioma is a known benign brain tumor, we hypothesized that these types of brain tumors might be appropriate in vivo models for comparison. Thus, this study analyzed the expression levels of *SOX2*, *ZEB1*, and *p21* in primary brain tumors to assess the significance of these molecules in brain tumorigenesis. The same comparison was conducted between GBM, as the most malignant glioma grade, and lower-grade gliomas treated as non-GBM cases. This approach aligns with recent literature that recognizes the distinct malignant features of GBM. Our study was in line with previous studies^[19-24], demonstrating the upregulation of stemness and EMT markers, specifically *SOX2* and *ZEB1* ($p = 0.016$ and $p = 0.017$), in glioma tumors compared to meningioma tumors. This is most likely due to cancer-specific mechanisms modulating the expression of these molecules. Furthermore, our findings support the data previously acquired by other investigators^[19-24].

2. MATERIALS AND METHODS

2.1. Patient samples

Seventy-five fresh brain tumor samples, 31 glioma (15 GBM and 16 non-GBM) and 44 meningioma samples, were collected from Shariati Hospital affiliated to Tehran University of Medical Sciences, Tehran, Iran, from 2014 to 2016. All pathology results and definitive diagnoses of tumor types and grades were confirmed by an expert neuropathologist.

2.2. RNA extraction, DNase treatment, and complementary DNA (cDNA) synthesis

All procedures related to RNA extraction, DNase I treatment, and cDNA synthesis have been described

previously and followed the MIQE guidelines^[19]. The steps of total RNA extraction were based on a modified Tripure Isolation Reagent protocol. Briefly, tissue samples were subjected to cycles of freeze and thaw, followed by homogenization with the lysis buffer for the simultaneous isolation of RNA, DNA, and protein in a single-step liquid-phase separation. Following centrifugation of the top aqueous phase containing RNA, RNA was precipitated using isopropanol and then purified using ethanol 70% for the removal of the residual salts. After air-drying, the RNA pellet was resuspended in RNase-free water. The yield and purity of the extracted RNA were ultimately assessed using a NanoDrop spectrophotometer (A260/A280 and A260/A230 ratios). To remove any residual genomic DNA contamination and obtain RNA with higher purity for downstream analyses, we treated the samples with RNase-free DNase I according to the manufacturer's instructions. For the synthesis of the first-strand cDNA, PrimeScript RT Reagent kit (Takara Bio Inc., Shiga, Japan) was applied to the RNA templates, and reverse transcription of the templates was carried out accordingly, utilizing random hexamers and oligo dT primers.

2.3. Quantitative real-time PCR (qRT-PCR)

Specific primers of mRNA expression were designed by primer-design (NCBI) and Gene Runner or adopted (Table 1). The primer efficiency and R² values for each of *GAPDH*, *p21*, *SOX2*, and *ZEB1* genes were assessed using a standard curve. qRT-PCR was performed by utilizing SYBR Premix Ex TagTM (Takara Bio Inc.) and Light Cycler® 96 System (Roche Life Science, Germany). The steps of real-time PCR cycles were as follows: initiation for 30 s at 95°C, followed by 40 cycles including 6 s at 95°C and 20 s at 60°C. Melting curves were analyzed at the end of each real-time PCR reaction to check the presence of any primer dimers and the specificity of the amplicon peaks. To ensure the removal of any genomic DNA from each RNA sample, we performed qRT-PCR reactions using *SOX2* primers. Absence of an amplification peak was a confirmation of the perfect function of DNase I treatment. Expression

data for each tissue sample was normalized to *GAPDH* gene expression as the housekeeping gene. All experiments were performed in duplicates.

2.4. Statistical analysis

Statistical analysis was performed with SPSS (IBM Corp., Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp., USA). Data were presented with mean, standard deviation, and range. The Q-Q plot and the Kolmogorov-Smirnov test were used to evaluate normal distribution. The analysis of variance (ANOVA) and the Mann-Whitney test were conducted to compare groups. The Bonferroni method was used for multiple comparisons. The ROC curve and the area under the curve (AUC) were used to evaluate the sensitivity and specificity of each mRNA. The Pearson correlation coefficient was assessed between the rankings of two variables. A *p* value less than 0.05 was deemed statistically significant for a confidence interval of 95%.

3. RESULTS

3.1. Patients' demographic and histopathologic data

The summarized demographic data of the patients are presented in Table 2.

3.2. Gene expression analysis results

Each primer pair was validated using standard curves. All the primers demonstrated high linearity with R² values ranging from 0.95 to 0.99. The calculated amplification efficiencies derived from the slopes fell between 93% and 118%, rendering the primers reliable for use in downstream tests. The expression levels of *ZEB1*, *SOX2*, and *p21* were evaluated by qRT-PCR in glioma and meningioma samples, and *GAPDH* was used as the housekeeping gene. *GAPDH*^[25] was selected as a control gene based on the existing literature, including the one by Said et al, suggesting the use of *GAPDH* since its expression is not affected by various conditions and remains stable. According to Mathur et al.^[26], *GAPDH* is among the panel of genes that are frequently used as endogenous controls in 80% of gene expression in glioma tumors. In this study, we used a different

Table 1. Primer sequences for the analysis of *SOX2*, *ZEB1*, and *p21* mRNA expression using RT-PCR

Gene name	Primer	Sequence	Tm	Product length (bp)
<i>p21</i>	Forward	GACTCTCAGGGTTCGAAAACG	58	93
	Reverse	GGATTAGGGCTTCCTCTTGG	57	
<i>ZEB1</i>	Forward	CATTTTCTCTGAGGCACCTG	56	91
	Reverse	TGAAAATGCATCTGGTGTTC	57	
<i>SOX2</i>	Forward	CAGCTCGCAGACCTACATGA	59	152
	Reverse	TGGAGTGGGAGGAAGAGGTA	58	

Table 2. Patients' demographic data in glioma vs. meningioma and GBM vs. non-GBM subgroups

Age	Glioma vs. meningioma	GBM vs. non-GBM
Mean	43.1 vs. 58.4	45.3 vs. 40.9
Minimum	10 vs. 31	10 vs. 23
Maximum	77 vs. 80	64 vs. 77
Male/female ratio	61%/39% vs. 30%/70%	53%/47% vs. 69%/31%

approach to find significant differential expression in a spectrum of primary brain tumors, concerning aggressiveness, including glioma versus meningioma and GBM versus non-GBM. At first, we focused on the aggressive features of primary brain tumors and studied the expression levels of *ZEB1*, *SOX2*, and *p21* in glioma samples, as malignant primary brain tumors, versus meningioma samples, as benign primary brain tumors. We found that *ZEB1* and *SOX2* as EMT and stemness factors, were upregulated in glioma samples compared to meningioma samples, with $p = 0.017$ and $p = 0.016$ and fold change of 2.8 and 16.9, respectively. On the

contrary, *p21* displayed an elevated expression level with $p = 0.027$ in the most aggressive glioma grade, GBM. GBM is now considered a distinct entity with an utterly different biological landscape compared to the lower-grade non-GBM tumor samples, which therefore justifies the comparison carried out between the two subgroups. The relevant fold change was found to be approximately 2.5. The results of gene expression analyses are presented in Figure 1 and Table 3. In order to differentiate glioma from meningioma, we measured the specificity and sensitivity of our studied genes using the ROC curve and AUC (Fig. 2 and Table 4).




Fig. 1. The log-transformed $2^{-\Delta\Delta CT}$ values for the differentially expressed genes. (A) *SOX2* and (B) *ZEB1* in glioma vs. meningioma; (C) *p21* in GBM vs. non-GBM subgroups.

Table 3. Relevant *p* values and effect sizes of *SOX2*, *ZEB1*, and *p21* expression in glioma vs. meningioma and GBM vs. non-GBM

Comparison	Gene	Test	Statistic (U)	<i>p</i> value	Effective size (Rank-biserial correlation)
Glioma vs. meningioma	<i>P21</i>	Mann-Whitney U	603	0.492	0.0953
	<i>ZEB1</i>	Mann-Whitney U	460	0.017	-0.3255
	<i>SOX2</i>	Mann-Whitney U	425	0.016	-0.3320
GBM vs. non-GBM	<i>P21</i>	Mann-Whitney U	64	0.027	-0.467
	<i>ZEB1</i>	Mann-Whitney U	93	0.299	0.225
	<i>SOX2</i>	Mann-Whitney U	106	0.593	-0.117

4. DISCUSSION

The two most common types of primary brain tumors are glioma and meningioma, and most importantly, the deadliest tumor is glioma grade IV, known as GBM. Glioma is the most common malignant primary brain tumor in adults, with most cases showing malignant features^[27]. In contrast, 80% of meningiomas are benign, making them suitable candidates for a comparison of biological characteristics.

There are several mechanisms contributing to glioma pathogenesis, including EMT. Multiple genes, such as *ZEB1*, *SOX2*, and *p21*, are known to play roles in EMT. These genes were selected for this study because they had previously been studied by our research group in skin cancer^[28]. Thus, these three molecular targets were chosen for further investigation in glioma and compared to benign meningioma tumors to elucidate possible underlying factors contributing to their differing behaviors from an EMT perspective.

Herein, we reported a significant upregulation of *ZEB1* and *SOX2* in glioma samples compared to meningioma samples, which is in agreement with previous studies^[19-24]. Additionally, we observed elevated expression of *p21* in GBM in comparison to non-GBM glioma samples.

Regarding the null findings of *p21*, it is important to

note that this gene plays a dual role in cancer, acting both as a tumor suppressor and as a factor that promotes tumor survival under stress conditions. The lack of significant difference in *p21* expression between gliomas and meningiomas suggests that this gene is not a reliable marker for distinguishing tumor types. This observation is in agreement with the reports indicating that *p21* expression varies widely across brain tumors and may be influenced more by microenvironmental stressors or therapeutic exposure than by the inherent biology of the baseline tumor. However, the notable increase in *p21* levels in GBM compared to non-GBM reflects the heightened DNA damage, hypoxia, and cellular stress typically associated with GBM.

SOX2, a key transcription factor and a stemness molecule belonging to the SOX protein family and containing the SOX2 DNA-binding domain, has been reported to be involved in tumor invasion and migration in malignant gliomas. Downregulation of *SOX2* using small interfering RNAs has been shown to reduce invasive capabilities^[29]. This gene has also been studied in medulloblastomas, which is implicated in resistance to chemotherapy in these tumors^[30]. Our findings demonstrated that *SOX2* expression was significantly upregulated in glioma compared to meningioma. Likewise, expression of SOX2 protein was verified in




Fig. 2. The ROC curves for *p21*, *SOX2*, and *ZEB1* genes in (A) glioma vs. meningioma and (B) GBM vs. non-GBM.

Table 4. ROC curve summary for gene expression ($\Delta\Delta CT$) in glioma vs. meningioma and GBM vs. Non-GBM

Comparison	Gene	AUC	Std. error	95% CI	p value
Glioma vs. meningioma	<i>P21</i>	0.548	0.0688	0.413-0.682	0.488
	<i>ZEB1</i>	0.663	0.0642	0.537-0.789	0.011
	<i>SOX2</i>	0.666	0.0657	0.537-0.795	0.012
GBM vs. non-GBM	<i>P21</i>	0.733	0.0923	0.552-0.914	0.011
	<i>ZEB1</i>	0.612	0.1103	0.396-0.829	0.308
	<i>SOX2</i>	0.558	0.1101	0.343-0.774	0.596

The table corresponds to the ROC curves presented in Figure 2.

GBM biopsies using immunohistochemistry and Western blotting^[31]. One study reported low or negative expression of SOX2 protein in meningioma, while another study reported its positive expression in 11 samples of grade I meningioma^[22,32]. Therefore, more studies are needed to confirm and clarify the importance of expression levels of SOX2 at the mRNA and protein levels across different brain tumors, particularly concerning its role in tumorigenesis.

ZEB1 is a transcription factor involved in tumor invasion and metastasis by inducing the EMT process in epithelial tumors. Additionally, it is believed to suppress the expression of E-cadherin, an epithelial biomarker, and promote mesenchymal traits by suppressing its targets, including *Crumb3*, *HUGL2*, and *PATJ*, all leading to tumor metastasis. *ZEB1* overexpression has been reported in various cancers, including bladder, colon, and breast^[33-39]. In a genomic study, Edwards and colleagues observed *ZEB1* deletion in 15% of glioma patients and suggested it as a poor prognostic marker^[40]. Similarly, another genomic study demonstrated *ZEB1* deletion in 15% of grades II and III gliomas and also in 50% of GBM samples^[39]. Our finding of increased *ZEB1* expression does not contradict previous studies, as *ZEB1* expression can be regulated independently of its copy number status. The elevated expression levels of *ZEB1* observed in our cohort support the notion that gliomas rely on stemness and EMT pathways to sustain their malignant phenotype, distinguishing them from the less aggressive meningioma tumors. Moreover, the absence of a significant difference in *ZEB1* and *SOX2* expression between GBM and non-GBM samples highlights that these genes may be more useful for distinguishing tumor type (glioma vs. meningioma), rather than explaining grade progression within gliomas. Overall, these findings, similar to previous studies on other organ systems^[28], highlight the importance of adopting a comprehensive approach when investigating brain tumors with opposing features, based on their aggressive characteristics. This approach could help identify key regulatory molecules involved in their

distinct behaviors, emphasizing the need for further studies on the mechanistic roles of *ZEB1* and *SOX2* genes at the protein level.

5. CONCLUSION

Our results suggest that the molecular pathways modulating *SOX2* and *ZEB1* markers are cancer-specific and likely distinct across tumor types. Future studies should include tumor grades, increase sample size, and examine classic EMT markers (E-cadherin and N-cadherin) as well as stemness and leader-cell behavior to better elucidate the molecular network driving aggressiveness and invasion in CNS tumors.

DECLARATION

Acknowledgments

Not applicable.

Generative AI and AI-assisted technologies

The authors declare that no AI-assisted technologies were used during the preparation of this manuscript.

Ethical approval

All experimental procedures in this study were in accordance with the Helsinki Declaration of 1975, as revised in 1983 and approved by the ethical standards of Tehran University of Medical Sciences, Tehran, Iran (ethical code: IR.TUMS.MEDICINE.REC.1395.1689).

Consent to participate

All participants signed the written informed consent.

Consent for publication

All authors reviewed the results and approved the final version of the manuscript.

Authors' contributions

SB: contributed to the study design, experimental procedures, and drafted the manuscript; NM: and FVR: contributed to the experimental procedures;

AM: contributed to the analysis of the findings; MT: contributed to the study design, supervision of the work, interpretation of the results, and revision of the manuscript; AK: contributed to the samples' surgical resection and diagnosis of the tumor types.

Data availability

All relevant data can be found within the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was financially supported by Tehran University of Medical Sciences, Tehran, Iran.

Supplementary information

The online version does not contain supplementary material.

REFERENCES

- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, et al. The 2007 WHO classification of tumors of the central nervous system. *Acta Neuropathol.* 2007;114(2):97-109.
- Ostrom QT, Gittleman H, Farah P, Ondracek A, Chen Y, Wolinsky Y, et al. CBTRUS statistical report: Primary brain and central nervous system tumors diagnosed in the United States in 2006–2010. *Neuro Oncol.* 2013;15(2):1-56.
- Arora RS, Alston RD, Eden TO, Estlin EJ, Moran A, Birch JM. Age-incidence patterns of primary CNS tumors in children, adolescents, and adults in England. *Neuro Oncol.* 2009;11(4):403-13.
- Lee CH, Jung KW, Yoo H, Park S, Lee SH. Epidemiology of primary brain and central nervous system tumors in Korea. *J Korean Neurosurg Soc.* 2010;48(2):145-52.
- Larjavaara S, Mäntylä R, Salminen T, Haapasalo H, Raitanen J, Jääskeläinen J, et al. Incidence of gliomas by anatomic location. *Neuro Oncol.* 2007;9(3):319-25.
- Gousias K, Markou M, Voulgaris S, Goussia A, Voulgari P, Bai M, et al. Descriptive epidemiology of cerebral gliomas in northwest Greece and study of potential predisposing factors, 2005–2007. *Neuroepidemiology.* 2009;33(2):89-95.
- Bradshaw A, Wickremsekera A, Tan ST, Peng L, Davis PF, Itinteang T. Cancer stem cell hierarchy in glioblastoma multiforme. *Front Surg.* 2016;3:21.
- Sheehan J, Shaffrey ME, Gupta B, Larner J, Rich JN, Park DM. Improving the radiosensitivity of radioresistant and hypoxic glioblastoma. *Future Oncol.* 2010;6(10):1591-601.
- Jampani P, Venkatesh BS, Sujun KC, Kabindra K. Review on glioblastoma multiforme (GBM). *World J Pharm Res.* 2018;7(17):326-37.
- Paw I, Carpenter RC, Watabe K, Debinski W, Lo H-W. Mechanisms regulating glioma invasion. *Cancer Lett.* 2015;362(1):1-7.
- Iwadate Y. Epithelial-mesenchymal transition in glioblastoma progression. *Oncol Lett.* 2016;11(3):1615-20.
- Nørøxe DS, Poulsen HS, Lassen U. Hallmarks of glioblastoma: A systematic review. *ESMO Open.* 2017;1(6):000144.
- Zhang P, Wei Y, Wang L, Debeb BG, Yuan Y, Zhang J, et al. ATM-mediated stabilization of ZEB1 promotes DNA damage response and radioresistance through CHK1. *Nat Cell Biol.* 2014;16(9):864-75.
- Peinado H, Olmeda D, Cano A. Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat Rev Cancer.* 2007;7(6):415-28.
- Aigner K, Dampier B, Descovich L, Mikula M, Sultan A, Schreiber M, et al. The transcription factor ZEB1 (deltaEF1) promotes tumor cell dedifferentiation by repressing master regulators of epithelial polarity. *Oncogene.* 2007;26(49):6979-88.
- Spaderna S, Schmalhofer O, Wahlbuhl M, Dimmler A, Bauer K, Sultan A, et al. The transcriptional repressor ZEB1 promotes metastasis and loss of cell polarity in cancer. *Cancer Res.* 2008;68(2):537-44.
- Siebzehnruhl FA, Silver DJ, Tugertimur B, Deleyrolle LP, Siebzehnruhl D, Sarkisian, MR, et al. The ZEB1 pathway links glioblastoma initiation, invasion and chemoresistance. *EMBO Mol Med.* 2013;5(8):1196-212.
- Kozyrska K, Pilia G, Vishwakarma M, Wagstaff L, Goschorska M, Cirillo S, et al. P53 directs leader cell behavior, migration, and clearance during epithelial repair. *Science.* 2022;375(6581):8876.
- Gross MW, Kraus A, Nashwan K, Mennel H-D, Engenhart-Cabillic R, Schlegel J. Expression of p53 and p21 in Primary Glioblastomas. *Strahlenther Onkol.* 2005;181(3):164-71.
- Terzi A, Saglam EA, Barak A, Soylemezoglu F. The significance of immunohistochemical expression of Ki-67, p53, p21, and p16 in meningiomas tissue arrays. *Pathol Res Pract.* 2008;204(5):305-14.
- Annovazzi L, Mellai M, Caldera V, Valente G, Schiffer D. SOX2 expression and amplification in gliomas and glioma cell lines. *Cancer Genomics Proteomics.* 2011;8(3):139-48.
- Xiao Z-Y, Chen X-J, Pan Q, Yang Q-Z, Li K-Z. Expression of nestin, CD133 and Sox2 in meningiomas. *Turk Neurosurg.* 2018;28(6):910-14.
- Nesvick CL, Zhang C, Edwards NA, Montgomery BK, Lee M, Yang C, et al. ZEB1 expression is increased in IDH1-mutant lower-grade gliomas. *J Neurooncol.* 2016;130(1):111-22.
- Wallesch M, Pachow D, Blucher C, Firsching R, Warnke J-P, Braunsdorf WEK, et al. Altered expression of E-cadherin-related transcription factors indicates partial epithelial-mesenchymal transition in aggressive meningiomas. *J Neurol Sci.* 2017;380:112-21.
- Said HM, Hagemann C, Stojic J, Schoemig B, Vince, GH, Flentje M, et al. GAPDH is not regulated in human glioblastoma under hypoxic conditions. *BMC Mol Biol.* 2007;8:55.

26. Mathur D. Selection of suitable housekeeping genes for expression analysis in glioblastoma using qRT-PCR. *Ann Neurosci*. 2014;21(2):62-3.
27. Pardal R, Clarke MF, Morrison SJ. Applying the principles of stem-cell biology to cancer. *Nat Rev Cancer*. 2003;3(12):895-902.
28. Ahmadi-Beni R, Vand-Rajabpour F, Ahmadifard M, Daneshpazhooch M, Noormohammadpour P, Rahmati J, et al. Decreased *Sox2* messenger RNA expression in basal cell carcinoma. *Indian J Dermatol*. 2020;65(3):178-82.
29. Han X, Fang X, Lou X, Hua D, Ding W, Foltz G, et al. Silencing *SOX2* induced mesenchymal-epithelial transition and its expression predicts liver and lymph node metastasis of CRC patients. *Plos One*. 2012;7(8):41335.
30. Vanner RJ, Remke M, Gallo M, Selvadurai HJ, Coutinho F, Lee L, et al. Quiescent *sox2(+)* cells drive hierarchical growth and relapse in sonic hedgehog subgroup medulloblastoma. *Cancer Cell*. 2014;26(1):33-47.
31. Berezovsky AD, Poisson LM, Cherba D, Webb CP, Mikkelsen AD, deCarvalho AC. *Sox2* promotes malignancy in glioblastoma by regulating plasticity and astrocytic differentiation. *Neoplasia*. 2014;16(3):193-206.
32. Shivapathasundram G, Wickremesekera AC, Brasch HD, Marsh R, Tan ST, Itinteang T. Expression of embryonic stem cell markers on the microvessels of WHO grade I meningioma. *Front Surg*. 2018;(5):65.
33. Spoelstra NS, Manning NG, Higashi Y, Darling D, Singh, M, Shroyer KR, et al. The transcription factor ZEB1 is aberrantly expressed in aggressive uterine cancers. *Cancer Res*. 2006;66(7):3893-902.
34. Wellner U, Brabletz T, Keck T. ZEB1 in pancreatic cancer. *Cancers*. 2010;2(3):1617-28.
35. Shen A, Zhang Y, Yang H, Xu R, Huang G. Overexpression of ZEB1 relates to metastasis and invasion in osteosarcoma. *J Surg Oncol*. 2012;105(8):830-4.
36. Zhang J, Lu C, Zhang J, Kang J, Cao C, Li M. Involvement of ZEB1 and E-cadherin in the invasion of lung squamous cell carcinoma. *Mol Biol Rep*. 2013;40(2):949-56.
37. Zhou YM, Cao L, Li B, Zhang RX, Sui CJ, Yin ZF, et al. Clinicopathological significance of ZEB1 protein in patients with hepatocellular carcinoma. *Ann Surg Oncol*. 2012;19(5):1700-6.
38. Okugawa Y, Toiyama Y, Tanaka K, Matsusita K, Fujikawa H, Saigusa S, et al. Clinical significance of zinc finger E-box binding homeobox 1 (ZEB1) in human gastric cancer. *J Surg Oncol*. 2012;106(3):280-5.
39. Edwards LA, Kim S, Madany M, Nuno M, Thomas T, Li A, et al. ZEB1 is a transcription factor that is prognostic and predictive in diffuse gliomas. *Front Neurol*. 2019;9:1199.
40. Edwards LA, Li A, Berel D, Madany M, Kim N-H, Liu M, et al. ZEB1 regulates glioma stemness through LIF repression. *Sci Rep*. 2017;7(1):69.