

Clonal Diversity and Genetic Landscape of *mecC*-Positive *S. aureus* in Human Samples from Iran: First Report of CC/ST423/t742 in Tehran, Iran

Mehdi Goudarzi^{1*}, Sima Sadat Seyedjavadi², Mozhgan Raigani³, Zahra Salehi², Masoud Dadashi⁴, Mohammad Javad Nasiri¹

¹Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ²Department of Mycology, Pasteur Institute of Iran, Tehran, Iran; ³Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran; ⁴Department of Microbiology, School of Medicine, Alborz University of Medical Sciences, Karaj, Iran

OPEN ACCESS

Article type: Research Article

Received: September 22, 2025

Revised: December 7, 2025

Accepted: February 15, 2026

Published online: February 17, 2026

How to cite:

Mirfakhraie R, Hassani Fard Katiraei S, Mirnezami M, Bayat S, Soosanabadi M, Kookhaei A, Gholami M. Whole-Exome Sequencing Identified a Novel Mutation in an Iranian Patient with Epidermolysis Bullosa. *Iran. Biomed. J.* 2026; 30(2 & 3): 174-187.

ABSTRACT

Background: The emergence of *mecC*-carrying *Staphylococcus aureus* strains is a serious concern in public health. In this study, we investigated the clonal diversity, genetic characteristics, antimicrobial resistance, virulence determinants, biofilm-forming ability, and molecular epidemiology of *mecC*-positive *Staphylococcus aureus* isolates recovered from hospitalized patients in Iran.

Methods: In a descriptive study, 45 *mecC*-carrying *S. aureus* strains obtained from 1,500 *S. aureus* which were isolates from different clinical samples, were analyzed. The characterization included antimicrobial susceptibility tests, biofilm formation assessment, *spa* typing, multilocus sequence typing (MLST) and PCR analysis to detect resistance, and biofilm-related, and virulence genes.

Results: Our results showed that 17.8%, 17.8%, and 64.4% of the isolates were classified as weak, intermediate, and strong biofilm producers, respectively. The prevalence of the *ant(4')-Ia* (80%), and *tet(M)* (73.3%) genes was found in more than half of the isolates. MLST revealed that the 45 isolates belonged to four clonal complexes, including CC8 (44.4%), CC130 (24.5%), CC121 (17.8%), and CC425 (13.3%). The majority of *S. aureus* isolates belonged to CC8/ST239 (44.4%). *tst*-positive strains belonged to CC121/ST599/t5930 (17.8%) and CC8/ST239/t037 (11.1%), while *eta*-positive isolates belonged to CC/ST130/t843 (15.6%). Fusidic acid-resistant *S. aureus* isolates belonged to CC8/ST239/t037 (11.1%), followed by CC8/ST239/t7688 (6.7%) lineages, carrying *fusB*. Vancomycin-intermediate *S. aureus* strains belonged to CC/ST599/t5930 (6.7%) and CC/ST130/t843 (4.4%) clones.

Conclusion: The study highlights the need for close and continuous monitoring of the genetic diversity and characteristics of the *mecC*-carrying *S. aureus* strains. Besides, urgent measures are required to prevent the transmission of these strains to treat associated infections effectively.

DOI: 10.61882/ibj.5161

Keywords: Hospitals, Methicillin-resistant *Staphylococcus aureus*, *Staphylococcus aureus*



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

Corresponding Author: Mehdi Goudarzi

Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Koodak-yar St., Daneshjoo Blvd, Velenjak, Chamran HWY, Tehran, Iran; Tel.: (+98-21) 23872556; Fax: (+98-21) 23872556; E-mail: goudarzim@yahoo.com; ORCID ID: 0000-0001-6720-9341

1. INTRODUCTION

Staphylococcus aureus is a significant pathogen that contributes to both hospital-acquired and community-associated infections worldwide. It is implicated in a wide spectrum of clinical presentations, ranging from superficial skin infections to severe, life-threatening conditions^[1]. A critical challenge in managing *S. aureus* infections is the emergence and widespread dissemination of multidrug-resistant (MDR) strains, which consistently compromise therapeutic efficacy. Among these, methicillin-resistant *S. aureus* (MRSA) strains are responsible for the majority of *S. aureus*-related infections^[1,2].

Resistance in MRSA is primarily mediated by the *mecA* gene, which encodes an altered penicillin-binding protein (PBP2a) with reduced affinity for β -lactam antibiotics, rendering these treatments ineffective^[2]. Recently, a novel genetic determinant, initially identified as *mecALGA251* and now referred to as *mecC*, has been discovered^[3,4]. This gene encodes a transpeptidase and is transferable between livestock-associated MRSA or other *Staphylococci* species and human MRSA strains. Notably, *mecC* shares 63% nucleotide and 63% protein sequence identity with the conventional *mecA*/PBP2a^[3]. Consequently, this divergence renders *mecC* undetectable by standard *mecA*-specific PCR assays or PBP2a slide agglutination tests, often leading to misidentification as methicillin-sensitive *S. aureus*^[3]. Following its initial identification in Iran, *mecC*-positive MRSA has subsequently been documented in multiple European countries, including the United Kingdom, Denmark, Germany, France, and the Netherlands, as well as in regions beyond Europe, such as Australia and parts of Asia^[4-6]. These strains have been isolated from both human clinical specimens and various animal sources, particularly livestock and wildlife, underscoring their zoonotic nature and the significance of interspecies transmission in their spread^[4,6,9].

While the prevalence of *mecC*-positive MRSA is generally lower than that of *mecA*-mediated MRSA, growing attention has been directed toward their clinical importance due to diagnostic limitations and the risk of unrecognized circulation in healthcare environments. Although the emergence, epidemiology, and genetic variability of the *mecA* gene have been extensively studied, the prevalence and clinical implications of *mecC*-carrying MRSA isolates remain insufficiently understood^[3,5]. In Iran, information on *mecC*-positive *S. aureus* remains scarce, as the majority of surveillance and molecular studies have primarily addressed methicillin resistance associated with the *mecA* gene. Consequently, limited data are available regarding the occurrence, genetic features, and virulence-related traits of *mecC*-carrying *S. aureus* isolates in clinical

settings across the country. This lack of evidence underscores the necessity for focused investigation on enhancing epidemiological insight and improving diagnostic recognition of these emerging MRSA variants in the region.

2. MATERIAL AND METHODS

2.1. Sample collection

This study was conducted over eleven years from 2013 to 2024 in Tehran, Iran. *S. aureus* strains were isolated from various clinical samples (skin wounds, burn wounds, surgical wounds, blood, pus, and urine) referred to the clinical laboratories of hospitals affiliated with Shahid Beheshti University of Medical Sciences (SBMU), Tehran, Iran. The identification process involved catalase testing and coagulase testing. For additional confirmation, the isolates were subcultured on DNAse and mannitol-salt agar. To genetically confirm the isolates, we amplified the *nucA* gene using PCR. The forward primer sequence was 5'-GTGATGGTGATACGGTT, and the reverse primer was AGCCAAGCCTTGACGAACTAAAGC, as described previously^[8]. The PCR products were examined through electrophoresis on 1.5% agarose gels. The confirmed *S. aureus* isolates, supplemented with 20% glycerol, were stored at -70°C for further analysis, as described formerly^[8].

2.2. Screening for MRSA strains and *mecC* gene detection

MRSA strains were screened using the disk diffusion method with cefoxitin (30 μ g) disks on Mueller-Hinton agar (Merck, Germany), following the Clinical and Laboratory Standards Institute (CLSI) guidelines (2022). The presence of the *mecC* gene in all isolates was further investigated through PCR analysis, using forward (5'-GTCCCTAACAAAACACCCAAAGA-3') and reverse (5'-GAAGATCTTTTCGTTTTTCAGC-3') primers, as outlined before^[9].

2.3. Determination of isolate susceptibility

All *mecC*-carrying isolates were evaluated for the determination of minimum inhibitory concentration (MIC) values for various antibiotics using the microbroth dilution method, in accordance with the CLSI guidelines. The antibiotics tested included fusidic acid, ciprofloxacin, erythromycin, clindamycin, nitrofurantoin, gentamicin, penicillin, chloramphenicol, tetracycline, rifampin, vancomycin, and mupirocin (including both low- and high-level resistance)^[8]. Additionally, the D-zone test was conducted in accordance with the CLSI standards to further characterize resistance patterns. For quality control, reference strains of *S. aureus* (ATCC 29213, ATCC 25923, and ATCC 43300) were included in each experiment to ensure the reliability and accuracy of the testing procedures.

2.4. Detection of biofilm formation

The biofilm formation was evaluated using a microtiter plate assay as explained earlier^[8]. Biofilm formation by *mecC*-carrying strains was determined by measuring the optical density (OD) at 490 nm. The cutoff optical density was calculated by adding three times of the standard deviation (SD) to the mean OD of the blank control wells. Isolates were then categorized according to their biofilm-forming capacity as follows: non-biofilm producers ($OD \leq 0.059$), weak biofilm producers ($OD_c < OD \leq 2 \times OD_c$), moderate biofilm producers ($2 \times OD_c < OD \leq 4 \times OD_c$), and strong biofilm producers ($OD > 4 \times OD_c$). The *S. epidermidis* ATCC 35984 strain was used as the positive control in the biofilm formation assays.

2.5. DNA extraction, virulence, resistance, and biofilm genes

The phenol-chloroform assay was used for the extraction of chromosomal DNA as previously described^[8]. The detection of toxin-encoding genes (*pvl*, *etb*, *tst*, and *eta*) and resistance determinants (*mupA*, *msr(A)*, *vanA*, *erm(A)*, *fusA*, *fusB*, *msr(B)*, *vanB*, *mupB*, *erm(B)*, *erm(C)*, *fusC*, *tetL*, *tetK*, *tetO*, *tetM*, *aph* (3')-IIIa, *aac* (6')-Ie/aph (2''), *int* (4')-Ia) was performed using PCR. The *mecC*-carrying isolates were evaluated for the presence of *clfA*, *clfB*, *fnbA*, *fnbB*, *ebp*, *can*, and *lpp* using conventional PCR^[8,10,11]. Specific primers were designed for each target gene, and their sequences were verified using the BLAST tool available on the NCBI GenBank database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

2.6. Genotypic characterization *spa* typing

spa typing was performed to classify different types of *S. aureus* protein A, following the protocol established previously^[12]. Specific primers and optimized PCR conditions were used for the assay, with each PCR reaction including a negative control that replaced genomic DNA with water. Following amplification, the PCR products were purified using the QIAquick PCR Purification Kit (Toronto, ON, Canada). The purified products were then sequenced using an ABI Prism 377 automated sequencer (Applied Biosystems, Foster City, CA, USA). The obtained sequences were processed and edited with Chromas software (version 1.45, Australia). Finally, the sequences were submitted to the Ridom SpaServer database (<http://www.spaserver.ridom.de>) for comprehensive analysis and *spa* profile assignment.

2.7. Multilocus sequence typing (MLST)

MLST was performed on all *mecC*-carrying strains following the method described before^[13]. To determine

sequence types (STs), the sequences of seven housekeeping genes (*aroE*, *gmK*, *pta*, *tpiA*, *yqiL*, *arcC*, and *glpF*) were compared with the allelic profiles available in the MLST database (<https://pubmlst.org/>). Sequence type clustering was analyzed using the eBURST program (version 3), which identifies related STs and groups them into clonal complexes. The clonal complex for *S. aureus* was further verified by cross-referencing the ST with the PubMLST database.

2.8. Phylogenetic analysis

The bioinformatics data were thoroughly analyzed. The multilocus sequences (*AroE*, *AroE*, *GlpF*, *Gmk*, *Pta*, *Tpi*, and *YqiL*) were aligned using ClustalW on the CIPRES Science Gateway v.3.14^[14] platform. The final alignment comprised 2386 base pairs (bp) for the combined analysis of seven genes. Phylogenetic tree construction was performed using the maximum likelihood (ML) method implemented in the IQTREE v.1.9.6^[15]. The selection of the evolutionary model and ML phylogenetic analysis, with 10,000 ultrafast bootstrap replicates, was conducted on the IQTREE web server. The best-fitting evolutionary model was determined to be TPM2u+F+I, and the analysis was run with 10,000 bootstrap replicates (parameters set to '-alrt 10,000 -bb 10,000 -nt AUTO' with all other options set to default).

3. RESULTS

A total of 45 *mecC*-carrying *S. aureus* strains were obtained from 1,590 *S. aureus* isolates collected from different clinical samples. All *mecC*-carrying *S. aureus* isolates were confirmed to be MRSA. Among these isolates, 25 (55.6%) were obtained from female patients, and the remaining 20 (44.4%) were from male patients. A total of six different clinical sample types were collected in this study. Skin wound specimens were the most common, accounting for 33.3% of all samples, followed by burn wound samples (20.0%), surgical wound samples (15.6%), blood samples (13.3%), pus samples (11.1%), and urine samples (6.7%). Our findings showed that a low percentage of *mecC*-carrying strains (13.3%) were isolated from invasive infections, whereas the remaining isolates (86.7%) were related to non-invasive infections. The mean age of the patients was 40 years. Patients were divided into four age groups: 5 (11.1%) aged ≤ 20 years, 23 (51.1%) aged 21–45 years, 12 (26.7%) aged 46–65 years, and 5 (11.1%) aged ≥ 65 years.

According to our MIC results, the highest and lowest resistance rates among the 45 *mecC*-carrying *S. aureus* isolates were observed for penicillin (100%), and chloramphenicol (11.1%), respectively (Table 1). Notably, none of the tested isolates were resistant to vancomycin; however, five isolates had an MIC value

Table 1. Antimicrobial susceptibility patterns of *mecC*-carrying *S. aureus* isolated from different clinical samples, MIC (µg/ml)

Antibiotic	MIC (µg/ml)			N (%)
	Range	50%	90%	
Chloramphenicol	1-64	4	8	5 (11.1)
Ciprofloxacin	0.25-32	1	4	12 (26.7)
Clindamycin	0.25-64	4	4	24 (53.3)
Erythromycin	0.5-64	16	16	31 (68.9)
Fusidic acid	0.125-4	1	2	8 (17.8)
Gentamicin	1-128	32	32	40 (88.9)
Mupirocin	4-512	128	256	10 (22.2)
Nitrofurantoin	8-512	32	128	6 (13.3)
Penicillin	0.125-16	1	4	45 (100)
Rifampin	0.5-32	2	4	10 (22.2)
Tetracycline	0.5-128	16	32	42 (93.3)
Vancomycin	0.125-64	2	4	5 (11.1)

of 4 µg/mL, which confirmed them as vancomycin-intermediate *S. aureus* (VISA) isolates. Our results also showed that 10 isolates (22.2%) were resistant to mupirocin, and all exhibited high-level resistance, which was classified as high-level mupirocin-resistant *S. aureus* (HLMUPR). Overall, 88.9% of the *mecC*-carrying *S. aureus* isolates were classified as MDR, with MDR defined as resistance to three or more antimicrobial agents. According to our results, 24 isolates (53.3%) exhibited constitutive macrolide-lincosamide-streptogramin B (cMLS_B) phenotypes, and 7 (15.6%) isolates indicated inducible macrolide-lincosamide-streptogramin B (iMLS_B) phenotypes. The results of the micro broth dilution method indicated

12 distinct resistance profiles. The top three resistance profiles included resistance to clindamycin, erythromycin, gentamicin, penicillin, and tetracycline, which were observed in 24.4% (11/45) of the isolates. In addition, resistance to clindamycin, erythromycin, gentamicin, penicillin, and tetracycline was detected in 15.6% (7/45) of the isolates. Resistance patterns to ciprofloxacin, erythromycin, gentamicin, mupirocin, penicillin, and tetracycline, as well as to gentamicin, penicillin, and tetracycline, were observed in 11.1% (5/45) of the isolates. Details of the 12 simultaneous resistance patterns and the distribution of clinical samples are summarized in Table 2.

Table 2. Antibiotic resistance profiles of *mecC*-carrying *S. aureus* strains and their distribution in different samples

No. of antibiotics	Resistance profile	No. of isolates (%)	Type of samples (n; %)
Eight	CHO, ERY, FLU, GEN, MUP, NIT, PEN, TET	3 (6.7)	BW (3; 100)
Seven	CHO, ERY, GEN, MUP, PEN, RIF, TET	3 (6.7)	B (1; 33.3), SuW (1; 33.4), SW (1; 33.3)
	CLI, ERY, FUS, GEN, PEN, RIF, TET	2 (4.4)	SW (2; 100)
	CIP, CLI, ERY, FUS, GEN, PEN, TET	3 (6.7)	SuW (1; 33.3), SW (1; 33.4), BW (1; 33.3)
Six	CLI, ERY, GEN, PEN, TET, RIF	5 (11.1)	BW (2; 40), SuW (1; 20), B (1; 20), P (1; 20)
	CIP, ERY, GEN, MUP, PEN, TET	4 (8.9)	SW (2; 50), U (1; 25), B (1; 25)
	CIP, GEN, NIT, PEN, TET, VAN	3 (6.7)	SuW (1; 33.3), SW (1; 33.4), U (1; 33.3)
	CHO, CIP, GEN, PEN, TET, VAN	2 (4.4)	B (1; 50), U (1; 50)
Five	CLI, ERY, GEN, PEN, TET	11 (24.4)	SW (4; 36.4), BW (3; 27.2), SuW (2; 18.2), B (2; 18.2)
Three	GEN, PEN, TET	4 (8.9)	SW (2; 50), P (2; 50)
Two	PEN, TET	2 (4.4)	SuW (1; 50), SW (1; 50)
One	PEN	3 (6.7)	SW (1; 33.3), P (2; 66.7)

^aPEN: penicillin; ERY: erythromycin; GEN: gentamicin; TET: tetracycline; CLI: clindamycin; NIT: nitrofurantoin; CIP: ciprofloxacin; RIF: rifampin; MUP: mupirocin; VAN: vancomycin; CHO: chloramphenicol; B: blood; BW: burn wounds; SuW: surgical wounds; SW: skin wounds; P: pus; U: urine



Fig. 1. PCR products of antibiotic resistance and toxin genes. Lane 1: *tetO* gene; lane 2: *tetL* gene; lane 3: *tetK* gene; lane 4: *fusB* gene; lane 5: *ant(4')-Ia* gene; lane 6: *aac(6')-Ie/aph(2'')* gene; lane 7: *erm(B)* gene; lane 8: *erm(C)* gene; lane 9: *erm(A)* gene; lane 10: *msrB* gene; lane 11: *msrA* gene; lane 12: *tst* gene; lane 13: *eta* gene; lane 14: *tetM* gene; lane 15: *aph(3')-IIIa* gene; lane 16: *aph(3')-IIIa* gene; lane M: 100 bp DNA ladder (Thermo Fisher Scientific, UK) showing fragments of 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1200, and 1500 bp.

Antibiotic resistance gene screening indicated that of 45 tested isolates, 36 (80%) had *ant(4')-Ia* gene, while 33 (73.3%) *tet(M)*, 14 (31.1%) *erm(A)*, 12 (26.7%) *aac(6')-Ie/aph(2'')*, 10 (22.2%) *mupA*, 8 (17.8%) *fusB*, 8 (17.8%) *erm(B)*, 6 (13.3%) *tet(L)*, 5 (11.1%) *tet(K)*, 5 (11.1%) *msrB*, 4 (8.9%) *msrA*, 4 (8.9%) *tet(O)*, 3 isolates (6.7%) *tet(S)*, 3 isolates (6.7%) *tet(A)*, and 1 *aph(3')-IIIa* (2.2%), as shown in Figure 1. All isolates were found to carry the *mecC* gene. Regarding antibiotic resistance genes, 14 genes were detected. The three most frequently detected profiles were: *ant(4')-Ia*, *tet(M)*, *erm(A)* (11%; 5/45), *ant(4')-Ia*, *tet(M)*, *tet(K)*, *aac(6')-Ie/aph(2'')* (11.1%; 5/45), and *ant(4')-Ia*, *tet(M)* (11%; 5/45). All HLMUPR strains were found to carry the *mupA* gene, and none of the VISA isolates harbored the *vanA* gene.

The biofilm assay showed that all *mecC*-carrying *S. aureus* strains were biofilm producers. Among the 45 total isolates, 29 (64.4%) were classified as strong biofilm producers, while 8 (17.8%) were categorized as moderate producers, and 8 (17.8%) as weak biofilm producers. All *mecC*-carrying *S. aureus* strains with iMLSB phenotypes were strong biofilm producers (15.6%; 7/45). Of the 24 cMLSB isolates, 17 (70.8%) indicated strong biofilm production, 4 (16.7%) moderate ability, and 3 (12.5%) weak ability. All fusidic acid-resistant *S. aureus* isolates carrying the *mecC* gene, including those resistant to HLMUPR, exhibited strong ability in biofilm formation. The distribution of resistance patterns among different biofilm-producing isolates is presented in Table 3.

The prevalence of biofilm carriage genes was recorded as follows: *clfA* in all isolates (100%), *clfB* in 40 isolates (88.9%), *fnbA* in 37 isolates (82.2%), *fnbB* in 24 isolates (53.3%), *cna* in 23 isolates (51.1%), *bap* in 10 isolates (22.2%), and *ebp* in 5 isolates (11.1%). Six patterns of biofilm-related genes were detected. Among these genes, *clfA*, *clfB*, *fnbA*, *fnbB*, and *cna* had a higher frequency (26.7%; 12/45). Figure 2 indicates the presence of biofilm genes in *mecC*-carrying *S. aureus* strains. Toxigenic *mecC*-carrying *S. aureus* strains were detected in 20 isolates (44.4%), with the majority of *tst* gene (28.9%, 13/45), followed by the *eta* (15.6%, 7/45).

All isolates were analyzed using *spa* and MLST typing methods. In the present survey, five *spa* types and four STs were identified. The invasive *mecC*-carrying *S. aureus* strains were associated with *spa* types t742 (6.7%; 3/45) and t7688 (6.7%; 3/45). The TST-carrying isolates belonged to *spa* types t5930 and t037, accounting for 17.8% (8/45), and 11.1% (5/45), respectively. All the isolates carrying *eta* toxin belonged to *spa* type t843, representing 15.6% (7/45) of the total isolates. MLST analysis revealed four STs among *mecC*-carrying *S. aureus* strains: ST239 (44.4%, 20/45), ST130 (24.5%, 11/45), ST599 (17.8%, 8/45), and ST425 (13.3%, 6/45). These STs were categorized into clonal complexes: CC8 (44.4%), CC130 (24.5%), CC121 (17.8%), and CC425 (13.3%).

Table 3. Distribution of resistance patterns among biofilm-producing *mecC*-carrying *S. aureus* strains

Biofilm status	Phenotypic resistance (no.:%)	Number (%)
Strong producer	CLI, ERY, FUS, GEN, MUP, NIT, PEN, TET (3; 10.3)	29 (64.4)
	CIP, CLI, ERY, FUS, GEN, PEN, TET (3; 10.3)	
	CHO, ERY, GEN, MUP, PEN, RIF, TET (3; 10.3)	
	CLI, ERY, FUS, GEN, PEN, RIF, TET (2; 6.9)	
	CIP, ERY, GEN, MUP, PEN, TET (4; 13.9)	
	CHO, CIP, GEN, PEN, TET, VAN (2; 6.9)	
	CIP, GEN, NIT, PEN, TET, VAN (3; 10.3)	
	CLI, ERY, GEN, PEN, TET, RIF (3; 10.3)	
Moderate producer	CLI, ERY, GEN, PEN, TET, RIF (2; 25)	8 (17.8)
	CLI, ERY, GEN, PEN, TET (2; 25)	
	GEN, PEN, TET (3; 37.5)	
	PEN (1; 12.5)	
Weak producer	PEN (2; 25)	8 (17.8)
	PEN, TET (2; 25)	
	GEN, PEN, TET (3; 37.5)	
	CLI, ERY, GEN, PEN, TET (3; 10.3)	

PEN: penicillin; ERY: erythromycin; GEN: gentamicin; TET: tetracycline; CLI: clindamycin; NIT: nitrofurantoin; CIP: ciprofloxacin; RIF: rifampin; MUP: mupirocin; VAN: vancomycin; CHO: chloramphenicol

3.1. Analysis of *mecC*-positive clones CC8

In the present study, the CC8, corresponding to CC8/ST239/t7688 (33.3%; 15/45) and CC8/ST239/t037 (11.1%; 5/45), was the most prevalent clone among *mecC*-carrying *S. aureus* strains. The CC8/ST239-SCC*mec* XI strain was recovered from skin wounds (35%), burn wounds (20%), and surgical wounds, blood, and urine (each 5%). All the CC8 isolates were resistant to mupirocin and indicated the HLMUPR

phenotype carrying the *mupA* gene. Additionally, all the CC8/ST239/t037 isolates were found to harbor the *tst* gene. Of eight fusidic acid-resistant *S. aureus* isolates, three (37.5%; 3/8) elongated to the CC8/ST239/t7688 and five (62.5%; 5/8) belonged to CC8/ST239/t037, both of which carried the *fusB* gene. All CC8/ST239/t037 isolates were identified as strong biofilm producers. In contrast, among the CC8/ST239/t7688 isolates, 66.7% were strong biofilm




Fig. 2. Representative amplification of biofilm encoding genes from *mecC*-carrying *S. aureus* isolates. Lane M: 100-bp DNA ladder (Thermo Fisher Scientific, UK); lane 1: *clfA*-encoding gene; lane 2: *clfB*-encoding gene; lane 3: *fnbA*-encoding gene; lane 4: *fnbB*-encoding gene; lane 5: *bap*-encoding gene; lane 6: *can*-encoding gene; lane 7: *ebp*-encoding gene; lane 8: positive control of *clfB*-encoding gene; lane 9: positive control of *can*-encoding gene; lane 10: positive control of *ebp*-encoding gene; lane 11: negative control.

producers and 33.7% were weak biofilm producers. The present research indicated that the iMLS_B phenotype was more prevalent in CC8/ST239/t7688 strains compared to the cMLS_B phenotype (46.6% vs. 20%). Conversely, CC8/ST239/t037 strains displayed a 100% prevalence of the cMLS_B phenotype, with none exhibiting the iMLS_B phenotype. Regarding resistance genes, *ant(4')-Ia*, *tet(M)*, and *fusB* were detected in all the CC8/ST239/t037 isolates. More than three-fourths of the CC8/ST239/t7688 isolates carried *ant(4')-Ia* and *tet(M)* genes. The distribution of biofilm-related genes among the CC8/ST239 isolates was diverse. Figure 3 details the distribution of resistance profiles and biofilm genes among the ST239 clonal lineage.

3.2. CC130

The present survey showed that all the isolates belonging to the CC130 clone were classified as CC/ST130/t843. The CC/ST130-SCC_{mec} XI/t843 strain was isolated from burn wounds (27.3%), pus (27.3%), skin wounds (27.3%), and surgical wounds (18.1%). Also, two VISA isolates (40%; 2/5) were identified within this clone. The cMLS_B phenotype was detected in 81.8% of isolates (9/11). Notably, none of the CC130

isolates exhibited the iMLS_B phenotype. Resistance gene analysis showed the presence of *ant(4')-Ia* (81.8%), *tet(M)* (90.9%), *tet(K)* (27.3%), *tet(S)* (9.1%), *aac(6')-Ie/aph(2'')* (45.4%) among the tested isolates (Table 4). Virulence gene analysis indicated that the *eta* gene was found only in CC/ST130/t843 (63.6%; 7/11) isolates. According to our analysis, all the isolates exhibited strong biofilm production. The distribution of resistance profiles and biofilm-related genes is presented in Figure 4 (A and B). Surprisingly, we detected a relatively high prevalence of *bap* (36.4%) among these isolates.

3.3. CC121

The third most common *mecC*-positive *S. aureus* clone was ST599/t5930, which exhibited VISA in 37.5% of isolates (3/8). The CC/ST599-SCC_{mec} XI/t5930 strain was isolated from skin wounds (50%), burn wounds (25%), and surgical wounds (each 12.5%). Notably, none of these isolates exhibited the HLMUPR phenotype (Fig. 4C and 4D). All of these isolates carried the *tst* gene. Additionally, 37.5% of the CC121 isolates were high-level biofilm producers. More than three-fourths of the isolates harbored the *ant(4')-Ia*.




Fig. 3. Distribution of resistance profiles and biofilm encoding genes among CC8 lineages. (A) Left: distribution of phenotypic resistance patterns in CC8/ST239/t7688 isolates; right: distribution of biofilm genes in CC8/ST239/t7688 isolates; (B) left: distribution of phenotypic resistance patterns in CC8/ST239/t037 isolates; right: distribution of biofilm genes in CC8/ST239/t037 isolates.

Table 4. Characteristics of the 45 *mecC*-positive *S. aureus* strains obtained from clinical isolates

Clonal lineage	Toxin genes	Biofilm formation degree (n; %)	Genotype (n; %)	Adhesion genes (n; %)	Total n (%)
CC/ST130/t843	<i>eta</i> (7; 63.6)	Strong (11; 100)	<i>ant(4')-Ia</i> (9; 81.8), <i>tet(M)</i> (10; 90.9), <i>tet(K)</i> (3; 27.3), <i>tet(S)</i> (1; 9.1), <i>aac(6')-Ie/aph(2'')</i> (5; 45.4)	<i>clfA</i> (11; 100), <i>clfB</i> (11; 100), <i>fnbA</i> (9; 81.8), <i>fnbB</i> (7; 63.6), <i>cna</i> (9; 81.8), <i>bap</i> (4; 36.4)	11 (24.5)
CC8/ST239/t7688	-	Strong (10; 66.7), Weak (5; 33.3)	<i>ant(4')-Ia</i> (13; 86.7), <i>tet(M)</i> (13; 86.7), <i>fusB</i> (3; 20), <i>erm(A)</i> (6; 40), <i>mupA</i> (7; 46.7), <i>erm(C)</i> (3; 20), <i>erm(B)</i> (7; 46.7), <i>tet(O)</i> (2; 13.3), <i>aac(6')-Ie/aph(2'')</i> (2; 13.3), <i>msr(B)</i> (2; 13.3), <i>msrA</i> (3; 20)	<i>clfA</i> (15; 100), <i>clfB</i> (15; 100), <i>fnbA</i> (9; 60), <i>fnbB</i> (10; 66.7), <i>cna</i> (6; 40), <i>bap</i> (2; 13.3)	15 (33.3)
CC8/ST239/t037	<i>tst</i> (5; 100)	Strong (5; 100)	<i>ant(4')-Ia</i> (5; 100), <i>tet(M)</i> (5; 100), <i>mupA</i> (3; 60), <i>fusB</i> (5; 100), <i>erm(A)</i> (3; 60), <i>msr(B)</i> (2; 40)	<i>clfA</i> (5; 100), <i>clfB</i> (5; 100), <i>fnbA</i> (5; 100), <i>fnbB</i> (5; 100), <i>bap</i> (2; 40), <i>cna</i> (3; 60)	5 (11.1)
CC/ST425/t742	-	Moderate (5; 83.3), Weak (1; 16.7)	<i>ant(4')-Ia</i> (5; 100), <i>tet(M)</i> (6; 100), <i>tet(K)</i> (1; 16.7), <i>tet(O)</i> (1; 16.7), <i>aac(6')-Ie/aph(2'')</i> (3; 50), <i>erm(B)</i> (1; 16.7), <i>msr(A)</i> (1; 16.7), <i>erm(A)</i> (3; 50)	<i>clfA</i> (5; 100), <i>fnbA</i> (6; 100), <i>fnbB</i> (4; 66.7), <i>ebp</i> (2; 33.3), <i>cna</i> (2; 33.3)	6 (13.3)
CC121/ST599/t5930	<i>tst</i> (8; 100)	Strong (3; 37.5), Moderate (2; 37.5), Weak (2; 25)	<i>ant(4')-Ia</i> (6; 75), <i>tet(M)</i> (4; 50), <i>tet(K)</i> (2; 25), <i>tet(O)</i> (1; 12.5), <i>tet(S)</i> (2; 25), <i>aac(6')-Ie/aph(2'')</i> (3; 37.5), <i>msr(B)</i> (1; 12.5), <i>erm(A)</i> (2; 25), Without resistance gene (1; 12.5)	<i>clfA</i> (8; 100), <i>clfB</i> (5; 62.5), <i>fnbA</i> (8; 100), <i>fnbB</i> (2; 25), <i>bap</i> (2; 25), <i>ebp</i> (3; 37.5), <i>cna</i> (3; 37.5)	8 (17.8)

Biofilm-related genes detected in isolates of this clone included *clfA* (100%), *fnbA* (100%), *clfB* (62.5%), *fnbB* (25%), *bap* (25%), *ebp* (37.5%), and *cna* (37.5%), as shown in Figure 5. Table 5 shows details of different clonal lineages based on *spa* and MLST typing methods.

3.4. CC425

Another clone identified in *mecC*-carrying *S. aureus* strains was CC/ST425/t742 (13.3%). The CC/ST425-SCC*mec* XI/t742 strain was recovered from blood (50%), as well as surgical wounds, skin wounds, and pus (16.7% each). More than three-fourths (75%) of the isolates were moderate biofilm producers. The most prevalent resistance genes including *tet(M)* (100%), *ant(4')-Ia* (50%), and *erm(A)* (50%). A phylogenetic tree based on seven combined loci is presented in Figure 4 (E and F). The tree was constructed using the

RAXML method with a dataset of 3,186 bp. The phylogenetic analysis revealed that all STs were classified into four well-supported clades. The ST599 clade occupied a basal position relative to the other STs (Fig. 5). The ST239 clade formed a highly supported group together with ST425, and they were placed as sister taxa in the ML tree. Notably, ST239 included two strains that showed different genetic resistance patterns.

4. DISCUSSION

MRSA infections are widespread in both healthcare settings and the community. The *mecC* gene has emerged as a novel determinant of methicillin resistance in *Staphylococcus* species^[3]. This study investigates the molecular epidemiology and genetic characterization of clinical *mecC*-carrying *S. aureus* strains isolated from selected hospitals in Tehran,



Fig. 4. Distribution of resistance profiles and biofilm encoding genes among CC lineages. Distribution of (A) resistance profiles and (B) biofilm encoding genes among CC130 lineages. Distribution of (C) resistance profiles and (D) of biofilm-encoding genes among CC121/ST599/t5930 isolates. Distribution of (E) resistance profiles and (F) biofilm-encoding genes among CC/ST425/t742 isolates.

providing novel insights into circulating *mecC*-carrying *S. aureus* strains previously unreported in Iran. The findings include the first report on the prevalence and genetic diversity of *mecC*-positive *S. aureus* isolates in clinical samples from Iran. Notably, *mecC*-positive *S. aureus* strains predominantly exhibited strong biofilm-forming capabilities. Additionally, a high prevalence of the CC8 clonal complex was observed among these isolates, highlighting their significant role in infections and outbreaks in healthcare settings.

The *mecC* gene has been detected in *S. aureus* isolates from both humans and dairy cattle^[3]. Studies have

indicated an increasing prevalence of *mecC*-carrying *S. aureus*^[3,14]. In our research, a 2.8% prevalence of *mecC*-carrying *S. aureus* isolated from clinical samples was found, with all being MRSA. Over the past two decades, the prevalence of *mecC* among *S. aureus* in Iran has increased from 0.6% to 0.9%^[8,15]. In 2015, a systematic review and meta-analysis study by Diaz et al. showed a low level of *mecC*-carrying *S. aureus* strains originated from human samples (0.004%; 95% CI = 0.002-0.007) but also highlighted a higher prevalence in animal samples (0.10%; 95% CI = 0.033-0.174)^[14]. Research from different geographic area, such as Pakistan




Fig. 5. A phylogenetic tree generated through ML analysis of four sequence types (ST 139, ST 239, ST 425, and ST 599) showing a combination of seven loci (*ArcC, AroE, GlpF, Gmk, Pta, Tpi, and YqiL*). The analysis was conducted with the IQ-TREE software applying the TPM2u+F+I model and performing 10,000 ultrafast bootstrap replicates. Bootstrap values greater than 95% are indicated on the tree. ST 239 included two strains that showed different genetic resistance patterns.

(96.8%), Egypt (6%), UK (0.45%), Denmark (5.9%), Germany (0.9%), Belgium (0.18%), Spain (15%), Austria (2%), Finland (0.74%), and Slovenia (5%), reported various prevalence rates of *mecC*-carrying *S. aureus* isolated from human samples, indicating a wide geographical spread of these strains [1, 17]. Notably, our findings revealed a concerning rise in the prevalence of *mecC*-carrying *S. aureus* in clinical settings in recent years [18]. However, *mecC* remained more prevalent in Iran than in several other European and some Asian countries [5, 19-21]. Although the origin of *mecC*-carrying *S. aureus* remains unclear, earlier evidence suggests that horizontal gene transfer from non-*aureus* *Staphylococcus* species to *S. aureus* may have significantly contributed to the increasing incidence and prevalence of *mecC*-mediated resistance in clinical settings in recent years [3]. As a result, clinical microbiologists should remain vigilant about the potential transfer of *mecC* from other methicillin-resistant *Staphylococcus* pathogens [3, 14, 19].

Earlier research has indicated that livestock serves as a reservoir of *mecC*-carrying *S. aureus* and a major source of zoonotic infections in humans [19]. It seems reasonable to attribute the high prevalence rate of *mecC* in the present study to contact with livestock and the growing trend of keeping pets, which could facilitate the transmission of *mecC*-carrying *S. aureus* strains from animals to humans. In our study, MRSA isolates exhibited high resistance to erythromycin, penicillin, tetracycline, gentamicin, and clindamycin and

appeared to be multidrug-resistant, consistent with previous reports from this region [8, 15]. We showed the resistance profile of clinical *mecC*-carrying *S. aureus* isolates for various antibiotics. These findings suggest either an overuse of antibiotics in human and veterinary medicine or potential evolutionary adaptations of the bacteria. Notably, intermediate resistance to vancomycin has already emerged in our clinical settings, with 11.1% of isolates. This is a concerning development, as the emergence of intermediate or resistant strains in this region poses a significant threat, given that vancomycin remains the primary treatment for MRSA infections. A systematic review and meta-analysis study conducted by Shariati et al. from 1997 to 2019 indicated a 3.6-fold increase in the prevalence of VISA strains from 1.2% before 2010 to 4.3% after 2010 [22]. However, the increase in VISA strains may be due to several genetic and phenotypic alterations in wild-type bacteria, changes in the thickness of the cell wall, easy availability of these strains, and the lack of guidelines for antibiotic use. Evidence also shows an increasing trend in resistance to fusidic acid in MRSA strains. In a study performed in Kuwait, Udo and colleagues reported an increasing trend of resistance to fusidic acid from 22% in 1994 to 92% in 2004 [23]. In contrast to earlier surveys conducted in Iran that indicated resistance rates of 2.5%-8% for fusidic acid [11, 24], our study showed a prevalence rate of 17.8% for fusidic acid-resistant isolates, all of which carried the *fusB* gene. Consistent with other research [25], our

study highlights a relatively high prevalence of resistance to fusidic acid in MRSA strains. This increase reflects a broader shift in antibiotic usage patterns and a growing prevalence of staphylococcal infections, underscoring the urgent need for revised treatment protocols and enhanced antimicrobial stewardship to combat rising resistance.

In the current survey, 15.6% of the isolates indicated the iMLSb phenotype. The presence of the iMLSb phenotype among *mecC*-carrying *S. aureus* strains has been reported by several researchers in the UK [6], Iran^[15], India^[26], and Pakistan^[27]. Our findings, along with previous studies, support the widely accepted belief that the emergence of widespread resistance in clinical pathogens is driven by antibiotic use in human and veterinary medicine. The *mecC*-positive CC8/ST239/t7688 strain, which accounted for 33.3% of the isolates, has previously been reported in Iran^[18] and other parts of Asia^[28]. Similar patterns have been observed in Iran and other countries, including Switzerland, where this strain was detected on the hands of healthcare workers and in hospital environments. Additionally, it has been identified as a successful pathogen responsible for infections at various hospital sites in Iran^[29]. Evidence suggests that t7688/ST239 is a widely distributed endemic clone across many Asian countries^[18,23]. The presence of multiple resistance genes and a multidrug profile underscores the need for close monitoring of the CC8/ST239/t7688 strain.

Our analysis indicated that more than three-fourths of the CC8/ST239/t7688 isolates carried *ant(4')-Ia* and *tet(M)* genes. Similar findings have been observed in a study conducted by Ghasemzadeh-Moghaddam et al. in Iran, which reported the presence of resistant genes in CC8/ST239/t7688 lineages^[18]. Consistent with previous studies in Iran that reported the presence of CC8/ST239-t7688 strains^[18], our findings confirmed that all *mecC*-positive CC8/ST239/t7688 isolates were MRSA. In this study, we identified CC8/ST239/t037 in 11.1% of *mecC*-positive strains. The persistence of ST239 in hospital settings, particularly in Iran, is a significant concern due to its multidrug resistance and rapid dissemination. All isolates exhibited strong biofilm formation, resistance to multiple antibiotic classes, and resistance to fusidic acid. In agreement with our results, an Iranian study on 308 *S. aureus* reported the presence of ST239/t037 in *mecC*-positive *S. aureus*^[15]. However, previous reports have documented the emergence of this clone in China, Kuwait, and Saudi Arabia^[30,31], further supporting the hypothesis that it may have been introduced from neighboring countries. An earlier study by Goudarzi et al. in Iran reported fusidic acid-resistant t030 strains^[10]. Similarly, between

2010 and 2013, spa type t030 carrying the *fusB* gene was detected in 7.1% of MRSA isolates from clinical samples in a hospital in China^[32]. In the present study, spa type t037 (7.6%) emerged as the most prevalent genotype among fusidic acid-resistant *mecC*-positive strains, with a 100% prevalence of *fusB* carriage. These findings are consistent with earlier research by Chen et al. in Taiwan, which identified t037 (62%) and t002 (29%) as the most common genotypes among fusidic acid-resistant MRSA isolates from hospitalized patients^[33].

The second most common *mecC*-positive clone was CC/ST130/t843, characterized by the presence of multiple resistance genes, strong biofilm production, and the ETA toxin. A study by Lindstedt et al. in Sweden, in which 45 *mecC*-positive *S. aureus* isolates from patients over a year were analyzed, found that 76% of *mecC*-positive *S. aureus* belonged to spa types t373 and t843. This study also revealed a decline in the incidence of *mecC*-positive *S. aureus* over the past three years, dropping from 9.8% in 2014 to 3.4% in the same year^[34]. Similarly, research by Petersen et al. in Denmark, conducted between 2007 and 2011, identified CC30 (87.5%) and CC2361 (12.5%) as the predominant clones among *mecC*-positive *S. aureus* isolates, with no virulence genes detected^[4]. These findings align with those of García-Álvarez et al., who reported that CC130 *mecC* MRSA is the most prevalent lineage among human isolates in the UK and Denmark^[9]. Furthermore, a study by Dermota et al. analyzing 395 MRSA strains isolated from various clinical samples between 2006 and 2013 identified only six MRSA isolates carrying the *mecC* gene, all of which belonged to CC/ST130. In contrast, Dermota's research in Slovenia indicated that none of the *mecC*-positive MRSA isolates carried antibiotic resistance genes, and all were classified as CC/ST130^[35]. Although earlier epidemiologic research has infrequently reported *mecC*-carrying *S. aureus* isolates associated with ST599, we identified a *tst*-positive ST599 isolate that harbors the *mecC* gene and is characterized by spa type 5930. Previous studies have documented the presence of CC/ST599 isolates in Iran and many European countries, including such as Germany, Austria, the UK, and Belgium^[3,8,20]. Our observations regarding the *tst* carriage by ST599 isolates are comparable with the results obtained by Goudarzi et al. from Iran^[8], Sabat et al. from France^[36], and Kerschner et al. from Austria^[20].

In the present study, we identified the *mecC*-positive *S. aureus* clone ST425/t742, which belonged to CC425. This finding is supported by a study by Silva et al. on *S. aureus* isolates recovered from 78 surface water samples in Portugal. Their investigation into the clonal

diversity of *mecC*-positive *S. aureus* strains reported the presence of CC/ST425^[37]. However, in contrast to our findings, they indicated that CC425 isolates were more prevalent than CC130 isolates in surface waters. Our results are also in line with the observations of Porrero et al., who showed that the CC425 lineage was the most common among *mecC*-MRSA strains isolated from animals and Urban Wastewater^[7]. Our *mecC*-carrying *S. aureus* strains were negative for toxin-encoding genes, which is in accordance with Silva's study.

5. CONCLUSION

The present study describes the variability of clonal lineages of *mecC*-carrying *S. aureus* isolated from different clinical sources, including the first detection of CC/ST423/t742 in Iran. The identification of livestock-associated *S. aureus* lineages in healthcare settings underscores the zoonotic potential of this emerging resistant microorganism, along with other related animal-derived clades, which pose a significant risk to human health. These findings underscore the importance of proper and rational clinical use of antibiotics in humans and livestock, which can contribute to the evolution of certain *mecC*-carrying *S. aureus* lineages. The widespread multidrug resistance observed in *mecC*-carrying *S. aureus* strains, especially in healthcare settings, highlights the need for continuous surveillance and robust antimicrobial management. Monitoring *mecC*-positive *S. aureus* strains in animal veterinary practices, and food safety laboratories is crucial and should be included in preventive strategies to prevent potential health risks. Therefore, studying the detection and genetic diversity of *mecC*-positive *S. aureus* using robust phenotypic and molecular screening assays would be highly beneficial.

DECLARATIONS

Acknowledgments

Not applicable.

Generative AI and AI-assisted technologies

In this study, no artificial intelligence technology was used in the production of the submitted work.

Ethical approval

All the experimental procedures in this study were approved by the Research Ethics Committee of Shahid Beheshti University of Medical Sciences, Tehran, Iran (ethical code: R.SBMU.MSP.REC.1402.116.).

Consent to participate

All participants voluntarily agreed to participate in this study. Written informed consent was obtained from all participants prior to enrollment in the study.

Consent for publication

All authors reviewed the results and approved the final version of the manuscript.

Authors' contributions

MG: conceptualization, funding acquisition, methodology, project administration, resources, supervision, validation, writing—original draft, writing—review and editing. SSS: data curation, formal analysis, methodology, writing original draft, writing—review and editing. MR: data curation, formal analysis, methodology, writing original draft, writing—review and editing. ZS: formal analysis, writing original draft, writing—review and editing. Masoumeh Navidinia: investigation and curation, writing—original draft. MD: formal analysis, investigation, methodology, project administration. MJN: formal analysis, writing original draft.

Data availability

All relevant data can be found within the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was financially supported by a research grant from Deputy of Research, School of Medicine, Shahid Beheshti University of Medical Sciences, Iran (Grant No. 43020470). The funding agency has no role in the design of the project, work execution, analysis, interpretation of the data, and manuscript writing or submission.

Supplementary information

The online version does not contain supplementary material.

REFERENCES

1. Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler Jr VG. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin Microbiol Rev 2015;28:603-61.
2. Lee AS, De Lencastre H, Garau J, Kluytmans J, Malhotra-Kumar S, Peschel A, et al. Methicillin-resistant *Staphylococcus aureus*. Nat Rev Dis Primers. 2018;4:1-23

3. Paterson GK, Harrison EM, Holmes MA. The emergence of *mecC* methicillin-resistant *Staphylococcus aureus*. Trends Microbiol. 2014;22:42-7
4. Petersen A, Stegger M, Heltberg O, Christensen J, Zeuthen A, Knudsen L, et al. Epidemiology of methicillin-resistant *Staphylococcus aureus* carrying the novel *mecC* gene in Denmark corroborates a zoonotic reservoir with transmission to humans. Clin Microbiol Infect. 2013;19:E16-E22.
5. Lozano C, Fernández-Fernández R, Ruiz-Ripa L, Gómez P, Zarazaga M, Torres C. Human *mecC*-carrying MRSA: Clinical implications and risk factors. Microorganisms. 2020;8:1615.
6. Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler Jr VG. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. Clin Microbiol Rev 2015;28:603-61
7. Paterson G, Morgan F, Harrison E, Cartwright E, Török M, Zadoks R, et al. Prevalence and characterization of human *mecC* methicillin-resistant *Staphylococcus aureus* isolates in England. J Antimicrob Chemother. 2014;69:907-10.
8. Porrero MC, Valverde A, Fernández-Llarió P, Díez-Guerrero A, Mateos A, Lavin S, et al. *Staphylococcus aureus* carrying *mecC* gene in animals and urban wastewater, Spain. Emerg Infect Dis. 2011;17:1003-10.
9. Goudarzi M, Navidini M, Hashemi A, Pouriran R. First report of methicillin-resistant *Staphylococcus aureus* carrying the *mecC* gene in human samples from Iran: prevalence and molecular characteristics. New Microbes New Infect. 2021;39:100832
10. García-Álvarez L, Holden MT, Lindsay H, Webb CR, Brown DF, Curran MD, et al. Methicillin-resistant *Staphylococcus aureus* with a novel *mecA* homologue in human and bovine populations in the UK and Denmark: a descriptive study. Lancet Infect Dis. 2011;11:595-603
11. Goudarzi M, Kobayashi N, Dadashi M, Pantůček R, Nasiri MJ, Fazeli M, et al. Prevalence, genetic diversity, and temporary shifts of inducible clindamycin resistance *Staphylococcus aureus* clones in Tehran, Iran: a molecular-epidemiological analysis from 2013 to 2018. Front Microbiol. 2020;11:663
12. Goudarzi M, Tayebi Z, Dadashi M, Miri M, Amirpour A, Fazeli M. Characteristics of community-acquired methicillin-resistant *Staphylococcus aureus* associated with wound infections in Tehran, Iran: High prevalence of PVL+ t008 and the emergence of new *spa* types t657, t5348, and t437 in Iran. Gene Rep. 2020;19:100603.
13. Harmsen D, Claus H, Witte W, Rothganger J, Claus H, Turnwald D, et al. Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. J Clin Microbiol. 2003;41:5442-8
14. Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. J Clin Microbiol. 2000;38:1008-15
15. Diaz R, Ramalheira E, Afreixo V, Gago B. Methicillin-resistant *Staphylococcus aureus* carrying the new *mecC* gene—a meta-analysis. Diagn Microbiol Infect Dis. 2016;84:135-40
16. Moghaddam PZ, Azimian A, Sepahy AA, Iranbakhsh A. Isolation and genetic characterization of vancomycin-resistant and *mecC*+ methicillin-resistant *Staphylococcus aureus* strains in clinical samples of Bojnurd, Northeastern Iran. Jundishapur J Microbiol. 2021;14.
17. Idrees MM, Saeed K, Shabir M, Akhtar M, Qammar K, Hassan J, et al. Evaluation of *mecC* and *mecC*-associated methicillin-resistant *Staphylococcus aureus* in clinical specimens. Punjab, Pakistan. Biomedicines. 2023;11:878
18. Kerschner H, Kerschner J, Gardete S, Holmes MA, Holden MT, Apfalter P, et al. Properties of a novel PBP2A protein homolog from *Staphylococcus aureus* strain LGA251 and its contribution to the β -lactam-resistant phenotype. J Biol Chem. 2012;287:36854-63
19. Ghasemzadeh-Moghaddam H, Azimian A, Bayani G, Dashti V, Nojoomi S, Shirazi N, et al. High prevalence and expression of antiseptic resistance genes among infectious t037/ST239 methicillin-resistant *Staphylococcus aureus* (MRSA) strains in North Khorasan Province, Iran. Iran J Basic Med Sci. 2022;25:775
20. Larsen J, Raisen CL, Ba X, Sadgrove NJ, Padilla-González GF, Simmonds MS, et al. Emergence of methicillin resistance predates the clinical use of antibiotics. Nature. 2022;602:135-41
21. Kerschner H, Harrison EM, Hartl R, Holmes MA, Apfalter P. First report of *mecC* MRSA in human samples from Austria: molecular characteristics and clinical data. New Microbes New Infect. 2015;3:4-9
22. Lindgren A-K, Gustafsson E, Petersson A, Melander E. Methicillin-resistant *Staphylococcus aureus* with *mecC*: a description of 45 human cases in southern Sweden. Eur J Clin Microbiol Infect Dis 2016;35:971-5
23. Shariati A, Dadashi M, Moghadam MT, van Belkum A, Yaslianifard S, Darban-Sarokhalil D. Global prevalence and distribution of vancomycin resistant, vancomycin intermediate and heterogeneously vancomycin intermediate *Staphylococcus aureus* clinical isolates: a systematic review and meta-analysis. Sci Rep. 2020;10:12689.
24. Udo EE, Al-Sweih N, Mokaddas E, Johnny M, Dhar R, Gomaa HH, et al. Antibacterial resistance and their genetic location in MRSA isolated in Kuwait hospitals, 1994-2004. BMC Infect Dis. 2006;6:1-8
25. Zamani S, Mohammadi A, Hajikhani B, Abiri P, Fazeli M, Nasiri MJ, et al. Mupirocin-Resistant *Staphylococcus aureus* in Iran: A Biofilm Production and Genetic Characteristics. Biomed Res Int. 2022;2022:7408029
26. Hajikhani B, Goudarzi M, Kakavandi S, Amini S, Zamani S, van Belkum A, et al. The global prevalence of fusidic acid resistance in clinical isolates of *Staphylococcus aureus*: a systematic review and meta-analysis.

- Antimicrob Resist Infect Control. 2021;10:75.
26. Bali N, Borkakoty B, BaSHir H, Nazir S, WANI S, MIR A, et al. Isolation of *mecC* Gene carrying Methicillin Resistant *Staphylococcus aureus* in Clinical Samples from a Tertiary Care Institute, Northern India. *J Clin Diagn Res.* 2021;15
 27. Khan AA, Ali A, Tharmalingam N, Mylonakis E, Zahra R. First report of *mecC* gene in clinical methicillin resistant *S. aureus* (MRSA) from tertiary care hospital Islamabad, Pakistan. *J Infect Public Health.* 2020;13:1501-7
 28. Chen C-J, Huang Y-C. New epidemiology of *Staphylococcus aureus* infection in Asia. *Clin Microbiol Infect.* 2014;20:605-23
 29. Mirzaii M, Emaneini M, Jabalameli F, Halimi S, Taherikalani M. Molecular investigation of *Staphylococcus aureus* isolated from the patient, personnel, air and environment of an ICU in a hospital Tehran. *J Infect Public Health.* 2015;8:202-7
 30. Monecke S, Coombs G, Shore AC, Coleman M, Sampaio P, Borg M, et al. A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PloS one.* 2011;6:e17922
 31. Boswihi SS, Udo EE, Al-Sweini M. Shifts in the clonal distribution of methicillin-resistant *Staphylococcus aureus* in two hospitals: 1992-2010. *PloS one.* 2016;11:e0157443
 32. Yu F, Liu Y, Li C, Guo J, Li X, Ding Y, et al. Dissemination of fusidic acid resistance among *Staphylococcus aureus* clinical isolates. *BMC Microbiol.* 2015;15:1-6
 33. Chen H-J, Hung W-C, Tseng S-P, Tsai J-C, Hsueh P-R, Teng L-J. Fusidic acid resistance determinants in *Staphylococcus aureus* clinical isolates. *Antimicrob Agents Chemother.* 2010;54:4985-91
 34. García-Garrote F, Cercenado E, Marín M, Bal M, Trincado P, Corredoira J, et al. Methicillin-resistant *Staphylococcus aureus* carrying the *mecC* gene: emergence in Spain and report of a fatal case of bacteraemia. *J Antimicrob Chemother.* 2014;58:549-56
 35. Džurina Z, Zelená I, Černelj I, Grmek-Kosnik I, Ribic H, Ponišvar J, et al. Detection of methicillin-resistant *Staphylococcus aureus* carrying the *mecC* gene in clinical samples in Slovenia. *Epidemiol Infect.* 2015;143:1105-8
 36. Ghahat AJ, Koksai M, Akkerboom V, Monecke S, Kriegeskorte A, Hendrix R, et al. Detection of new methicillin-resistant *Staphylococcus aureus* strains that carry a novel genetic homologue and important virulence determinants. *J Clin Microbiol.* 2012;50:3374-7
 37. Silva V, Ferreira E, Manageiro V, Reis L, Tejedor-Junco MT, Sampaio A, et al. Distribution and clonal diversity of *Staphylococcus aureus* and other staphylococci in surface waters: Detection of ST425-t742 and ST130-t843 *mec C*-positive MRSA strains. *Antibiotics.* 2021;10:1416

COPY EDITING