



DECEMBER 11-12, 2025
۲۱ و ۲۰ آذر ماه ۱۴۰۴



دومین کنگره
پلازما پزشکی ایران

The 2nd Congress on Plasma Medicine

دبیرخانه دائمی کنگره
پلازما پزشکی ایران
www.plasmamedsym.ir



Effects of Cold Plasma on the Survival of Cancer Cells in vitro—A Systematic Review of Recent Experimental Findings

Erfan Ghane^{1*}, Fatemeh Norozian³, Mohammad Haqparast¹, Lily Darvish¹, Ali Alizadeh², Hasti Ghane³

¹Department of Radiology, Faculty of Paramedicine, Hormozgan University of Medical Sciences, Hormozgan, Iran

²General Director of Health Insurance, Hormozgan, Iran

³Department of Medicine, Bandar Abbas University of Medical Sciences, Bandar Abbas, Iran

OPEN ACCESS

Citation:

Ghane E, Norozian F, Haqparast M, Darvish L, Alizadeh A, Ghane H. Effects of Cold Plasma on the Survival of Cancer Cells in vitro—A Systematic Review of Recent Experimental Findings. *Iran Biomed J. Supplementary* (2026): 36.

ABSTRACT

Introduction: Cold atmospheric plasma (CAP) has emerged as a promising non-thermal oncological tool due to its ability to generate reactive oxygen and nitrogen species, UV photons, charged particles, and transient electric fields. These components interact synergistically with cancer cells, inducing oxidative stress, activating cell death pathways, and uncovering tumor-specific vulnerabilities. Recent preclinical studies demonstrate that both direct exposure to CAP and the use of plasma-activated liquids suppress proliferation, enhance apoptosis and ferroptosis, and selectively damage malignant cells while sparing most healthy tissues.

Materials and Methods: CAP triggers the accumulation of reactive oxygen species (ROS), lipid peroxidation, DNA damage, cytochrome c release, and caspase-dependent apoptosis. Also, ferroptosis has been reported to be modulated by iron metabolism and lipid-ROS pathways. The weaker antioxidant defense and altered redox homeostasis in cancer cells contribute to their selective vulnerability to these processes.

Results and Discussion: Our results indicated that both CAP and plasma-activated media reduced cell viability in hepatocellular, breast, cervical, colon, and lung cancer models. CAP modulated the USP49/HDAC3 signaling pathway, activated p53 pathways, and enhanced apoptotic markers. Additionally, the combination of CAP with chemotherapy improved drug sensitivity and overcome chemoresistance. In vivo studies further confirmed reduced tumor volume, decreased microvessel density, and enhanced vascular permeability. Importantly, the safety and feasibility of CAP treatment was affirmed in phase I clinical trials.

Conclusion: Current experimental and translational findings support CAP as a selective, multi-mechanistic, and minimally invasive anticancer strategy. Its ability to induce apoptosis, ferroptosis, and immune-stimulatory responses highlights its potential as an adjunct or standalone therapeutic tool. Future research should focus on device standardization, dose-response optimization, mechanistic modeling, and controlled clinical trials to fully establish CAP as a clinical cancer therapy modality.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

Keywords: Apoptosis, Cold atmospheric plasma, Ferroptosis, Oxidative stress, Plasma-activated medium

Corresponding Author: Erfan Ghane

Department of Radiology, Faculty of Paramedicine, Hormozgan University of Medical Sciences, Hormozgan, Iran

E-mail: hane.erfan1995@gmail.com



Iranian Biomedical Journal Supplementary (February 2026): 36