



DECEMBER 11-12, 2025  
۲۱ و ۲۰ آذر ماه ۱۴۰۴



دومین کنگره  
پلازما پزشکی ایران

The 2<sup>nd</sup> Congress on Plasma Medicine

دبیرخانه دائمی کنگره  
پلازما پزشکی ایران  
www.plasmamedsym.ir



# Differential Susceptibility of Terbinafine-Resistant and Sensitive Isolates of the *T. mentagrophytes* Complex to Cold Atmospheric Plasma

Fatemeh Hemmatyar<sup>1</sup>, Zahra Salehi<sup>2</sup>, Seyed Mohammad Atyabi<sup>3\*</sup>

<sup>1</sup>Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran

<sup>2</sup>Department of Mycology, Pasteur Institute of Iran, Tehran, Iran

<sup>3</sup>Department of Nanobiotechnology, Pasteur Institute of Iran, Tehran, Iran

## OPEN ACCESS

### Citation:

Hemmatyar F, Salehi Z, Atyabi SM. Differential Susceptibility of Terbinafine-Resistant and Sensitive Isolates of the *T. mentagrophytes* Complex to Cold Atmospheric Plasma. *Iran Biomed J. Supplementary* (2-2026): 38.

## ABSTRACT

**Introduction:** *Trichophyton mentagrophytes* is one of the most common dermatophytes in Iran and a major cause of tinea corporis, tinea cruris, and onychomycosis. The increasing emergence of terbinafine-resistant isolates has led to considerable therapeutic challenges and highlighted the need for alternative therapies. Cold atmospheric plasma (CAP) has recently been introduced as a promising antifungal approach. This study reports the effectiveness of CAP against both terbinafine-resistant and -sensitive isolates of the *T. mentagrophytes* complex.

**Materials and Methods:** Fifty clinical isolates and one reference susceptible strain (PFCC 5809) were studied. Terbinafine susceptibility was determined using the CLSI M38-A3 broth microdilution method, defining resistance as minimum inhibitory concentrations greater than 0.25 µg/mL. The resistant and susceptible isolates were subjected to CAP for 180 and 210 seconds. Growth inhibition was evaluated using mycelial dry weight in Sabouraud maltose broth and colony-forming unit (CFU) on Sabouraud dextrose agar. All studies were conducted in duplicate, with statistical significance set at  $p < 0.05$ .

**Results and Discussion:** Exposure to CAP decreased the growth of fungi in both resistant and susceptible isolates; however, the resistant strains showed greater inhibition. The most significant reduction was observed after 210 seconds of treatment, supported by dry weight and CFU analysis.

**Conclusion:** CAP exhibited potent inhibitory action against the terbinafine-resistant strains of *T. mentagrophytes*. In the light of increasing antifungal resistance and clinical burden caused by dermatophytosis, CAP may offer a promising alternative or adjuvant therapy. Further studies are required to optimize treatment parameters and evaluate clinical applicability.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

**Keywords:** Cold atmospheric plasma, Fungal growth inhibition, Plasma-based antifungal therapy, Terbinafine-resistant dermatophytes, *Trichophyton mentagrophytes* complex

**Corresponding Author:** Seyed Mohammad Atyabi

Department of Nanobiotechnology, Pasteur Institute of Iran, Tehran, Iran; E-mail: mohammadatyabi@yahoo.com



**Iranian Biomedical Journal Supplementary (February 2026): 38**