

A Novel Defensin-Like Peptide Associated with Two Other New Cationic Antimicrobial Peptides in Transcriptome of the Iranian Scorpion Venom

Masoumeh Baradaran¹, Amir Jalali², Maryam Naderi soorki³ and Hamid Galehdari^{*1}

¹Toxicology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; ²Department of Pharmacology and Toxicology, Toxicology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; ³Department of Genetics, Faculty of Sciences, Shahid Chamran University of Ahvaz, Ahvaz, Iran

Received 21 June 2016; revised 27 June 2016; accepted 2 July 2016

ABSTRACT

Introduction: Scorpion venom is a source of bioactive peptides, and some antimicrobial peptides (AMPs) have been found in the venom gland of scorpions. Therefore, the discovery of new anti-infective agents is an essential need to overcome the problem of antibiotic resistance of clinical isolates. Here, we describe three new cationic AMPs, including meuVAP-6, meuAP-18-1, and meuPep34 from the venom gland of the Iranian scorpion, *Mesobuthus eupeus*. **Methods:** The cDNA sequences encoding all the three peptides were obtained from the cDNA library of scorpion venom gland and were deposited in the GenBank database. **Results:** MeuVAP-6 and meuAP-18-1 are non-disulphide-bridged antimicrobial peptides, while meuPep34 is a cysteine-rich defensin-like peptide. **Discussion:** All three identified AMPs are rich in arginine and tryptophan. The overall results from the length, net charge, and hydrophobicity index suggested that meuPep34 could be the most active AMPs with the potential ability of biofilm inhibition. The data from molecular characterization of identified AMPs can provide a platform with application in drug discovery programs. **DOI: 10.18869/acadpub.ibj.21.3.190**

Keywords: Antimicrobial cationic peptides, Defensin-like peptide, *Mesobuthus eupeus*

Corresponding Author: Hamid Galehdari

Toxicology Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; Tel.: (+98-613) 3338965;

E-mail: galehdari187@yahoo.com

INTRODUCTION

Defensins are a group of antimicrobial peptides (AMPs) and a part of innate immunity in vertebrates and invertebrates^[1]. The first scorpion defensin was isolated from the hemolymph of the North African scorpion, *Leiurus quinquestriatus*, based on the similarity with insect defensin. It was also active against Gram-positive *M. luteus* but inactive against Gram-negative *E. coli*^[2]. Later, some other AMPs with different effects were identified from various scorpion venomes^[3].

AMPs as a part of the immunity system of all animals have the length of four to more than 100 amino acids and also are wide-spectrum molecules^[4]. Finding new effective AMPs is valuable because resistance to available antibiotics is rising constantly,

and the need for new antibiotics to replace is increasing^[5,6]. One promising class of antibiotics is AMPs^[7]. In the current study, we investigated the transcriptome of the Iranian scorpion *Mesobuthus eupeus* venom gland with the aim of finding new AMPs. This study represents the molecular and bioinformatics analysis of three founded AMPs.

MATERIAS AND METHODS

Double-stranded cDNA was synthesized from the total RNA extracted from the venom glands of *M. eupeus*. Following the cDNA cloning in appropriate cloning vector, the recombinant vectors were transformed into the bacterial host to prepare the cDNA library. The cDNA library was then sequenced

and analyzed to find the AMPs. Nucleotide BLAST and protein BLAST were carried out using online BLAST tool in NCBI (blast.ncbi.nlm.nih.gov), while open reading frames (ORFs) were detected using the online ORF finder tool (www.ncbi.nlm.nih.gov/orffinder/). Molecular weight, theoretical isoelectric pH, and the net charge of peptides (at neutral pH) were predicted using peptide property calculator online tool at Innovagen (www.innovagen.com/proteomics-too/ls). SignalP4.1 (http://www.cbs.dtu.dk/services/SignalP/) was also used to identify any signal peptide (http://www.cbs.dtu.dk/services/SignalP/). The hydrophobicity of each peptide was calculated by GPMaw lite online tool available in http://www.alphalyse.com/gpmaw_lite.html. The secondary structure of peptides was predicted by using the GOR secondary structure prediction method software (npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_gor4.html)^[8]. The alignment of every peptide was performed using Alignment tool in UniProt (www.uniprot.org/align). Transmembrane domains were predicted by the TMHMM tool (http://www.cbs.dtu.dk/services/

TMHMM/). The schematic diagrams of protein domain structures were drawn by using the Illustrator for biological sequences (IBS, version 1.0) software (http://ibs.biocuckoo.org/).

RESULTS

BLASTn analysis and protein BLAST identified four transcripts with acceptable similarity to the known scorpion AMPs ($E < 10^{-3}$). One of the transcripts namely CLAP has previously been identified and described by our team and deposited in NCBI with the accession number of KC108907^[9]. Other transcripts were named meuVAP-6, meuAP-18-1, and meuPep34. As shown in Figure 1, the cDNA sequences of all the three peptides have been deposited in GenBank under the accession numbers KU513845 (meuVAP-6), KU513846 (meuAP-18-1), and KU513849 (meuPep34). The full length cDNA of meuVAP-6 is 314 bp that encodes a 70-amino acid peptide containing a 23-amino acid signal peptide, while that




Fig. 1. cDNA and translated open-reading frame (ORF) for meuVAP-6 (A), meuAP-18-1 (B), and meuPep34 (C). Nucleotides of ORF are single underlined, and stop codons are indicated by asterisks. AAA and TTT motifs in 3'UTR are highlighted with yellow color.

of meuAP-18-1 is 339 bp encoding a 76-amino acid peptide consisting of a 25-amino acid signal peptide. The full length cDNA of 508 bp encoding a 103-amino acid peptide, meuPep34, was also found, with a 20-amino acid predicted signal.

The amino acid sequence alignments of meuVAP-6, meuAP-18-1, and meuPep34 with homologues in the UniProt database and the schematic image of peptides are indicated in Figures 2A, 2B, and 2C, respectively. The comparative sequence analysis of meuVAP-6 and meuAP-18-1 with other scorpion peptides revealed that both of them have some similarities with scorpion venom AMPs. Therefore, it can be concluded that meuVAP-6 and meuAP-18-1 are AMPs. The protein BLAST search and the alignment of meuPep34 with its counterparts indicated that meuPep34 is a peptide from scorpion as it is highly similar to AbCp-7 from scorpion *Androctonus bicolor*. The three-dimensional structure modeling of meuPep34 (Fig. 3) showed that meuPep34 has structural similarity with Bubble, a defensin protein, from *Penicillium brevicompactum*^[10] with the coverage of 29% and the sequence DNA similarity of 41%. As a result, meuPep34 is supposed to be a new scorpion defensin-like peptide. In the predicted three-dimensional model, meuPep34 has three β -sheet strands.

MeuVAP-6 and meuAP-18-1 have no cysteine residue; therefore, they are classified in non-disulphide-bridged antimicrobial peptides (NDBPs), while meuPep34 contains 8 cysteine residues in which the pattern of CXCX₇CX₁₂CX₇CX₁₃CX₄CX₁₀C was found.

Predicted secondary structures and some physiochemical properties of meuVAP-6, meuAP-18-1, and meuPep34 are summarized in Table 1. Based on the positive net charges and the existence of basic amino acids (arginine and tryptophan [RW]) among the amino acids of all the three peptides (Fig. 2A, 2B, and 2C), we concluded that the described peptides are all entirely cationic.

DISCUSSION

Scorpion venom gland is considered as a source of novel AMPs^[3]. Considering the importance of AMPs as new anti-infective compounds, this study was designed to identify some novel anti-microbial peptides expressed in scorpion venom gland. In the current study, we presented three new cationic AMPs, including meuVAP-6, meuAP-18-1, and meuPep34.

Protein blasts of meuVAP-6 and meuAP-18-1 against homologues in the UniProt database indicates

that N-terminal and C-terminal of both peptides are most highly conserved (Fig. 2a and 2B). Meucin-13 and Meucin-18-1 from Chinese *M. eupeus* with the identity of 91.4% and 92.1% are the most similar peptides to meuVAP-6 (Fig. 2A) and meuAP-18-1 (Fig. 2B), respectively. Regarding these similarities, it can be suggested that meuVAP-6 and meuAP-18-1 are the new variants of known peptides in the Chinese scorpion, or they are completely novel peptides. Moreover, the finding of identical peptides in the venom of different scorpion species has been indicated to be common^[11]. Furthermore, the finding of similar peptides in scorpions from the same strain might be due to subspecies.

As mentioned above, meuVAP-6 and meuAP-18-1 are NDBPs. Zeng *et al.*^[12] have classified the scorpion NDBPs into six subfamilies. Based on this classification, meuVAP-6 and meuAP-18-1 are clustered in third subfamily containing long chain peptides, including hadrurin, parabutoporin, BmKbpb, pandinin 1, opistoporin 1, and opistoporin 2. In NDBPs grouping of Harrison *et al.*^[3], meuVAP-6 and meuAP-18-1 are also placed in the long-chain peptide group. Zeng *et al.*^[12] have described some features for cDNA of NDBPs and their encoded precursors. These features are extended for meuVAP-6 and meuAP-18-1 and includes AAA or TTT motif at 3'UTR of cDNA (Fig. 1), the existence of a small neutral residue (glycine, alanine, or serine) at the cleavage site of signal peptide, the presence of one positively charged residue, Lys, in N-terminal region, as well as one negatively charged residue (glutamic acid or aspartic acid) at C-terminal region of each precursor (Fig. 2A and 2B). It has been claimed that the presence of a rich AAA or TTT motif at 3'UTR of cDNA have positive effect on the stability of RNA^[12]. Although meuPep34 is not a NDBP, it is very rich in AAA or TTT motif at 3'UTR of its cDNA (Fig. 1), which may affect the RNA stability.

Based on the existence of signal peptide^[13], it is suggested that all of the three described peptides are secretory peptides, i.e. when they are inserted into endoplasmic reticulum, the signal peptides are removed by a signal peptidase enzyme, and the mature peptide would be released in endoplasmic reticulum. Their transmembrane predicted domains (Fig. 2A, 2B, and 2C) consist of 23 amino acids and contain a large part of signal peptide plus four or six (six in meuVAP-6 and meuPep34, four in meuAP-18-1) beginning amino acids of mature peptides. It is proposed that the peptides attach to the membrane of endoplasmic reticulum through these domains.

It is known that the insertion of highly basic peptides into the cytosol perturb the integrity of the




Fig. 2 The left images are schematic image of (A) meuVAP-6, (B) meuAP-18-1, and (C) meuPep34 with its domains. The right images shows aa sequences of (A) meuVAP-6, (B) meuAP-18-1, (C) meuPep34 and aa of each domain. (A) Down image indicates aa sequences alignment of meuVAP-6 with homologues, including venom antimicrobial peptide-6 (Meucin-13), Chinese *M. eupeus* (AC. no.: E4VP07), NDBP11, *Lychas mucronatus* (AC. no.: A0A0U1S7U5), Peptide BmKn1, *Mesobuthus martensii* (AC. no.: Q9GQW4), Peptide BmKn2, *Mesobuthus martensii* (AC. no.: Q6JQN2), Amphiphatic peptide Tx348, *Buthus occitanus israelii* (AC. no.: B8XH50), non-disulfide-bridged peptide androcin 10-2, *Androctonus bicolor* (AC. no.: A0A0K0LBS0), non-disulfide-bridged peptide androcin 10-1, and *Androctonus bicolor* (AC. no.: A0A0K0LCH4). (B) Down image shows the aa sequences alignment of meuAP-18-1 with homologues, including antimicrobial peptide meucin-18-1, Chinese *M. eupeus* (AC. no.: F6K5S5), antimicrobial peptide marcin-18, *Mesobuthus martensii* (AC. no.: F6K5S6), antimicrobial peptide marcin-18-1, *Mesobuthus martensii* (AC. no.: F6K5S7), antimicrobial peptide megin-18, *Mesobuthus gibbosus* (AC. no.: A0A059U8Y9), antimicrobial peptide AMP2, *Odontobuthus dorae* (AC. no.: A0A0U4LVY4), non-disulfide-bridged peptide androcin 18-1, *Androctonus bicolor* (AC. no.: A0A0K0LBT6), non-disulfide-bridged peptide androcin 18-4, *Androctonus bicolor* (AC. no.: A0A0K0LBT1), non-disulfide-bridged peptide androcin 18-2, *Androctonus bicolor* (AC. no.: A0A0K0LBT8). (C) Down image displays aa sequences alignment of meuPep34 with homologue peptide cellular protein AbCp-7 from *Androctonus bicolor* (Ac.no.: A0A0K0LC44). Conserved residues are represented in dark gray. Arginine and tryptophan are indicated with black color. Transmembrane domain is highlighted with yellow highlight. Residues with similar physicochemical properties are shown in light gray. Dash shows that there is not any residue in that position. The percentages of similarity with meuVAP-6, meuAP-18-1, and meuPep34 are brought in the right of alignment. Cysteine residues are indicated with red rectangle.

and 1W could be the potent inhibitor of biofilms. Moreover, as it can be conferred from the amino acid alignment (Fig. 2C), just one uncharacterized scorpion homologue exists for meuPep34. However, based on structural modeling that indicated a similarity with a known defensin, we suggest that meuPep34 is possibly a defensin. The alignment of meuPep34 with its homologue revealed that unlike the meuPep34, its homologue has seven cysteine residues, which is one residue less than meuPep34. The difference in numbers of cysteines may affect the final structure of the peptide. In the present study, we have identified new peptides in the Iranian Scorpion Venom with putative antibacterial activity. Therefore, the result of this study may introduce novel antimicrobial peptides to pharmaceutical industry.

ACKNOWLEDGMENTS

This study was financially supported by Ahvaz Jundishapur University of Medical Sciences (Ahvaz, Iran). We are thankful of Shahid Chamran University authorities for assistance in preparing laboratory equipments. We also thank Dr. Babak Vazirian for his kind supports.

CONFLICT OF INTEREST. None declared.

REFERENCES

- Ganz T. Defensins: antimicrobial peptides of innate immunity. *Nature reviews immunology* 2003; **3**(9):710-720.
- Bontems F, Roumestand C, Gilquin B, Menez A and Toma F. Refined structure of charybdotoxin: common motifs in scorpion toxins and insect defensins. *Science* 1991; **254**(5037): 1521-1523.
- Harrison PL, Abdel-Rahman MA, Miller K, Strong PN. Antimicrobial peptides from scorpion venoms. *Toxicon* 2014; **88**: 115-137.
- Wenzel M, Chiriac AI, Otto A, Zweytick D, May C, Schumacher C, Gust R, Albada HB, Penkova M, Kramer U, Erdmann R, Metzler-Nolte N, Straus SK, Bremer E, Becher D, Brotz-Oesterhelt H, Sahl HG and Bandow JE. Small cationic antimicrobial peptides delocalize peripheral membrane proteins. *Proceedings of the national academy of sciences USA* 2014; **111**(14): E1409-E1418.
- Tavares LS, Silva CSF, de Souza VC, da Silva VL, Diniz CG and Santos MO. Strategies and molecular tools to fight antimicrobial resistance: resistome, transcriptome, and antimicrobial peptides. *Frontiers in microbiology* 2013; **4**: 412.
- Bahar AA and Ren D. Antimicrobial Peptides. *Pharmaceuticals* 2013; **6**(12): 1543-1575.
- Malmsten M. Antimicrobial peptides. *Upsala journal of medical sciences* 2014; **119**(2): 199-204.
- Garnier J, Gibrat JF, Robson B. GOR method for predicting protein secondary structure from amino acid sequence. *Methods in enzymology* 1996; **266**: 540-553.
- Baradaran M, Jalali A, Jolodar A, Ghasemian S. New caerin-like antibacterial peptide from the venom gland of the Iranian scorpion *Mesobuthus eupeus*: cDNA amplification and sequence analysis. *African journal of biotechnology* 2012; **11**: 10176-10181.
- Olsen JG, Flensburg C, Olsen O, Bricogne G, Henriksen A. Solving the structure of the bubble protein using the anomalous sulfur signal from single-crystal in-house Cu K α diffraction data only. *Acta crystallographica section biological crystallography* 2004; **60**(Pt 2): 250-255.
- Kuzmenkov AI, Vassilevski AA, Kudryashova KS, Nekrasova OV, Peigneur S, Tytgat J, Feofanov AV, Kirpichnikov MP, Grishin EV. Variability of potassium channel blockers in mesobuthus eupeus scorpion venom with focus on Kv1.1: An integrated transcriptomic and proteomic study. *Journal of biological chemistry* 2015; **290**(19): 12195-12209.
- Zeng XC, Corzo G and Hahin R. Scorpion venom peptides without disulfide bridges. *The international union of biochemistry and molecular biology life* 2005; **57**(1): 13-21.
- Nothwehr SF, Gordon JI. Targeting of proteins into the eukaryotic secretory pathway: signal peptide structure/function relationships. *Bioessays* 1990; **12**(10): 479-484.
- Mabrouk K, Van Rietschoten J, Vives E, Darbon H, Rochat H, Sabatier JM. Lethal neurotoxicity in mice of the basic domains of HIV and SIV Rev proteins. Study of these regions by circular dichroism. *FEBS letters* 1991; **289**(1): 13-17.
- Fjell CD, Hiss JA, Hancock RE, Schneider G. Designing antimicrobial peptides: form follows function. *Nature reviews drug discovery* 2011; **11**(1): 37-51.
- Taheri-Araghi S, Ha B-Y. Cationic antimicrobial peptides: a physical basis for their selective membrane-disrupting activity. *Soft matter* 2010; **6**: 1933-1940.
- Dathe M, Nikolenko H, Meyer J, Beyermann M, Bienert M. Optimization of the antimicrobial activity of magainin peptides by modification of charge. *FEBS letters* 2001; **501**(2-3):146-150.
- Taheri-Araghi S, Ha BY. Physical basis for membrane-charge selectivity of cationic antimicrobial peptides. *Physical review letters*. 2007; **98**(16): 168101.
- Shai Y. Mechanism of the binding, insertion and destabilization of phospholipid bilayer membranes by alpha-helical antimicrobial and cell non-selective membrane-lytic peptides. *Biochimica et biophysica acta* 1999; **1462**(1-2): 55-70.
- Zaslhoff M. Antimicrobial peptides of multicellular organisms. *Nature* 2002; **415**(6870): 389-395.
- Brogden KA. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nature reviews*

- microbiology* 2005; **3**(3): 238-250.
22. de la Salud Bea R, Ascuitto MR, de Johnson LE. Synthesis of analogs of peptides from *Buthus martensii* scorpion venom with potential antibiotic activity. *Peptides* 2015; **68**: 228-232.
 23. Lee MT, Chen FY and Huang HW. Energetics of pore formation induced by membrane active peptides. *Biochemistry* 2004; **43**(12): 3590-3599.
 24. Liu Z, Brady A, Young A, Rasimick B, Chen K, Zhou C, Kallenbach NR. Length effects in antimicrobial peptides of the (RW)_n Series. *Antimicrobial agents and chemotherapy* 2007; **51**(2): 597-603.
 25. Lewis K. Riddle of biofilm resistance. *Antimicrob agents chemotherapy* 2001; **45**(4): 999-1007
 26. Hou S, Zhou C, Liu Z, Young AW, Shi Z, Ren D, Kallenbach NR. Antimicrobial dendrimer active against *Escherichia coli* biofilms. *Bioorganic and medicinal chemistry letters* 2009; **19**(18): 5478-5481.
 27. Hou S, Liu Z, Young AW, Mark SL, Kallenbach NR, Ren D. Effects of Trp- and Arg-containing antimicrobial-peptide structure on inhibition of *Escherichia coli* planktonic growth and biofilm formation. *Applied and environmental microbiology* 2010; **76**(6): 1967-1974.