DOR: 20.1001.1.1028852.2017.21.4.7.9

Plackett-Burman Design and Response Surface Optimization of Medium Trace Nutrients for Glycolipopeptide Biosurfactant Production

Maurice George Ekpenyong^{*1}, Sylvester Peter Antai¹, Atim David Asitok¹ and Bassey Offiong Ekpo^{2,3}

¹Environmental Microbiology and Biotechnology Unit, Department of Microbiology, Faculty of Biological Sciences, University of Calabar, P.M.B.1115 Calabar, Nigeria; ²Exploration, Research and Services Section, Research and Development (R&D) Division, Nigerian National Petroleum Corporation (NNPC), Port-Harcourt, Nigeria; ³Environmental Geochemistry Unit, Department of Pure and Applied Chemistry, Faculty of Physical Sciences, University of Calabar, P.M.B. 1115, Calabar, Nigeria

Received 29 August 2016; revised 8 October 2016; accepted 10 October 2016

ABSTRACT

Background: A glycolipopeptide biosurfactant produced by *Pseudomonas aeruginosa* strain IKW1 reduced the surface tension of fermentation broth from 71.31 to 24.62 dynes/cm at a critical micelle concentration of 20.80 mg/L. The compound proved suitable for applications in emulsion stabilization in food, as well as in cosmetic and pharmaceutical formulations. **Methods**: In the present study, Plackett-Burman design (PBD) and response surface method (RSM) were employed to screen and optimize concentrations of trace nutrients in the fermentation medium, to increase surfactant yield. **Results**: The PBD selected 5 significant trace nutrients out of the 12 screened. The RSM, on the other hand, resulted in the production of 84.44 g glycolipopeptide/L in the optimized medium containing 1.25 mg/L nickel, 0.125 mg/L zinc, 0.075 mg/L iron, 0.0104 mg/L boron, and 0.025 mg/L copper. **Conclusion**: Significant second-order quadratic models for biomass (P<0.05; adjusted R^2 =94.29%) and biosurfactant (R^2 =99.44%) responses suggest excellent goodness-of-fit of the models. However, their respective non-significant lack-of-fit (Biomass: F=1.28; P=0.418; Biosurfactant: F=1.20; P=0.446) test results indicate their adequacy to explain data variations in the experimental region. The glycolipopeptide is recommended for the formulation of inexpensive pharmaceutical products that require surface-active compounds. **DOI: 10.18869/acadpub.ibj.21.4.249**

Keywords: Pseudomonas aeruginosa, Surface-active agents, Fermentation, Nickel, Copper

Corresponding Author: Maurice George Ekpenyong

Environmental Microbiology and Biotechnology Unit, Department of Microbiology, Faculty of Biological Sciences, University of Calabar, P.M.B.1115 Calabar, Nigeria. Tel.: (+234-70)38218972; E-mail: maurygg2002@yahoo.com or maurygg2002@unical.edu.ng

INTRODUCTION

Biosurfactants are amphiphilic compounds that reduce the free surface enthalpy per unit area of surfaces and interfaces^[1]. They are derived from biological sources including plants, animals, and microorganisms; however, world commercial interest has focused, principally, on microbial derivatives from bacteria, yeasts, and molds^[2].

Microbial surfactants are of diverse chemical nature, including glycolipids^[3], flavolipids^[4], lipopeptides^[5], and glycolipopeptides^[6-8]. Each biosurfactant performs specialized function(s) in their respective producing organisms. Such functions include solubilization, emulsification, pathogenesis, antibiosis, swarming motility, as well as wetting of and attachment to surfaces^[9].

Commercial interest in biosurfactant production has

heightened over the years owing to their increasingly widening applications. Surface-active compounds are applied for different purposes in pharmaceutical, detergent, food, cosmetic, petroleum, and agricultural sectors, as well as the environment^[10,11]. A threatening bottleneck in microbial production of biosurfactants is the economics of production^[1]. Several strategies exist to increase biosurfactant yield, that include the use of low-cost substrates, medium optimization, strain improvement, and development of fermentation conditions^[12,13].

Optimization of fermentation media involving major and trace nutrients, as well as environmental and fermentation conditions has been documented^[14-16]. A good number of biosurfactant fermentation media compositions incorporating both major and minor nutrients have been reported without information on source of trace element composition. A remarkable study on trace mineral composition of a biosurfactant medium has been reported by Joshi *et al.*^[17].

Trace elements are required in small amounts and play essential roles in cellular metabolism, mostly as enzyme co-factors and/or side groups in some microbial metabolites^[18]. Selection of significant trace elements for incorporation into fermentation media is an uphill task from the viewpoint of variable size. A classical method for screening large variables is the use of Plackett-Burman design (PBD). It is a small-sized two-level factorial experimental design programmed to identify critical physicochemical parameters from N number of variables in N+1 experiments without recourse to the interaction effects between and among the variables. Since the sample size is traditionally small, the interaction effects are completely shrouded in the main effects. PBD therefore simply screens the design space to detect large main effects^[19]. The selected parameters are further optimized by the means of an appropriate design technique of a response surface method (RSM). This method is a collection of statistical techniques that uses design of experiments to build models, evaluate the effect of factors and predict optimum conditions for the factors^[20]. The objective of the present study was to optimize the conditions of trace nutrients in a fermentation medium using PBD and RSM to improve biosurfactant yield.

MATERIALS AND METHODS

Producing organism

The bacterium, *Pseudomonas aeruginosa* strain IKW1, was earlier isolated^[6,21]. The stock culture of the bacterium was retrieved and subcultured in Tryptic Soy Agar by the quadrant-streak plate technique at

30°C for 36 h. The bacterium was passed a second time through the reactivation step in the same medium for 24 h, after which a loop-full of culture was used to inoculate freshly prepared Luria broth in 50-mL Erlenmeyer flask containing 10 mL of medium. Flasks were incubated on a rotary shaker at 150 rpm at room temperature (28±2°C) for 18 h.

Optimization experiments Screening trace elements

The PBD incorporated into MINITAB 17 statistical software (trial version) was used to screen 12 trace nutrients in 20 randomized experimental runs. The nutrients were purchased from Sigma Aldrich, USA and included NaCl, KCl, CaCl₂, MgSO₄.7H₂O, CuSO₄.5H₂O, NiCl₂.6H2O, FeCl₃, ZnCl₂.7H₂O, K₃BO₃, MoNa₂O₄.2H₂O, CoCl₂, and MnSO₄.4H₂O. Each was tested only at two levels, low and high. Biosurfactant concentration (g/L) served as the only Fermentation medium response variable. formulated according to the experimental design and based on 50 mL of each formulation dispensed into 250-mL Erlenmeyer flask. Phosphorus (1.0 g/L (NH₄)₂HPO₄/NH₄H₂PO₄) and carbon (5% v/v waste frying sunflower oil with composition of [% w/w] stearic acid [2.21], palmitic acid [6.11], volatile fractions (16.23), oleic acid [22.34], and linoleic acid [50.76]; saponification value 76 and density [30°C] 283.3 kg/m³) sources were added to the media, and flasks were corked and sterilized by autoclaving at 121°C for 15 min. Upon cooling, filter-sterilized urea (1.19 g/L) and inoculum $(10\% \text{ v/v-}10^8 \text{ cells/mL})$ were added to the medium, and flasks were incubated on a rotary shaker agitating at 150 rpm at room temperature for 72 h. All arrangements were made in triplicates.

Biosurfactant quantification was performed as described elsewhere^[6]. Data were analyzed using the same statistical software that generated the design. Significant trace elements were selected for RSM and optimization.

Response surface design and fermentation studies

The experimental design employed to fit the multiple regression models of the fermentation study was a 2⁵⁻¹ half-fractional factorial central composite rotatable design. The test variables were Ni²⁺, Zn²⁺, Fe³⁺, B³⁺, and Cu²⁺; each at five levels. The selection of factor levels for RSM followed, on one hand, the method of the path of steepest ascent, a procedure that moves nutrient levels sequentially in the direction of maximum increase in the response investigated, and on the other, a consideration for the introduction of biomass concentration as a second response variable to facilitate the determination of biosurfactant production

yield, Yp/x. Results of preliminary screening for the effects of increasing concentrations of metals on the growth of the bacterium were also considered in the selection of factor levels of significant trace nutrients from PBD for biomass and biosurfactant formation. Actual levels of each factor were calculated using the equation of Myers and Montgomery^[20].

Coded value = actual level
$$-\frac{high \ level + low \ level}{2}$$

$$\div \frac{high \ level - low \ level}{2} \qquad \textit{Eqn. 1}$$

The coded levels were determined as follows: $X_1 = (\text{Ni}^{2^+} - 0.75)/0.25, \quad X_2 = (\text{Zn}^{2^+} - 0.075)/0.025, \quad X_3 = (\text{Fe}^{3^+} - 0.075)/0.025, \quad X_4 = (\text{B}^{3^+} - 0.03)/0.01, \text{ and } X_5 = (\text{Cu}^{2^+} - 0.03)/0.01, \quad X_5 = (\text{Cu}^{2^+} - 0.03)/0.01)$ 0.075)/0.025. The response variables were biomass Y_1 (g/L) and biosurfactant Y₂ (g/L) concentrations. The experimental design required 32 experimental runs, which were set up in 250-mL Erlenmeyer flasks, each containing 50 mL of fermentation medium composed (g/L) of Na₂HPO₄/KH₂PO₄ (2:1; 4.5), MgSO₄.7H₂O NaCl (0.5), $CaCl_2$ (0.5), KClMoNa₂O₄.2H₂O (0.05 mg/L), CoCl₂.6H₂O (0.05)mg/L), and MnSO₄.4H₂O (0.05 mg/L). Different concentrations of the five significant trace elements obtained from PBD screening were then added to the flasks according to the actual values of the factor levels specified by the coded values in Table 1. Medium pH was adjusted to 7.0 with KOH pellets^[4]. Waste frying

oil was subsequently added to each flask at 5% (v/v) concentration and flasks, without nitrogen source, were sterilized by autoclaving at 121° C for 15 min. Upon cooling, filter-sterilized urea (1.19 g/L) and inoculum (10% v/v– 10^{8} cells/mL) were introduced into flasks prepared in triplicates, and then the flasks were incubated on rotary shakers agitating at 150 rpm at room temperature for 72 h.

Determination of biomass (Y1) and biosurfactant (Y2) concentrations

Small portions of 72-h fermentation broth from each experimental setting (10 mL) were centrifuged at 8,000 $\times g$ for 10 min. Cell-free supernatants were collected, and the cell pellets were washed twice in demineralized water. The supernatants were subjected to 0.45- μM and 0.2- μM Millipore membrane filtrations and subsequently to acid precipitation with 6N HCl, pH 2.0. Biomass and biosurfactant concentrations were determined from cell pellets and acid precipitates of supernatants, respectively as described previously $^{[6]}$.

Statistical analysis

All data generated from the factorial experiment were subjected to multiple regression analysis using least squares to build the regression models. A second-order (quadratic) function was used to fit the data generated. Experimental design, data analysis, interaction plotting, and optimization of factor conditions were done with MINITAB 17 statistical software, while Excel 2007 was used for confirmation

Table 1. Placket-Burman design matrix (randomized) for trace element contribution to biosurfactant formation in coded units

Run	A	В	C	D	E	F	G	Н	J	K	L	M	BSC (g/L)
1	-1	-1	-1	1	-1	1	-1	1	1	1	1	-1	23.58
2	1	1	1	1	-1	-1	1	1	-1	1	1	-1	25.96
3	-1	-1	1	1	-1	1	1	-1	-1	-1	-1	1	24.21
4	1	-1	-1	1	1	-1	1	1	-1	-1	-1	-1	21.43
5	-1	1	1	-1	-1	-1	-1	1	-1	1	-1	1	24.19
6	1	1	-1	1	1	-1	-1	-1	-1	1	-1	1	28.96
7	-1	1	1	1	1	-1	-1	1	1	-1	1	1	31.73
8	1	1	-1	-1	-1	-1	1	-1	1	-1	1	1	32.75
9	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	17.45
10	1	-1	1	-1	1	1	1	1	-1	-1	1	1	36.02
11	1	-1	1	1	1	1	-1	-1	1	1	-1	1	24.72
12	-1	1	-1	1	-1	1	1	1	1	-1	-1	1	23.48
13	1	1	1	-1	-1	1	1	-1	1	1	-1	-1	22.31
14	1	-1	1	1	-1	-1	-1	-1	1	-1	1	-1	27.08
15	-1	-1	-1	-1	1	-1	1	-1	1	1	1	1	33.27
16	-1	1	-1	1	1	1	1	-1	-1	1	1	-1	25.31
17	-1	1	1	-1	1	1	-1	-1	-1	-1	1	-1	28.78
18	1	1	-1	-1	1	1	-1	1	1	-1	-1	-1	21.08
19	1	-1	-1	-1	-1	1	-1	1	-1	1	1	1	33.10
20	-1	-1	1	-1	1	-1	1	1	1	1	-1	-1	23.29

A, boron; B, calcium, C, cobalt; D, copper; E, iron; F, potassium; G, magnesium; H, manganese; J, molybdenum; K, sodium; L, nickel; M, zinc; '1', high value; '-1', low value; BSC, biosurfactant concentration

of model fits where predicted responses were plotted against experimentally-derived data. All hypotheses were tested at 95% confidence level.

For the five factors considered in the optimization experiment, the quadratic model took the form below:

 $Y = bo + b1X1 + b2X2 + b3X3 + b4X4 + b5X5 + b11X1^2 + b22X2^2 + b33X3^2 + b44X4^2 + b55X5^2 + b12X1X2 + b13X1X3 + b14X1X4 + b15X1X5 + b23X2X3 + b24X2X4 + b25X2X5 + b34X3X4 + b35X3X5 + b45X4X5$ Eqn. 2

where *b*0 is the value of the fixed response at the central point (0, 0, 0, 0); *b*1, *b*2, *b*3, *b*4, and *b*5 the coefficients of the linear terms; *b*11, *b*22, *b*33, *b*44, and *b*55 the coefficients of the quadratic terms; *b*12, *b*13, *b*14, b15, *b*23, *b*24, *b*25, *b*34, *b*35, and *b*45 are the coefficients of the cross products (interactive terms).

Verification experiments

Studies to confirm the validity of the results of the optimization experiment were done by setting up glycolipopeptide production in the optimized medium. The medium had the same composition as that described in Materials and Methods with concentrations of nickel, zinc, iron, boron, and copper incorporated in accordance with the results of the various optimized conditions. Fermentation was conducted in 250-mL Erlenmeyer flasks containing 50 mL of fermentation medium supplemented with 5%

(v/v) waste frying sunflower oil as carbon source. Cooled sterilized media were supplemented with filtersterilized urea (1.19 g/L) and then inoculated with 10% (v/v-10⁸ cells/mL) inoculum of *Pseudomonas aeruginosa* strain IKW1. Flasks were incubated as earlier described and biomass and biosurfactant concentrations determined as described in Ekpenyong *et al.*^[6]. Means of triplicate determinations of concentrations of biomass and biosurfactant from corroborating experiments were compared with those predicted by the regression models.

RESULTS AND DISCUSSION

Plackett-Burman design

The design matrix of the PBD for the effects of 12 trace nutrients on biosurfactant production and their responses are shown in Table 1. Results showed the highest biosurfactant concentration of 36.02 g/L in run 10. However, the results of the trace element modeling experiment by PBD revealed that only 5 out of 12 nutrient elements significantly influenced the glycolipopeptide production. The non-selection of the remaining seven elements suggests their non-significant (P>0.05) contributions to the response under investigation at the confidence level selected for the study.




Fig. 1. Normal plot of standardized effects of significant trace nutrients of a Plackett-Burman design for glycolipoeptide-biosurfactant production. Bo is used loosely to indicate boron and not as a chemical symbol.




Fig. 2. Main effects plots of contributions of significant trace elements to glycolipopeptide-biosurfactant production by *Pseudomonas aeruginosa* strain IKW1. BSC, biosurfactant concentration

Results of the trace element screen test by PBD indicated the same results as trace element modeling experiment. Figure 1 is the normal plot of the standardized effect of the significant nutrients, showing the magnitude and direction of their significant effects. The Figure reveals that nickel has the highest significant positive effect on biosurfactant production by the bacterium since its effect is positioned the furthest to the right of the response line. Other nutrients with significant enhancement effect on biosurfactant production were zinc, iron, and boron, which is in agreement with the Joshi *et al.*'s study^[17]. However, the Figure reveals a significant reductive effect of copper on glycolipopeptide production by the bacterium since its effect is positioned to the left of the biosurfactant response line. Figure 2 shows the main effects plots of significant trace nutrients on the response variable and confirms the results displayed in Figure 1. Nickel has been shown to make the highest contribution to biosurfactant production, whereas copper exerts a negative effect on biosurfactant production.

The analysis of variance (ANOVA) result for significant trace elements is demonstrated in Table 2 and confirms nickel as having the most significant (P<0.05) enhancement effect on biosurfactant production given by its very large F value. The linear regression coefficient of determination, adjusted R^2 of 91.81%, indicates that the model equation (below), given in un-coded units, was significant and could explain 91.81% of the variability in the response data. The equation reveals that nickel has the largest coefficient that is preceded by a positive sign, confirming once again its strong enhancement effect on biosurfactant formation.

$$BSC$$
 = 26.435 + 0.906 $Bo - 0.789 Cu + 1.024 Fe + 3.323 Ni$ + 2.808 Zn $Eqn. 3$

Response surface optimization

Experimental data of the response surface method

Table 3 shows the actual factor levels corresponding to coded factor levels for a 2⁵⁻¹ half-fractional factorial,

Table 2. Analysis of variance (ANOVA) of the regression model from the Plackett-Burman design for trace element contribution to biosurfactant formation in un-coded units

Source	DF	Adj. SS	Adj. MS	F value	P value
Model	5	428.38	85.677	43.61	0.000
Linear	5	428.38	85.677	43.61	0.000
Bo	1	16.42	16.417	8.36	0.012
Cu	1	12.45	12.450	6.34	0.025
Fe	1	20.97	20.972	10.67	0.006
Ni	1	220.85	220.847	112.41	0.000
Zn	1	157.70	157.697	80.27	0.000
Error	14	27.51	1.965		
Total	19	455.89			

Model Summary: S, 1.40166; R^2 , 93.97%; adjusted R^2 , 91.81%; predicted R^2 , 87.69%; P<0.05, 5% significance level. Bo is used loosely to indicate boron and not as a chemical symbol. DF, degrees of freedom; SS, sum of squares; MS, mean sum of squares.

Table 3. Actual factor levels corresponding to coded factor levels for 2⁵⁻¹ half-fractional factorial central composite rotatable design of response surface method

Variable (mg/L)	Actual values							
variable (mg/L)	Code	-2	-1	0	1	2		
NiCl ₂ .6H ₂ O	X_1	0.250	0.50	0.750	1.00	1.250		
$ZnSO_4.7H_2O$	X_2	0.025	0.05	0.075	0.10	0.125		
FeCl ₃	X_3	0.025	0.05	0.075	0.10	0.125		
K_3BO_3	X_4	0.010	0.02	0.030	0.04	0.050		
CuSO ₄ .5H ₂ O	X5	0.025	0.05	0.075	0.10	0.125		

Table 4. Actual factor levels corresponding to coded factor levels for the CCRD of the response surface optimization showing biomass and biosurfactant concentrations

Run order	X1	X2	X3	X4	X5	NiCl ₂ (mg/L)	ZnSO ₄ (mg/L)	FeCl ₃ (mg/L)	K ₃ BO ₃ (mg/L)	CuSO ₄ (mg/L)	BMC (g/L)	BSC (g/L)
1	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	16.86	41.03
2	-2	0	0	0	0	0.250	0.075	0.075	0.030	0.075	14.36	29.55
3	-1	-1	-1	1	-1	0.500	0.050	0.050	0.040	0.050	8.96	36.04
4	1	-1	1	1	-1	1.000	0.050	0.100	0.040	0.050	12.98	38.32
5	0	0	0	2	0	0.750	0.075	0.075	0.050	0.075	13.52	28.39
6	0	0	0	-2	0	0.750	0.075	0.075	0.010	0.075	18.52	36.57
7	0	0	0	0	-2	0.750	0.075	0.075	0.030	0.025	15.39	46.47
8	-1	1	-1	-1	-1	0.500	0.100	0.050	0.020	0.050	11.35	38.32
9	-1	1	1	1	-1	0.500	0.100	0.100	0.040	0.050	18.26	35.31
10	1	-1	-1	-1	-1	1.000	0.050	0.050	0.020	0.050	10.62	32.19
11	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	18.76	41.22
12	0	-2	0	0	0	0.750	0.025	0.075	0.030	0.075	9.67	36.82
13	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	18.87	41.46
14	-1	1	-1	1	1	0.500	0.100	0.050	0.040	0.100	13.28	27.54
15	-1	-1	1	1	1	0.500	0.050	0.100	0.040	0.100	13.41	26.79
16	0	0	2	0	0	0.750	0.075	0.125	0.030	0.075	16.59	40.45
17	1	1	-1	1	-1	1.000	0.100	0.050	0.040	0.050	12.38	31.07
18	1	1	-1	-1	1	1.000	0.100	0.050	0.020	0.100	16.58	30.01
19	0	0	-2	0	0	0.750	0.075	0.025	0.030	0.075	8.95	28.12
20	0	2	0	0	0	0.750	0.125	0.075	0.030	0.075	15.38	46.05
21	1	1	1	1	1	1.000	0.100	0.100	0.040	0.100	8.47	39.67
22	0	0	0	0	2	0.750	0.075	0.075	0.030	0.125	11.04	37.58
23	1	-1	1	-1	1	1.000	0.050	0.100	0.020	0.100	11.94	41.40
24	-1	-1	1	-1	-1	0.500	0.050	0.100	0.020	0.050	21.07	29.56
25	1	1	1	-1	-1	1.000	0.100	0.100	0.020	0.050	19.53	57.21
26	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	18.94	42.08
27	-1	1	1	-1	1	0.500	0.100	0.100	0.020	0.100	17.39	37.99
28	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	19.04	41.68
29	-1	-1	-1	-1	1	0.500	0.050	0.050	0.020	0.100	9.47	31.07
30	0	0	0	0	0	0.750	0.075	0.075	0.030	0.075	18.83	40.78
31	1	-1	-1	1	1	1.000	0.050	0.050	0.040	0.100	9.57	30.24
32	2	0	0	0	0	1.250	0.075	0.075	0.030	0.075	12.35	39.00

X1, nickel; X2, zinc; X3, iron; X4, boron; X5, copper; BMC, biomass concentration; BSC, biosurfactant concentration

a central composite rotatable design of RSM. The experimental responses from the 32 experimental runs of the surface methodology are presented in Table 4. Maximum biomass concentration of 21.07 g/L was obtained at conditions set at $(X_1, X_2, X_3, X_4, X_5)=(-1,$ -1, 1, -1, -1) corresponding to 0.5 mg/L nickel, 0.05 mg/L zinc, 0.1 mg/L iron, 0.02 mg/L boron, and 0.05 mg/L copper. In these conditions, biosurfactant concentration obtained was 29.56 g/L. However, the highest biosurfactant concentration of 57.21 g/L was 1, -1, -1) corresponding to 1 mg/L nickel, 0.1 mg/L zinc, 0.1 mg/L iron, 0.02 mg/L boron, and 0.05 mg/L copper and a corresponding biomass concentration of 19.53 g/L. Glycolipopeptide production yield, Yp/x, under these conditions would be 2.93. The attainment of peak biomass and biosurfactant concentrations at different experimental runs suggest the requirement of different trace nutrient conditions for cellular growth and metabolite synthesis, especially metabolites obtained during or near idiophasic metabolism of organisms.

Regression model for biomass concentration, Y₁

The data presented in Table 4 was subjected to multiple regression analyses using least squares regression to fit a second-order (quadratic) regression model for biomass concentration, Y_1 . The model stipulated 20 predictors; however, the result of ANOVA of the model in Table 5 reveals that 5 interactive terms were removed by stepwise selection, since their contributions were not significant at P=0.05. The T values of the predictor coefficients (data not shown) suggest that all model predictors, except the linear terms of zinc and iron, made significant negative contributions to the model, implying their reductive effects on biomass formation. The model equation is presented in un-coded units as equation 4 below:

$$Y1$$
= $18.346 - 0.631 Ni + 1.277 Zn + 1.922 Fe$
- $1.277 Bo - 0.989 Cu - 1.095 Ni * Ni$
- $1.303 Zn * Zn - 1.241 Fe * Fe$
- $0.429 Bo * Bo - 1.130 Cu * Cu$
- $1.456 Ni * Fe - 0.619 Ni * Bo - 0.670 Zn * Fe$
- $0.811 Fe * Bo - 1.639 Fe * Cu$
Eqn. 4

Table 5. Analysis of variance of the 2^{5-1} half-fractional factorial central composite rotatable design of an response surface method for biomass regression model in un-coded units

Source	DF	Adjusted SS	Adjusted MS	F value	P value
Model	15	435.332	29.0222	35.12	0.000
Linear	5	199.895	39.9790	48.38	0.000
Ni	1	9.551	9.5508	11.56	0.004
Zn	1	39.117	39.1171	47.34	0.000
Fe	1	88.627	88.6273	107.26	0.000
Во	1	39.117	39.1171	47.34	0.000
Cu	1	23.483	23.4828	28.42	0.000
Square	5	134.701	26.9401	32.60	0.000
Ni*Ni	1	35.179	35.1787	42.57	0.000
Zn*Zn	1	49.773	49.7729	60.23	0.000
Fe*Fe	1	45.202	45.2022	54.70	0.000
Bo*Bo	1	5.395	5.3951	6.53	0.021
Cu*Cu	1	37.463	37.4633	45.34	0.000
Two-way interaction	5	100.737	20.1473	24.38	0.000
Ni*Fe	1	33.931	33.9306	41.06	0.000
Ni*Bo	1	6.126	6.1256	7.41	0.015
Zn*Fe	1	7.182	7.1824	8.69	0.009
Fe*Bo	1	10.530	10.5300	12.74	0.003
Fe*Cu	1	42.968	42.9680	52.00	0.000
Error	16	13.221	0.8263		
Lack-of-fit	11	9.748	0.8862	1.28	0.418
Pure error	5	3.473	0.6946		
Total	31	448.554			

Model Summary: S, 0.909022; R², 97.05%; adjusted R², 94.29%; Predicted R², 86.75%; *P*<0.05, 5% significance level. Bo is used loosely to indicate boron and not as a chemical symbol. DF, degrees of freedom; SS, sum of squares; MS, mean sum of squares

The metal, iron, is described as the only macro-bioelement of the heavy metals and the most biologically relevant trace nutrient^[18]. The effect of its linear term on biomass formation was understandably pronounced, and the significant reductive effect of its quadratic term suggests possible toxicity at high concentrations and therefore its trace requirement. Abalos *et al.*^[14] have reported a similar result trend during production of rhamnolipid from a strain of *Pseudomonas aeruginosa*.

The requirement of zinc for biomass formation derives from its role in DNA-binding proteins and a variety of enzymes. Zinc is also known to act as a Lewis base needed to activate water for involvement in aqueous reactions^[6].

The significance of a model is given by its goodness-of-fit test, often expressed as the coefficient of determination R^2 , which is the percentage of the variations in the response that can be explained by independent factors and their interactions. In this study, the biomass regression model (Table 5) was highly significant (F=35.12; P=0.000) with an adjusted R^2 of 94.29%, indicating that only 5.71% of the variability in the response is not explainable by the model.

A well-fitted model estimated by adjusted R^2 might not adequately explain data variations in the region of experimentation. The lack-of-fit test is therefore frequently used as a support test for adequacy of the fitted model. The ANOVA table for the biomass regression model in this study shows a non-significant lack-of-fit (F=1.28; P=0.418) for the model, which suggests its adequacy for the explanation of data variations in the region of experimentation.

The regression plot of experimental against predicted biomass responses is presented as Figure 3 and reveals a coefficient of determination, R^2 of 0.9705, which is in agreement with the R^2 for the model in the ANOVA table.

Regression model for biosurfactant concentration, Y₂

The ANOVA for the biosurfactant regression model presented in Table 6 indicated that the model was highly significant (F=371.07; P=0.000). All the linear predictor terms and only the square terms of nickel, iron, and boron were significant. Six of the interactive predictor terms were also significant; however, Fe*Cu interaction, although included in the model, was not significant (P=0.051). The magnitude and direction of significant effect of model predictors are given by the T values of the coefficient estimates of the predictors (data not shown) and reveal that the linear, but not the quadratic terms of nickel, zinc, and iron, significantly enhanced biosurfactant formation. It is probably because of the enhancing effect of Fe*Cu that the term

was included in the model, to increase biosurfactant concentration as long as the adjusted R^2 of the model was not negatively affected. The model equation, in un-coded units, is therefore presented as equation 5 below.

```
Y2
= 41.445 + 2.350 Ni + 2.082 Zn + 3.101 Fe
-2.047 Bo - 2.129 Cu
-1.8826 Ni * Ni - 1.8801 Fe * Fe
-2.3313 Bo * Bo + 3.526 Ni * Fe - 0.641 Ni * Bo
+2.294 Zn * Fe - 1.694 Zn * Bo - 1.256 Zn * Cu
-1.211 Fe * Bo + 0.263 Fe * Cu

Eqn. 5
```

The model was shown to have an adjusted R^2 value of 99.44%, suggesting that only 0.56% of the variability in biosurfactant responses could not be explained by it, and that the observed variations were due to the factor effects and not due to noise. To confirm the adequacy of the biosurfactant model in explaining the variations about the data, an examination of the lack-of-fit test result in the ANOVA table was made and demonstrated a non-significant (F=1.20; P=0.446) lack-of-fit, indicating the adequacy of the model to explain data in the experimental region.

A plot of experimental biosurfactant concentrations against predicted concentrations obtained by solving equation 5 is presented in Figure 4 as the final ratification of the fit of the biosurfactant regression model and confirms a goodness-of-fit, R^2 (unadjusted) value of 99.97%.




Fig. 3. Experimental biomass concentration plotted against biomass concentration predicted by the fitted model.

Table 6. Analysis of variance of the 2⁵⁻¹ half fractional factorial central composite rotatable design (CCRD) of an response surface method for biosurfactant regression model in un-coded units

Source	DF	Adjusted SS	Adjusted MS	F value	P value
Model	15	1388.26	92.551	371.07	0.000
Linear	5	676.69	135.338	542.62	0.000
Ni	1	132.49	132.493	531.21	0.000
Zn	1	104.04	104.042	417.14	0.000
Fe	1	230.83	230.826	925.46	0.000
Во	1	100.57	100.573	403.23	0.000
Cu	1	108.76	108.758	436.05	0.000
Square	3	326.18	108.726	435.92	0.000
Ni*Ni	1	105.31	105.312	422.24	0.000
Fe*Fe	1	105.03	103.033	421.11	0.000
Bo*Bo	1	161.50	161.502	647.52	0.000
2-Way Interaction	7	385.39	55.056	220.74	0.000
Ni*Fe	1	198.88	198.881	797.38	0.000
Ni*Bo	1	6.57	6.566	26.33	0.000
Zn*Fe	1	84.23	84.227	337.69	0.000
Zn*Bo	1	45.93	45.935	184.17	0.000
Zn*Cu	1	25.23	25.226	101.14	0.000
Fe*Bo	1	23.45	23.450	94.02	0.000
Fe*Cu	1	1.11	1.108	4.44	0.051
Error	16	3.99	0.249		
Lack-of-Fit	11	2.90	0.263	1.20	0.446
Pure Error	5	1.09	0.219		
Total	31	1392.25			

Model Summary: S, 0.499416; R^2 , 99.71%; adjusted R^2 , 99.44%; predicted R^2 , 98.77%; P<0.05, 5% significance level. Bo is used loosely to indicate boron and not as a chemical symbol. DF, degrees of freedom; SS, sum of squares; MS, mean sum of squares.

Contour and surface plots for biomass (Y1) and biosurfactant (Y2) concentrations

Biomass response plots were made with the vertical axis representing biomass concentration (Y1) and two horizontal axes representing the most significant two- way interaction (X3, X5)=(Fe, Cu). The plots led to maximal biomass formation with the remaining three factors (X1, X2, X4)=(Ni, Zn, Bo) held at their optimum levels. Biomass concentration under this condition was in excess of 20 g/L. The contour and surface plots displayed in Figure 5 reveal that maximal biomass would be accumulated under the high levels of iron and low levels of copper.

The biosurfactant response plots (Fig. 6) were made with the vertical axis representing biosurfactant concentration (Y2) and two horizontal axes representing the two most significant variables (X4, X5)=(Bo, Cu) that led to maximal biosurfactant formation with the remaining three factors (X1, X2, X3)=(Ni, Zn, Fe) held at their optimum levels. Maximal biosurfactant concentration under this condition was in excess of 80 g/L. The contour and surface plots reveal that the highest biosurfactant concentration would be obtained when fermentation medium contains low levels of both copper and

boron, with the concentrations of nickel, zinc, and iron supplied at high levels.

Boron, a micro-nutrient with intermediary properties of metals and non-metals, is known as a regulator of certain pathways that require serine proteases or oxidoreductases involving coenzymes like nicotinamide adenine dinucleotide and





Fig. 4. Experimental biosurfactant concentrations versus theoretical values predicted by the regression model.






Fig. 5. Contour (A) and surface (B) plots of two-way interactions of independent variables for maximal biomass production. Bo is used loosely to indicate boron and not as a chemical symbol. BMC, biomass concentration

nicotinamide adenine dinucleotide phosphate for activity^[22]. These activities of boron in bacteria could very well be the reason for its very little requirement both for cellular growth and secondary metabolism. Copper, on the other hand, is mostly important in cytochrome C oxidases; oxygen-dependent terminal oxidases in the electron transport chain of aerobic organisms^[23]. The interaction therefore between boron and copper has to be very important in biosurfactant synthesis since the whole process is energy driven; copper to generate energy and boron to inhibit the process.


Nickel is a transition metal useful only in a few selected reactions. It associates with iron in NiFe hydrogenases; an enzyme system that splits molecular hydrogen into protons and electrons. The metal is also intrinsically bound to urease; an enzyme that catalyses the splitting of urea into carbon dioxide and ammonia thus supplying nitrogen in its available form for cellular metabolism^[18,24]. The preference of urea as a nitrogen source for glycolipopeptide

biosurfactant production by *Pseudomonas aeruginosa* strain IKW1 (unpublished data) demystifies the significant enhancement effect of nickel on biosurfactant production and suggests a correlation between urease activity and biosurfactant production in the bacterium.

Selection of optimum conditions for response variables

For situations where biomass accumulation is desired instead of the surface-active compound, which was not the case in this study, the highest predictable biomass concentration of 26.55 g/L was obtained at trace mineral conditions set at (X1, X2, X3, X4, X5)=(-1.0707, -0.0202, 2.0, -2.0, -1.8788) corresponding to 0.482 mg/L nickel, 0.074 mg/L zinc, 0.125 mg/L iron, 0.010 mg/L boron, and 0.028 mg/L copper.

For tertiary oil recovery and spilled oil remediation where maximum concentrations of both responses






Fig. 6. Contour (A) and surface (B) plots of two-way interactions of independent variables for maximal glycolipopeptide production. Bo is used loosely to indicate boron and not as a chemical symbol. BSC, biosurfactant concentration

Table 7. Design codes, actual values, experimental and predicted responses of validation experiments of an RSM for glycolipopeptide production

Parameters	BMC maximized	BSC maximized	BMC and BSC maximized		
X1	-1.0707	2	0.2626		
X2	-0.0202	2	0.9495		
X3	2	2	2		
X4	-2	-1.9596	-2		
X5	-1.8788	-2	-2		
Ni (mg/L)	0.482	1.25	0.816		
Zn (mg/L)	0.074	0.125	0.099		
Fe (mg/L)	0.125	0.125	0.125		
Bo (mg/L)	0.010	0.010	0.010		
Cu (mg/L)	0.028	0.025	0.025		
eBMC (g/L	25.74	19.14	25.14		
pBMC (g/L)	26.55	NP	23.40		
eBSC (g/L)	25.96	84.44	56.83		
pBSC (g/L)	NP	81.92	57.55		

 X_1 , nickel (Ni); X_2 , zinc (Zn); X_3 , iron (Fe); X_4 , boron (Bo); X_5 , copper (Cu); eBMC, experimental biomass concentration; pBMC, predicted biomass concentration; eBSC, experimental biosurfactant concentration; pBSC, predicted biosurfactant concentration; NP, not predicted. Bo is used loosely to indicate boron and not as a chemical symbol.

are desired, trace mineral conditions set at (X1, X2, X3, X4, X5)=(0.2626, 0.9495, 2.0, -2.0, -2.0) corresponding to 0.816 mg/L nickel, 0.099 mg/L zinc, 0.125 mg/L iron, 0.01 mg/L boron, and 0.025 mg/L copper, which would lead to 23.4 g/L biomass and 57.55 g/L biosurfactant.

Finally, for detergent, food, cosmetic, and pharmaceutical applications where maximum surface-active compound and zero microbial cells are desired, the response optimizer set conditions for maximum biosurfactant concentration of 81.92 g/L at (X1, X2, X3, X4, X5)=(2.0, 2.0, 2.0, -1.9596, -2) corresponding to 1.25 mg/L nickel, 0.125 mg/L zinc, 0.125 mg/L iron, 0.0104 mg/L boron, and 0.025 mg/L copper.

Verification experiments

Table 7 shows that maximum biosurfactant concentration obtained in the validation experiments under conditions (X1, X2, X3, X4, X5) =(2, 2, 2, -1.9596, -2) corresponding to 1.25 mg/L nickel, 0.125 mg/L zinc, 0.125 mg/L iron, 0.0104 mg/L boron, and 0.025 mg/L copper was 84.44 g/L with a corresponding biomass concentration of 19.14 g/L, giving a production yield $(Y_{p/x})$ of 4.41. When fermentation conditions were set to maximize both responses, biosurfactant yield of 2.26 was achieved. Compared to the 81.92 g/L biosurfactant concentration predicted by the optimizer, the concentration obtained in the confirmation experiment was only 0.03% higher, thus validating the prediction of the response optimizer. glycolipopeptide biosurfactant concentration of 84.44 g/L, when compared to 23.86 g/L obtained from a previous (control) experiment (unpublished data), reveals a ~3.54fold increase in biosurfactant concentration, thereby making separate optimization experiments for trace nutrients appropriate for developing the fermentation media for the production of microbial metabolites.

PBD selected nickel, zinc, iron, boron, and copper as the most significant (P<0.05; adjusted R^2 =91.82%) trace minerals for glycolipopeptide biosurfactant production. Optimization of the nutrients by RSM resulted in 84.44 g/L of the biosurfactant under conditions set at (X1, X2, X3, X4, X5)=(2, 2, 2, -1.9596, -2). These conditions corresponded to 1.2500 mg/L nickel, 0.125 mg/L zinc, 0.125 mg/L iron, 0.0104 mg/L boron, and 0.025 mg/L copper, giving a production yield, Yp/x of 4.41. In conclusion, PBD and RSM are dependable tools for selecting and optimizing conditions of nutrients for biosurfactant production.

ACKNOWLEDGEMENTS

The authors are grateful to the government of Cross River State, Nigeria for the research grant (CRS/SB/P/149/30) awarded for this work through their Ministry of Education, Scholarship and Grant unit. Sincere gratitude also goes to Mr. Obi Oyeka-Director of Bez Pharmacy and Stores, Calabar, Nigeria and Dr. Samuel Ekpata for their financial assistance towards this research.

CONFLICT OF INTEREST. None declared.

REFERENCES

- 1. Desai JD, Banat IM. Microbial production of surfactants and their commercial potential. *Microbiology and molecular biology reviews* 1997; **61**(1): 47-64.
- 2. Abouseoud M, Maachi R, Amrane A, Boudergua S, Nabi A. Evaluation of different carbon and nitrogen sources in production of biosurfactant by *Pseudomonas fluorescens*. *Desalination* 2008; **223**(1-3): 143-151.
- 3. Banat IM, Franzetti A, Gandolfi I, Bestetti G, Martinotti M G, Fracchia L, Smyth T J, Marchant R. Microbial biosurfactants production, applications and future potential. *Applied microbiology and biotechnology* 2010; **87**(2): 427-444.
- Bodour AA, Drecs KP, Maier RM. Distribution of biosurfactant-producing bacteria in undisturbed and contaminated arid southwestern soils. *Applied and* environmental microbiology 2003; 69(6): 3280-3287.
- Morikawa M, Daido H, Takao T, Murata S, Shimonishi Y, Imanaka T. A new lipopeptide biosurfactant produced by *Arthrobacter* sp. strain MIS38. *Journal of bacteriology* 1993; 175(20): 6459-6466.
- Ekpenyong M, Antai SP, Asitok, A. A *Pseudomonas aeruginosa* strain IKW1 produces an unusual polymeric surface-active compound in waste frying oil-minimal medium. *International journal of sciences* 2016; 5(6): 108-123.
- 7. Ilori MO, Amund DI. Production of a peptidoglycolipid bioemulsifier from *Pseudomonas aeruginosa* grown on hydrocarbon. *Zeitschrift für naturforschung c* 2001; **56**(7-8): 547-552.
- Thavasi R, Jayalakshmi S, Balasubramaian T, Banat IM. Biosurfactants production by *Corynebacterium kutscheri* from waste motor lubricant oil and peanut oil cake. *Letters in applied microbiology* 2007; 45(6): 686-691.
- 9. Ron EZ, Rosenberg E. Natural roles of biosurfactants. *Environmental microbiology* 2001; **3**(4): 229-236.
- Pacwa-Plociniczak M, Plaza GA, Piotrowska-Seget Z, Cameotra SS. Environmental applications of biosurfactants: recent advances. *International journal of* molecular sciences 2011; 12(1): 633-654.
- 11. Sachdev DP, Cameotra SS. Biosurfactants in agriculture. *Applied microbiology and biotechnology* 2013; **97**(3): 1005-1016.
- 12. Sekhon KK, Khanna S, Cameotra SS. Enhanced biosurfactant production through cloning of three genes and role of esterase in biosurfactant release. *Microbial cell factories* 2011; **10**: 49.

- Thavasi R, Subramanyam Nambaru VRM, Jayalakshmi S, Balasubramanian T, Banat IM. Biosurfactant production by *Pseudomonas aeruginosa* from renewable resources. *Indian journal of microbiology* 2011; 51(1): 30-36.
- 14. Abalos A, Maximo F, Manresa MA, Bastida J. Utilization of response surface methodology to optimize the culture media for the production of rhamnolipids by *Pseudomonas aeruginosa* AT10. *Journal of chemical technology and biotechnology* 2002; **77**(7): 777-784.
- 15. Deepika KV, Kalam S, Sridhar PR, Podile AR, Bramhachari PV. Optimization of rhamnolipid biosurfactant production by mangrove sediment bacterium *Pseudomonas aeruginosa* KVR-HR42 using response surface methodology. *Biocatalysis and agricultural biotechnology* 2016; **5**: 38-47.
- Rodrigues L, Teixeira J, Oliveira R, van der Mei H C. Response surface optimization of the medium components for the production of biosurfactants by probiotic bacteria. *Process biochemistry* 2006; 41(1): 1-10.
- 17. Joshi S, Yadav S, Nerurkar A, Desai AJ. Statistical optimization of medium components for the production of biosurfactant by *Bacillus licheniformis* K51. *Journal of microbiology and biotechnology* 2007; **17(2)**: 313-319.
- 18. Nies DH. Microbial heavy-metal resistance. *Applied microbiology and biotechnology* 1999; **51**(6): 730-750.
- 19. Cavazzuti M. *Optimization Methods: from Theory to Design*, London: Springer, 2013.
- Myers R H, Montgomery D C. Response Surface Methodology: Product and Process Optimization Using Designed Experiments. Second ed. New York: John Wiley & Sons, 2002.
- 21. Raorane CJ, Gavimath CC, Kukreja GP, Kalsekar DP, Kulkarni SM, Ravishankar, BE, Hooli RS. Isolation, identification and phylogenetic analysis of phthalic acid-degrading bacteria. *International journal of advanced biotechnology and research* 2012; **3**(4): 804-809.
- 22. Hunt CD. Dietary boron: progress in establishing essential roles in human physiology. *Journal of trace elements in medicine and biology* 2012; **26**(2-3): 157-160.
- Michel H, Behr J, Harrenga A, Kannt A. Cytochrome C oxidase: structure and spectroscopy. *Annual review of biophysics and biomolecular structure* 1998; 27: 329-356.
- 24. Maier T, Jacobi A, Sauter M, Bock A. The product of the hypB gene, which is required for nickel incorporation into hydrogenases, is a novel guanine nucleotide-binding protein. *Journal of bacteriology* 1993; **175**(3): 630-635.