

Molecular Identification and Antifungal Susceptibility Pattern of Non-*albicans* *Candida* Species Isolated from Vulvovaginal Candidiasis

Ziba Abbasi Nejat¹, Shirin Farahyar^{*2,3}, Mehraban Falahati², Mahtab Ashrafi Khozani², Aga Fateme Hosseini⁴, Azamsadat Faiazy⁵, Masoome Ekhtiari² and Saeideh Hashemi-Hafshenjani²

¹International Campus, Iran University of Medical Sciences, Tehran, Iran; ²Department of Medical Parasitology and Mycology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran; ³Microbial

Biotechnology Research Center, School of Medicine, Iran University of Medical Sciences, Tehran, Iran;

⁴Department of Biostatistics, School of Public Health, Iran University of Medical Sciences, Tehran, Iran; ⁵Department of Gynecology, Sayyad Shirazi Hospital, Golestan University of Medical Sciences, Gorgan, Iran

Received 18 October 2016; revised 4 January 2017; accepted 7 January 2017

ABSTRACT

Background: Vulvovaginal candidiasis (VVC) is an important health problem caused by *Candida* spp. The aim of this study was molecular identification, phylogenetic analysis, and evaluation of antifungal susceptibility of non-*albicans* *Candida* isolates from VVC. **Methods:** Vaginal secretion samples were collected from 550 vaginitis patients at Sayyad Shirazi Medical and Educational Center of Gorgan (Golestan Province, Iran) from May to October 2015. Samples were analyzed using conventional mycological and molecular approaches. Clinical isolates were analyzed with specific PCR using CGL primers, and the internal transcribed spacer region and the D1-D2 domain of the large-subunit rRNA gene were amplified and sequenced. Susceptibility to amphotericin B, fluconazole, itraconazole, and clotrimazole was determined by the guidelines of the Clinical and Laboratory Standard Institute. **Results:** In total, 35 non-*albicans* *Candida* isolates were identified from VVC patients. The isolates included 27 strains of *Candida glabrata* (77.1%), 5 *Candida krusei* (*Pichia kudriavzevii*; 14.3%), 2 *Candida kefyr* (*Kluyveromyces marxianus*; 5.7%), and 1 *Candida lusitanae* (*Clavispora lusitanae*; 2.9%). The fungicides itraconazole and amphotericin B were effective against all species. One isolate of *C. glabrata* showed resistance to fluconazole and clotrimazole, and 26 isolates of *C. glabrata* indicated dose-dependent susceptibility to fluconazole. *C. lusitanae* was susceptible in a dose-dependent manner to fluconazole and resistant to clotrimazole. **Conclusions:** Non-*albicans* *Candida* spp. are common agents of vulvovaginitis, and *C. glabrata* is the most common species in the tested patients. **DOI: 10.22034/ibj.22.1.33**

Keywords: *Candida glabrata*, Vulvovaginal candidiasis, *Candida krusei*

Corresponding Author: Shirin Farahyar

Department of Medical Parasitology and Mycology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran; Tel.: (+98-21) 88622653; Fax: (+98-21) 88602217; E-mail: farahyar.sh@iums.ac.ir

INTRODUCTION

The incidence of vulvovaginal candidiasis (VVC) caused by non-*albicans* *Candida* spp. has increased considerably^[1,2]. Based on evidence,

about two-thirds of women worldwide have experienced at least one episode of VVC during their lifetime^[3] and some with recurrent episodes^[1]. Recurrent episodes are more often caused by non-*albicans* *Candida* spp. against which azole antifungal

agents show low effectiveness^[4].

Candida albicans is the major cause of vulvovaginitis, and *Candida glabrata* and *Candida tropicalis* appear to be the most common non-*albicans* *Candida* species involved in this disease^[2,4-6]. Identification of many species of *Candida* by traditional methods is a challenge and is sometimes imprecise, particularly for uncommon microorganisms. Sequence analysis of the internal transcribed spacer (ITS) region of the rRNA gene and the D1-D2 domain of the large-subunit rRNA gene as well as PCR-RFLP on the ITS region have been used extensively for identification of fungal pathogens^[7-11].

Minimal inhibitory concentrations (MIC) of azoles against some strains of non-*albicans* *Candida* species are high, due to intrinsic resistance^[12]. The antifungal susceptibility of *Candida* spp. causing VVC varies^[12,13], and reliable identification and assessment of drug sensitivity of *Candida* vaginal isolates are of value in determining proper treatment. The aim of this study was molecular identification, phylogenetic analysis, and evaluation of antifungal susceptibility of non-*albicans* *Candida* isolates causing VVC.

MATERIALS AND METHODS

Patients and specimens collection

This study was conducted on 550 non-pregnant vaginitis patients referred to Sayyad Shirazi Medical and Educational Center of Gorgan (Golestan, Iran) from May to October 2015. Non-pregnant patients were entered to the study by a simple random sampling method. A questionnaire was completed for each patient about their age, the medical condition (recent antibiotic or antifungal therapy, urinary tract infections, immunodeficiency, diabetes, experiencing at least one episode or recurrent episodes of VVC), and other conditions. Specimens were obtained from vaginal mucosal discharge with a sterile cotton swab. The research protocol was approved by the Ethics Committee of Iran University of Medical Sciences (Tehran, Iran), under Ethics Committee number 93-04-198-25289.

Yeast identification

Microscopic examination was carried out to distinguish yeast forms or pseudohyphae. All samples were cultured on CHROMagar *Candida* (CHROMagar, France) for identification of mixed infections of *Candida* spp.^[14]. The isolates were identified by carbohydrate assimilation method using API 20C AUX system (Biomérieux, France)^[15].

DNA extraction

A single colony of each clinical isolate from CHROMagar *Candida* was subcultured on yeast extract peptone dextrose agar and incubated at 37 °C for 24-48 h. Genomic DNA was extracted from yeast cultures using the Qiagen DNA tissue kit (Germany). The extracted DNA was stored at -20 °C for further use.

Specific PCR

All clinical isolates with mauve, pink, or white colonies on CHROMagar *Candida* as well as *C. glabrata* CBS 138, as the reference strain, were analyzed by *C. glabrata*-specific PCR with CGL1-(5'-TTA TCA CAC GAC TCG ACA CT-3') and CGL2-(5'-CCC ACA TAC TGA TAT GGC CTA CAA-3')^[7] primers. The PCR thermal cycles were as follows: an initial denaturation at 96 °C for 5 min followed by 40 cycles of 30 s at 94 °C, 30 s at annealing temperature of 58 °C and 30 s at 72 °C. A final extension of 15 min at 72 °C was included at the end of PCR cycles.

Amplification and sequencing of ITS and D1-D2 regions

The universal primers ITS1 (5'-TCC GTA GGT GAA CCT GCG G -3') and ITS4 (5'-TCC TCC GCT TAT TGA TAT GC-3')^[7] were used to amplify the ITS1-5.8S-ITS2 region (annealing temperature 56 °C). Also, D1-D2 domain of 26S ribosomal RNA was amplified with NL1 (5'-GCA TAT CAA TAA GCG GAG GAA AAG-3') and NL4 (5'-GGT CCG TGT TTC AAG ACG G-3')^[16] primers by the following profile: 98 °C (5 min), 35 cycles of 98 °C (30 s), annealing temperature 60 °C (30 s), and 72 °C (30 s), followed by a final extension of 72 °C (5 min). The PCR products were sequenced by Macrogen (Korea). The resulting sequences were analyzed and compared with the reference data available from the GenBank database using the BLAST sequence search tool (<http://www.ncbi.nlm.nih.gov/BLAST>), and the results were submitted to the GenBank.

Phylogenetic analysis

The sequencing results of the D1-D2 and the ITS domains were analyzed and compared with the reference strains by neighbor-joining method using MEGA 7 (TreeView software).

Antifungal drug susceptibility testing

Tests of susceptibility to amphotericin B, fluconazole, clotrimazole, and itraconazole (Sigma, Germany) were conducted according to the Clinical and Laboratory Standards Institute (CLSI) guidelines (document M27-S3 and S4)^[17,18]. *C. glabrata* CBS 138 was used as the reference strain, and all tests were duplicated.

[Downloaded from ibj.pasteur.ac.ir on 2025-06-30]
[DOR: 20.1001.1.1028852.2018.22.1.3.6]
[DOI: 10.22034/ibj.22.1.33]

Table 1. Non-albicans *Candida* isolates and age distribution of vulvovaginal candidiasis patients

Ages (y)	<i>Candida</i> spp.			
	<i>C. glabrata</i>	<i>C. krusei</i>	<i>C. kefyr</i>	<i>C. lusitaniae</i>
<20	6	3	1	-
20-29	10	2	1	-
30-39	11	-	-	1

RESULTS

Patients

A total of 550 vaginal specimens of non-pregnant vaginitis patients were studied. Individuals with conditions such as infection by *Trichomonas vaginalis*, *Mycoplasma urealyticum*, or *Chlamydia*, as well as bacterial vaginosis or vulval skin disease were excluded from the study. In addition, 122 (22.2%) non-pregnant vaginitis patients showed VVC, and *C. albicans* isolates were identified in 87 (71.3%) VVC patients (data not shown). Non-albicans *Candida* isolates were found in 35 (28.7%) VVC patients aged 19-39 years from Gorgan (Table 1). All patients were negative for diabetes, immunodeficiencies, or any chronic disease and were not taken any antifungal treatment.

Yeast isolates

Thirty-five isolates of non-albicans *Candida* were obtained from 550 vulvovaginitis patients: 27 *C. glabrata* (77.1%), 5 *C. krusei* (*Pichia kudriavzevii*; 14.3%), 2 *C. kefyr* (*Kluyveromyces marxianus*; 5.7%), and 1 *Candida lusitaniae* (*Clavispora lusitaniae*; 2.9%) (Table 1).

Amplification with specific primers

The clinical isolates with mauve, pink, or white colonies on CHROMagar *Candida* and *C. glabrata* CBS 138 were analyzed with CGL1/2 specific primers, and the presence of the 423-bp fragment amplified with these primers confirmed those isolates identical to *C. glabrata* (Fig. 1).

PCR amplification and sequencing of ITS region and D1-D2 domain

PCR amplification of all clinical isolates with ITS1 and ITS4 primers yielded the fragments of 350-880 bp. The ITS fragments of *C. glabrata* were ~500 to ~879 bp (Fig. 2)^[8,10], while *C. krusei*, *C. kefyr*, and *C. lusitaniae* yielded the fragments of ~500, ~720, and ~370 bp, respectively (Fig. 2)^[8]. The ITS fragments of three clinical *C. glabrata* isolates showing ~500 and ~600 bp were compared to the reference data in the GenBank database using the BLAST. Three

C. glabrata isolates showed partial sequences of ITS region (~500 and ~600 bp), while the complete sequences of ITS region of *C. glabrata* was ~879 bp, and the partial sequences and complete sequences were submitted to the GenBank (Table 2). The D1-D2 region of the large-subunit rRNA gene amplified with NL1 and NL4 primers yielded the fragments of ~600 bp (Fig. 3). The ITS and D1-D2 region sequences of non-albicans *Candida* clinical isolates were compared to the reference data in the GenBank database using BLAST (<http://www.ncbi.nlm.nih.gov/BLAST>). All clinical isolates were correctly determined to species level. The sequences were submitted to the GenBank under accession numbers KU845721, KU904424-26, KU992386-95, KX008737-53, and KX016018-21 (Table 2).

Phylogenetic trees

The sequences of D1-D2 region were aligned for phylogenetic analysis. All *C. glabrata* strains showed 100% identity with KU729149, KU729145, and KU729137 reference strains. *C. krusei* (*Pichia kudriavzevii*) indicated 100% similarity to KU729202 and KU729201 reference strains. *C. kefyr* (*Kluyveromyces marxianus*) and *C. lusitaniae* (*Clavispora lusitaniae*) displayed 100% identity with KM279378 and KP070758 reference strains, respectively (Fig 4). Phylogenetic analysis of sequences corresponding to the ITS region demonstrated that all strains of the species were identical to the reference strains. *C. glabrata* strains showed similarity to KP675206, KP131703, KP675517, and LT577613 but *C. krusei* (*Pichia kudriavzevii*) to KX833111 and KX015902 reference




Fig. 1. The genomic DNA of clinical isolates of *Candida glabrata* and *C. glabrata* CBS 138 were analyzed with PCR using CGL1/2 specific primers, and a 423-bp fragment produced. Isolates 1, 2, 3, 4, 5, 6, and 7, *C. glabrata*; S, *C. glabrata* CBS 138 (as standard); M, marker 100 bp; N, negative control.

[Downloaded from ibj.pasteur.ac.ir on 2025-06-30]
[DOR: 20.1001.1.1028852.2018.22.1.3.6]
[DOI: 10.22034/ibj.22.1.33]




Fig. 2. Amplification of genomic DNA from clinical isolates using ITS1 and ITS4 primers. Isolates 1 and 8, *C. krusei* (500 bp); isolates 2, 3, 4, 5, and 6, *Candida glabrata* producing an ~879-bp fragment; isolate 7, *C. glabrata* CBS 138 (as standard); isolate 9, *Candida lusitanae* producing a ~370-bp fragment; M, marker 100 bp; N, negative control.

Table 2. Accession numbers of clinical isolates

Clinical isolates	Accession no.
<i>C. glabrata</i>	KU845721
	KU904426
	KU992388
	KU992389
	KU992390
	KU992391
	KU992392
	KU992393
	KU992394
	KU992395
	KX008737
	KX008738
	KX008739
	KX008740
	KX008741
	KX008744
	KX008745
	KX008748
	KX008749
	KX008750
	KX008751
<i>Pichia kudriavzevii</i> (<i>Candida krusei</i>)	KX008752
	KX008753
	KX016018
	KX016019
<i>Kluyveromyces marxianus</i> (<i>Candida kefir</i>)	KX016020
	KX016021
	KU904424
	KU992387
<i>Clavispora lusitanae</i> (<i>Candida lusitanae</i>)	KX008742
	KX008743
	KX008746

strains. *C. kefir* (*Kluyveromyces marxianus*) indicated identity with KJ849337 and KJ849335 reference strains, while *C. lusitanae* (*Clavispora lusitanae*) showed similarity to KP674503 reference strain (Fig. 5).

Antifungal drug susceptibility

Results of susceptibility testing of the 35 non-*albicans* *Candida* isolates showed one isolate of *C. glabrata* to be resistant to fluconazole (MIC \geq 64 μ g/ml) and clotrimazole (MIC \geq 4 μ g/ml), and 26 isolates of *C. glabrata* were susceptible to fluconazole (MIC \leq 32 μ g/ml) in a dose-dependent manner^[18]. Single clinical isolate of *C. lusitanae* showed dose-dependent susceptibility to fluconazole (MIC = 16-32 μ g/ml) and resistant to clotrimazole (MIC = 2 μ g/ml)^[17]. The MICs for one isolate of *C. krusei* were as follows: fluconazole \leq 32 μ g/ml and clotrimazole = 2 μ g/ml. Because the clinical isolates of *C. krusei* showed intrinsic resistant to fluconazole, and their MICs should not be interpreted using this scale; therefore, breakpoint was not provided by CLSI document M27-S4^[18]. Itraconazole and amphotericin B were active against all of the isolates (Table 3).

DISCUSSION

This study revealed that non-*albicans* *Candida* spp., as important agents, are commonly associated with vulvovaginitis; *C. glabrata* is the second in rate of occurrence after *C. albicans*. Other investigations have found that *C. glabrata* and *C. albicans* to be the most common species isolated from VVC patients^[13,19,20]. The overall proportion of non-*albicans* infection in vaginitis has been reported to be high^[2]. *C. glabrata*




Fig. 3. The D1-D2 region of clinical isolates amplified with NL1 and NL4 primers, yielded fragments ~600 bp. Isolate 1, *C. glabrata* CBS 138 (as standard); isolates 2, 3, 4, 5, 6, and 7, *Candida glabrata*; M, marker 100 bp; N, negative control.

and *C. tropicalis* have also been found in the normal vaginal flora of women in China^[21]. Infections caused by less common yeasts have been increasingly observed^[22], and identification of a variety of medically important yeast species by traditional approaches may be challenging. Molecular methods can improve discrimination of uncommon clinical isolates and closely related yeast species such as those in *Candida* complexes. Molecular diagnostics are also useful in carrying out large epidemiological studies of pathogenic yeasts. In this study, conventional methods and specific PCR with CGL primers were used for identification of *C. glabrata*. Sequencing the ITS and D1-D2 regions has proven to be a feasible method for the reliable identification of clinically important yeasts,





Fig. 4. Molecular phylogenetic analysis using Neighbor-Joining method with sequences of D1-D2 domain. The evolutionary history was inferred using the Neighbor-Joining method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. Bootstrap values greater than 50% from 1000 replicates are indicated at the nodes. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. Evolutionary analyses were conducted in MEGA7. (A) Accession numbers of *Candida glabrata* isolated in this study and reference strains (KU729149, KU729145, and KU729137), (B) accession numbers of *Kluyveromyces marxianus* (*Candida kefir*) isolated in this study and reference strain (KM279378), (C) Accession number of *Clavispora lusitanae* (*Candida lusitanae*) isolated in this study and reference strain (KP070758), (D) accession numbers of *Pichia kudriavzevii* (*Candida krusei*) isolated in this study (Table 1) and reference strains (KU729202 and KU72920).



principally the *C. glabrata* complex (*C. glabrata*, *C. bracarensis*, and *C. nivariensis*)^[22,23] and the *C. parapsilosis* complex (*C. parapsilosis*, *C. orthopsilosis*, and *C. metapsilosis*)^[24]. Richter *et al.*^[13] reported that 173 of 593 yeast isolates from vaginitis patients were non-*albicans* *Candida* spp., and that *C. albicans* was the most frequent cause of vaginal candidiasis, followed by *C. glabrata*, *C. parapsilosis*,

C. krusei, *Saccharomyces cerevisiae*, *C. tropicalis*, and *C. lusitaniae*^[13]. Vijaya *et al.*^[25] showed that *C. tropicalis* is the major non-*albicans* species of *Candida* associated with vaginal candidiasis. Other studies in Iran introduced *C. glabrata* as the most important non-*albicans* species in vaginal candidiasis patients^[19,20]. Shi *et al.*^[26] demonstrated that *C. albicans* is the main cause of vaginal candidiasis, followed by *C. glabrata*,

Table 3. Drug treatment susceptibility of *Candida* spp. isolated from vulvovaginal candidiasis patients

Clinical isolates	Fluconazole			Itraconazole			Clotrimazole			Amphotericin B		
	S	R	S-DD	S	R	S-DD	S	R	S-DD	S	R	S-DD
	n			n			n			n		
<i>C. glabrata</i> (n = 27)	0	1	26	27	0	0	26	1	0	27	0	0
<i>C. krusei</i> (n = 5)	--*	--	--	5	0	0	4	1	0	5	0	0
<i>C. kefyr</i> (n = 2)	2	0	0	2	0	0	2	0	0	2	0	0
<i>C. lusitaniae</i> (n = 1)	0	0	1	1	0	0	0	1	0	1	0	0

S, sensitive; R, resistant; S-DD, susceptible, dose-dependent; *Because of the clinical isolates of *C. krusei* showed intrinsically resistant to fluconazole; therefore, breakpoint is not provided by Clinical and Laboratory Standards Institute document M27-S4.

C. tropicalis and *C. parapsilosis* in China. In the current study, *C. glabrata* was the most common species of non-albicans *Candida*. *Candida* spp., especially *C. glabrata* and *C. albicans*, represent a primary source of infection leading to bloodstream infections and to morbidity and mortality in severely affected and immune-compromised individuals^[12,27-29]. Species *C. krusei* and *C. glabrata* have been indicated. To be resistant or to have low susceptibility to azole drugs^[4,12]. *C. lusitaniae* has also been shown to have resistance to amphotericin B, caspofungin, and azoles^[30]. A study in Japan revealed that one of the 19 *C. glabrata* clinical isolates of VVC patients showed resistance to fluconazole, and this isolate demonstrated cross-resistance to other antimycotic drugs tested^[31]. Another study displayed that non-albicans isolates, particularly *C. glabrata* strains, were susceptible in a dose-dependent manner and were resistant to fluconazole^[13]. Kalkanci et al.^[32] suggested that *C. glabrata* was the most vaginal isolate of non-albicans *Candida*, and 3 of 81 (3.7%) *C. glabrata* isolates were resistant to ketoconazole, and only one *C. glabrata* was fluconazole resistant. Also, five *C. glabrata* isolates showed susceptibility to fluconazole in a dose-dependent manner. A previous study indicated that nystatin was an appropriate option instead of imidazoles^[33]. An investigation from Iran found that clinical isolates of *Candida* spp. were susceptible to clotrimazole, miconazole, and nystatin^[19]. Razzaghi-Abyaneh et al.^[34] indicated that itraconazole was the most effective antimycotic drug for *C. krusei*, *C. glabrata*, and *C. guilliermondii* isolates of superficial candidiasis in Iran. In the current study, 26 isolates of *C. glabrata* were susceptible dose-dependent to fluconazole, and one isolate was resistant to fluconazole and clotrimazole. In addition, one isolate of *C. lusitaniae* was susceptible to fluconazole in a dose-dependent manner and resistance to clotrimazole^[35,36].

The phylogenetic analyses of the D1-D2 and the ITS domains indicated that clinical isolates of vaginal

candidiasis are genetically similar to reference *Candida* species. The phylogenetic analyses of the D1-D2 domain revealed that all *C. glabrata* isolates had 100% similarity to KU729149 (ATCC 90030), KU729145 (ATCC 66032), and KU729137 (ATCC 2001) reference strains. Clinical isolates of *C. krusei* (*Pichia kudriavzevii*) showed 100% identity with KU729202 (ATCC 34135) and KU729201 (ATCC 14243) reference strains. *C. kefyr* (*Khuyveromyces marxianus*) and *C. lusitaniae* (*Clavispora lusitaniae*) were similar to KM279378 (isolate U-MF11) and KP070758 (isolate 0Q10) reference strains, respectively. Based on the phylogenetic analyses of the ITS region, *C. glabrata* clinical isolates showed similarity to KP675206 (strain m36b), KP131703 (CNRMA6.53 isolate ISHAM-ITS_ID MITS649), KP675517 (strain M310B), and LT577613 (strain IQBasrah28) reference strains in the GenBank databases. *C. krusei* (*Pichia kudriavzevii*) showed similarity to KX833111 (strain DMic 165166) and KX015902 reference strains. *C. kefyr* (*Khuyveromyces marxianus*) amplified sequences matched completely with the corresponding sequences of the KJ849337 (strain ZT-Kma.4) and KJ849335 reference strains. *C. lusitaniae* (*Clavispora lusitaniae*) indicated 96% identity with KP674503 (strain B157B) reference strain. The phylogenetic trees were created using the sequences of different *Candida* clinical isolates and showed the formation of separate branches for each species.

Sequencing of the ITS region and D1-D2 domain appears to be the most effective method for identification of *Candida* spp. The phylogenetic trees based on sequences of D1-D2 and ITS domains showed similarity of *Candida* spp. to closely related reference species. Results suggested that amphotericin B and itraconazole retain good clinical effectiveness. Accurate identification and assessment of susceptibility of *Candida* spp. isolates are critical to treatment management, since some strains showed varying degrees of resistance to antifungal drugs.

ACKNOWLEDGEMENTS

This research was supported by Iran University of Medical Sciences, Tehran, Iran (grant no. 25289).

CONFLICT OF INTEREST. None declared.

REFERENCES

- MacNeill C, Carey JC. Recurrent vulvovaginal candidiasis. *Current women's health reports* 2001; **1**(1): 31-35.
- Paulitsch A, Weger W, Ginter-Hanselmayer G, Marth E, Buzina W. A 5-year (2000-2004) epidemiological survey of *Candida* and non-*Candida* yeast species causing vulvovaginal candidiasis in Graz, Austria. *Mycoses* 2006; **49**(6): 471-475.
- Sobel JD. Vulvovaginal candidosis. *Lancet* 2007; **369**(9577): 1961-1971.
- Deorukhkar SC, Saini S, Mathew S. Non-*albicans Candida* Infection: an emerging threat. *Interdisciplinary perspectives on infectious diseases* 2014; **2014**: 615958.
- Mahmoudi Rad M, Zafarghandi S, Abbasabadi B, Tavallaee M. The epidemiology of *Candida* species associated with vulvovaginal candidiasis in an Iranian patient population. *European journal of obstetrics, gynecology, and reproductive biology* 2011; **155**(2): 199-203.
- Diba K, Namaki A, Ayatollahi H, Hanifian H. Rapid identification of drug resistant *Candida* species causing recurrent vulvovaginal candidiasis. *Medical mycology journal* 2012; **53**(3): 193-198.
- Luo G, Mitchell TG. Rapid identification of pathogenic fungi directly from cultures by using multiplex PCR. *Journal of clinical microbiology* 2002; **40**(8): 2860-2865.
- Fujita SI, Senda Y, Nakaguchi SH, Hashimoto T. Multiplex PCR using internal transcribed spacer 1 and 2 regions for rapid detection and identification of yeast strains. *Journal of clinical microbiology* 2001; **39**(10): 3617-3622.
- Boyanton BL Jr, Luna RA, Fasciano LR, Menne KG, Versalovic J. DNA pyrosequencing-based identification of pathogenic *Candida* species by using the internal transcribed spacer 2 region. *Archives of pathology and laboratory medicine* 2008; **132**(4): 667-674.
- Korabecna M. The variability in the fungal ribosomal DNA (ITS1, ITS2, and 5.8 S rRNA gene): its biological meaning and application in medical mycology. *Communicating current research and educational topics and trends in applied microbiology* 2007; **105**: 783-787.
- Mohammadi R, Mirhendi H, Rezaei-Matehkolaei A, Ghahri M, Shidfar MR, Jalalizand N, Makimura K. Molecular identification and distribution profile of *Candida* species isolated from Iranian patients. *Medical mycology* 2013; **51**(6): 657-663.
- Pfaller MA, Castanheira M, Lockhart SR, Jones RN. *Candida glabrata*: multidrug resistance and increased virulence in a major opportunistic fungal pathogen. *Current fungal infection reports* 2012; **6**(3): 154-164.
- Richter SS, Galask RP, Messer SA, Hollis RJ, Diekema DJ, Pfaller MA. Antifungal susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent cases. *Journal of clinical microbiology* 2005; **43**(5):2155-2162.
- Gültekin B, Yazici V, Aydin N. Distribution of *Candida* species in vaginal specimens and evaluation of CHROMagar *Candida* medium. *Mikrobiyoloji bulteni* 2005; **39**(3): 319-324.
- Buesching WJ, Kurek K, Roberts GD. Evaluation of the modified API 20C system for identification of clinically important yeasts. *Journal of clinical microbiology* 1979, **9**(5): 565-569.
- Linton CJ, Borman AM, Cheung G, Holmes AD, Szekely A, Palmer MD, Bridge PD, Campbell CK, Johnson EM. Molecular identification of unusual pathogenic yeast isolates by large ribosomal subunit gene sequencing: 2 years of experience at the United kingdom mycology reference laboratory. *Journal of clinical microbiology* 2007; **45**(4): 1152-1158.
- Clinical and Laboratory Standards Institute (CLSI). Reference method for broth dilution antifungal susceptibility testing of yeasts; Third informational supplement. CLSI document M27-S3. Clinical and Laboratory Standards Institute, Wayne, Pennsylvania 2008; **28**.
- Clinical and Laboratory Standards Institute (CLSI). Reference method for broth dilution antifungal susceptibility testing of yeasts; fourth informational supplement. CLSI document M27-S4. Clinical and Laboratory Standards Institute, Wayne, Pennsylvania 2012.
- Zarei Mahmoudabadi A, Najafyan M, Alidadi M. Clinical study of *Candida* vaginitis in Ahvaz, Iran and susceptibility of agents to topical antifungal. *Pakistan journal of medical sciences* 2010; **26**(3): 607-610.
- Hedayati MT, Taheri Z, Galinimoghadam T, Aghili SR, Yazdani Cherati J, Mosayebi E. Isolation of different species of *Candida* in patients with vulvovaginal candidiasis from sari, Iran. *Jundishapur journal of microbiology* 2015; **8**(4): e15992.
- Wei YP, Feng J, Luo ZC. Isolation and genotyping of vaginal non-*albicans Candida* spp. in women from two different ethnic groups in Lanzhou, China. *International journal of gynaecology and obstetrics* 2010; **110**(3): 227-230.
- Lockhart SR, Messer SA, Gherna M, Bishop JA, Merz WG, Pfaller MA, Diekema DJ. Identification of *Candida nivariensis* and *Candida bracarenensis* in a large global collection of *Candida glabrata* isolates: comparison to the literature. *Journal of clinical microbiology* 2009; **47**(4): 1216-1217.
- Borman AM, Petch R, Linton CJ, Palmer MD, Bridge PD, Johnson EM. *Candida nivariensis*, an emerging pathogenic fungus with multidrug resistance to antifungal agents. *Journal of clinical microbiology* 2008; **46**(3): 933-938.
- Tavanti A, Davidson AD, Gow NA, Maiden MC, Odds

- FC. *Candida orthopsilosis* and *Candida metapsilosis* spp. nov. to replace *Candida parapsilosis* groups II and III. *Journal of clinical microbiology* 2005; **43**(1): 284-292.
25. Vijaya D, Dhanalakshmi TA, Kulkarni S. Changing trends of vulvovaginal candidiasis. *Journal of laboratory physicians* 2014; **6**(1): 28-30.
 26. Shi XY, Yang YP, Zhang Y, Li W, Wang JD, Huang WM, Fan YM. Molecular identification and antifungal susceptibility of 186 *Candida* isolates from vulvovaginal candidiasis in southern China. *Journal of medical microbiology* 2015; **64**(Pt 4): 390-393.
 27. Taj-Aldeen SJ, AbdulWahab A, Kolecka A, Deshmukh A, Meis JF, Boekhout T. Uncommon opportunistic yeast bloodstream infections from Qatar. *Medical mycology* 2014; **52**(5): 552-556.
 28. Pfaller MA, Messer SA, Moet GJ, Jones RN, Castanheira M. *Candida* bloodstream infections: comparison of species distribution and resistance to echinocandin and azole antifungal agents in Intensive Care Unit (ICU) and non-ICU settings in the SENTRY Antimicrobial Surveillance Program (2008-2009). *International journal of antimicrobial agents* 2011; **38**(1): 65-69.
 29. Arendrup MC. *Candida* and candidaemia. Susceptibility and epidemiology. *Danish medical journal* 2013; **60**(11): B4698.
 30. Asner SA, Giulieri S, Diezi M, Marchetti O, Sanglard D. Acquired Multidrug Antifungal Resistance in *Candida lusitanae* during Therapy. *Antimicrobial agents and chemotherapy* 2015; **59**(12): 7715-7722.
 31. Nagashima M, Yamagishi Y, Mikamo H. Antifungal susceptibilities of *Candida* species isolated from the patients with vaginal candidiasis. *Journal of infection and chemotherapy* 2016; **22**(2):124-126.
 32. Kalkanci A, Güzel A, Jabban I, Aydin M, Ilkit M, Kuştimur S. *Candida* vaginitis in non-pregnant patients: a study of antifungal susceptibility testing and virulence factors. *Journal of obstetrics and gynaecology* 2013; **33**(4): 378-383.
 33. Choukri F, Benderdouche M, Sednaoui P. In vitro susceptibility profile of 200 recent clinical isolates of *Candida* spp. to topical antifungal treatments of vulvovaginal candidiasis, the imidazoles and nystatin agents. *Journal de mycologie médicale/journal of medical mycology* 2014; **24**(4): 303-307.
 34. Razzaghi-Abyaneh M, Sadeghi G, Zeinali E, Alirezaee M, Shams-Ghahfarokhi M, Amani A, Mirahmadi R, Tolouei R: Species distribution and antifungal susceptibility of *Candida* spp. isolated from superficial candidiasis in outpatients in Iran. *Journal de mycologie médicale/journal of medical mycology* 2014; **24**(2): e43-e50.
 35. Costa C, Ribeiro J, Miranda IM, Silva-Dias A, Cavalheiro M, Costa-de-Oliveira S, Rodrigues AG, Teixeira MC. Clotrimazole drug resistance in *Candida glabrata* clinical isolates correlates with increased expression of the drug: H⁺ antiporters CgAqr1, CgTpo1_1, CgTpo3, and CgQdr2. *Frontiers in microbiology* 2016; **7**:526.
 36. Pelletier R, Peter J, Antin C, Gonzalez C, Wood L, Walsh TJ: Emergence of resistance of *Candida albicans* to clotrimazole in human immunodeficiency virus-infected children: *in vitro* and clinical correlations. *Journal of clinical microbiology* 2000; **38**(4):1563-1568.