

Association of a New Germline Variant in the *MUTYH* DNA Glycosylase Gene with Colorectal Adenoma Transformation into Malignancy

Amjad Mahasneh^{1*}, Fawaz N. Al-Shaheri² and Mohammed N. BaniHani³

¹Dept. of Applied Biological Sciences, Jordan University of Science and Technology, Irbid 22110, Jordan

²Dept. of Medical Laboratory Sciences, Jordan University of Science and Technology, Irbid 22110, Jordan

³Dept. of General Surgery and Urology, Jordan University of Science and Technology, Irbid 22110, Jordan

Received 1 September 2018; revised 6 November 2018; accepted 13 November 2018

ABSTRACT

Background: *MUTYH* DNA glycosylase germline mutations are linked to the recessive inheritance of multiple adenoma. Studies have revealed that germline mutations in this gene are ethnicity related. This study aimed to identify the germline mutations in *MUTYH* gene and determine their prevalence among Jordanian patients with colorectal adenoma. **Methods:** In this study, 150 colorectal adenoma patients and 150 cancer-free individuals with no previous history of polyps were recruited. Sanger DNA sequencing of the *MUTYH* gene (accession number NG_008189.1) was carried out using 3130xL Genetic Analyzer. Sequencing results were analyzed by ChromasPro, and mutational effects were predicted by online bioinformatics tools. **Results:** Two novel variants, g.87C>T and c.1264G>C, were identified. g.87C>T was also found in 60 (40%) patients and 10 (6.7%) controls. However, c.1264G>C was detected in 90 (60%) patients and 7 (4.7%) controls. Thus, a significant association was observed between these two variants and colorectal adenoma (*p* value for both variants was <0.0001). Moreover, the newly identified germline variant, c.1264G>C, was found to be significantly associated with colorectal adenoma transformation into malignancy (*p* < 0.0001). **Conclusion:** The data showed high prevalence of two germline mutations in *MUTYH* gene among Jordanians with colorectal adenoma, which may make them as potential early biomarkers for diagnosis of colorectal adenoma. **DOI:** 10.29252/ibj.23.6.412

Keywords: Colorectal adenoma, Germline mutations, *MUTYH* gene

Corresponding Authors: Amjad Mahasneh

Department of Applied Biological Sciences, Jordan University of Science and Technology, Irbid 22110, Jordan; E-mail: amjada@just.edu.jo

INTRODUCTION

Colorectal adenoma pathogenesis is a multistep process and can induce neoplastic transformation of normal cryptic epithelial cells into neoplasia^[1]. This stepwise transformation is due to multiple genetic and environmental factors, which disrupt the epithelial cells homeostasis by diminishing cellular apoptosis, persisting DNA replication, and losing the ability for differentiation and maturation^[2-4]. Colorectal adenoma can be classified into tubular, villous, and tubovillous adenoma^[5]. The age, gender,

smoking, diet, and obesity are environmental factors that contribute to colorectal neoplasia^[6-8]. Genetic mechanisms such as chromosomal instability, CpG island hypermethylation, and microsatellite instability assist in colorectal neoplastic transformation. Microsatellite instability results from defect in DNA mismatch repair system that includes *PCNA*, *MLH1*, *MSH6*, *XRCC1*, and *MUTYH* genes^[9-11].

MUTYH comprises 16 exons encoding 535 amino acids protein and is mapped to the chromosome 1(1p32.1-p34.3)^[12]. The *MUTYH* glycosylase, a base excision repair enzyme^[13], detects and repairs DNA

damage including those generated by normal metabolic reactions such as alkylation, deamination, or oxidation^[14]. The exposure of reactive oxygen species (produced during aerobic metabolism) to some chemicals/radiation influence the DNA integrity^[15]. As a consequence, 7,8-dihydroxy-8-oxoguanine (8-oxo-G) often pairs with adenine (A), resulting in transversion of guanine:cytosine into thymidine:adenine (G:C>T:A) after two replication rounds of DNA^[16]. The 8-oxo-G:A base-pair detection is initiated by MUTYH glycosylase^[14].

MUTYH germline mutations have been linked to multiple colorectal adenomas inheritance in Caucasian populations^[17]. Therefore, their identification greatly enhances our understanding of the colorectal cancer (CRC) potential causes and attributes in implementation of screening and management measures. Importantly, the early detection of CRC can be effective by establishing panels of genetic biomarkers. There are many genes and proteins currently used in the clinical settings in USA^[18], but they differ significantly in their predictive and prognostic values among populations^[19]. Supposing that Jordanian patients with colorectal adenoma have a novel germline mutation(s) in *MUTYH* gene, this study aimed to identify these mutations among those patients and to determine their prevalence in relation to their ethnic background.

MATERIALS AND METHODS

Study subjects

Colorectal adenoma patients (n = 150) were recruited from King Abdulla University Hospital (Ramtha, Jordan) during the period between January 2016 and February 2017. To select these patients, the biopsy samples from 400 subjects who visited the Endoscopy Unit at King Abdullah University Hospital were examined by a pathologist. Subjects (n = 150) who were confirmed pathologically to have tubular adenoma, villous adenoma, or tubulovillous adenoma were investigated in the study. Inclusion criteria include the number of polyps and the patient's previous treatment status. Patients with more than 10 polyps and without any surgical removal of polyps were enrolled in this study. However, those with a previous history of irritable bowel syndrome or Crohn's disease were excluded. Clinical data were collected from patient's history files. The study was approved in advance by the Institution Review Board at King Abdullah University Hospital and Jordan University of Science and Technology (Ramtha, Jordan). Informed written consents were obtained from all participants. Besides,

150 cancer-free individuals with no previous family history of polyps were recruited as controls. A structured questionnaire interview was established to collect data on the characteristics of the study participants.

Sample collection and handling

One blood sample was collected from each participant. Five milliliters of whole blood was collected in EDTA tubes. Samples were transported on ice to the DNA extraction laboratory at Princess Haya Biotechnology Center (Ramtha, Jordan) and either processed immediately or stored at 4 °C and extracted the next day.

Genomic DNA extraction

DNA was extracted from whole blood samples using Qiagen® genomic DNA purification kit (Qiagen, USA) according to the manufacturer's instructions. The concentration of DNA was determined using Nano Drop 2000 (Thermo Scientific, USA). To check the quality of extracted DNAs, two microliters from each extracted DNA sample was subjected to agarose gel electrophoresis. Extracted DNA samples were stored at -80 °C.

Polymerase chain reaction (PCR) procedure

All exons of *MUTYH* gene were amplified using PCR. The primers were designed using Primer3 Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/>). Eight amplicons were designed comprising the whole coding regions of *MUTYH* gene as well as the 5' untranslated region and exon-intron boundaries (Fig. 1). Amplification of the target sequences was performed using conventional PCR (Veriti™ Thermal Cycler from Applied Bioscience). PCR was carried out in a reaction volume of 25 µL containing 12.5 µL commercial 2× Master Mix (2.5 U *Taq* of DNA polymerase, 3 mM of MgCl₂, and 0.5 mM of dNTPs), and buffer (100 mM of KCl, 20 mM of Tris HCl [pH8.3]), as well as 2µL of each forward and reverse primer (10µM), 6.5 µl of nuclease-free water, and 2-3 µl of sample DNA. The list of primers and sequences for each amplicon are shown in Table 1.

Gel electrophoresis

Five microliters from each PCR product was loaded on 2% agarose gel. The electrophoresis was run at 140 volts for 40 minutes. The running Tris/Borate/EDTA buffer was diluted to 1×, and the bands were then detected under a UV light in Gel Doc™ XR system (BioRad, USA) using Red Safe stain (iNtRON). Images were processed and analyzed by Quantity One 4.6.9 software.

[Downloaded from ibj.pasteur.ac.ir on 2025-08-24]
[DOR: 20.1001.1.1028852.2019.23.6.3.3]
[DOI: 10.29252/ibj.23.6.412]




Fig. 1. Schematic representation of the designed *MUTYH* amplicons. *MUTYH* 5'untranslated region, 16 exons, and exons-intron boundaries were amplified in eight amplicons.

Cycler sequencing, cleaning and Sanger sequencing

Purified PCR products were cycle-sequenced using Big Dye Terminator Ready Reaction (R-R) mix. The reaction components were 4 μ L of sequencing buffer, 4 μ L of nuclease-free water, 1 μ L of R-R, 1 μ L of forward or reverse primer, and 4 μ L of purified DNA. The excess of Dye DeoxyTM terminator, primer dimmers, and the other impurities were removed from DNA sequencing reaction using Qiagen cleaning kit. Sequencing of cleaned cycle-sequenced products was carried out on an ABI prism 3130xL Genetic Analyzer (Applied Biosystems, USA) in the Genomics Sequencing Laboratory at Princess Haya Biotechnology Center (PHBC).

Sequencing data analysis

Sequence analysis was carried out using ChromasPro software (<http://technelysium.com.au/wp/chromas/>).

Bioinformatics and computational analysis of variants

Several bioinformatics tools were used in this study. Primer3Plus online software ([<http://www.primer3plus.com/>\) was used for primer design. MutationTaster software \(<http://www.mutationtaster.org/>\). PROVEAN \(<http://provean.jcvi.org/index.php>\) was applied for mutational effect prediction and PolyPhen-2 \(<http://genetics.bwh.harvard.edu/pph2/>\) for protein prediction analyses.](http://www.</p></div><div data-bbox=)

Statistical analysis

GraphPad Prism 6.0 was used for statistical analyses. In particular, *Chi-square* test was used. Statistical analyses with *p* value less than 0.05 were considered statistically significant.

RESULTS

Characteristics of the study participants

The recruited Jordanian subjects with colorectal adenoma were 81(54%) males and 69 (46%) females, and the controls were 78 (52%) males and 72 (48%) females. There was no age difference between the

Table 1. Primer sequences for *MUTYH* amplicons

Amplicon	Reverse primer	Forward primer	Size (bp)
1	ctttgatcacaacgctcaa	aggagacggaccgcaagt	410
2	gccaaagagtaaaccgtgag	cttgggccacaacctagtc	300
3	catactgccacagctgct	ggtctgacccatgacccttc	590
4	tggctatagaagtgcctaca	accccaacatcctaccagag	593
5	gcagtgttcctctctttagg	cacgcccagtatccaggta	690
6	agggaatcggcagctgag	aagggaagtacaacaagacaacaaa	459
7	acaaaagtactgggacatgaagt	cctggagtggagaatgtca	239
8	gctctacagcattccaggcta	aatcacttgaggccagaatca	410

Table 2. General characteristics of the participants (n = 300)

Variable	Control n (%)	Patient n (%)	p value
Gender			
Male	78 (52)	81 (54.0)	0.73*
Female	72 (48)	69 (46.0)	
Total	150	150	
Age groups (y)			
30-50	84 (56)	80 (53.3)	0.37
51-70	45 (30)	40 (26.7)	
<60	21 (14)	30 (20.0)	

*Odds ratio: 0.922 and relative risk: 0.960

patients and controls ($p = 0.37$, Table 2). Tubular adenoma was the most prevalent histopathological group among 103 (68.6%) patients. Also, 28 subjects (18.6%) had villous adenoma, and 19 (12.7%) had tubovillous adenoma. There was no significant difference in gender among the study subjects ($p = 0.29$; Table 3). Regarding the progression of colorectal adenoma into CRC, 107 (71.4%) subjects had progressed to CRC, while 43 (28.6%) had recovered after polypectomy. In addition, there was no significant association between CRC progression and gender ($p = 0.31$). On the other hand, 86 (80.3%) patients, who progressed into CRC, continued to metastasis, compared to 21 (19.7%) who recovered after surgery and chemotherapy ($p = 0.28$).

Mutational analyses

Interestingly, we identified five germline variants (g.87C>T, c.381A>C, c.724G>A, c.1264G>C, and c.1760C>T) in Jordanian subjects with colorectal adenoma (Table 4); two of which (g.87C>T and c.1760C>T) were novel. The c.1264G>C was the most common variant found in 90 patients (60%) with

colorectal adenoma. Three variants (c.381A>C, c.724G>A, and c.1264G>C) were found to be disease predisposing, and two variants (g.87C>T and c.1760C>T) were considered as benign (Table 4). Based on Polyphen-2 protein prediction pipeline, c.724G>A and c.1264G>C were probably damaging to DNA glycosylase. However, c.381A>C and c.1760C>T were benign. These findings are consistent with those obtained from PROVEAN v1.1.3 software (http://provean.jcvi.org/seq_submit.php), which indicates that c.724G>A and c.1264G>C are predicted to be deleterious to DNA glycosylase.

The heterozygous variant g.87C>T was identified in amplicon 1 in 60 subjects with colorectal adenoma and in 11 controls. Searching the ExAc and 1000 genome databases showed that g.87C>T was a novel variant found in this study. The variant was mapped to the

Table 4. Genotype and allele frequencies of g.87C>T in *MUTYH* gene among study subjects (n=300).

Genotypes and alleles	Control n (%)	Patient n (%)	p value
C/C	140 (93.3)	90 (60)	<0.0001
C/T	10 (6.7)	60 (40)	
T/T	0 (0)	0 (0)	
Total	150	150	
Allele C	290 (96.7)	240 (80)	<0.0001*
Allele T	10 (3.3)	60 (20)	
Total	300	300	
Familial	1 (10)	4 (6.7)	0.548**
Sporadic	9 (90)	56 (93.3)	
Total	10	60	

Odds ratio: 7.25 and relative risk: 4.26; **odds ratio: 0.642 and relative risk: 0.928

Table 3. Clinical characteristics of the patients (n=150)

Variable	Male n (%)	Female n (%)	Total n (%)	p value
Histopathology				
Tubular adenoma	60 (58.25)	43 (41.17)	103 (68.6)	0.29
Villous adenoma	13 (46.4)	15 (53.6)	28 (18.6)	
Tubovillous adenoma	8 (42.10)	11 (57.9)	19 (12.7)	
Total	81	69	150	
CRC progression				
Yes	55 (51.5)	52 (48.5)	107 (71.4)	0.31*
No	26 (60.4)	17 (39.6)	43 (28.6)	
Total	81	69	150	
Metastasis				
Yes	42 (48.9)	44 (51.1)	86 (80.3)	0.28**
No	13 (61.9)	8 (38.1)	21 (19.7)	
Total	55	52	107	

*Odds ratio: 0.692 and relative risk: 0.850; **odds ratio: 0.58 and relative risk: 0.788

Table 5. Genotype and allele frequencies of germline mutation c.1264G>C in *MUTYH* gene in study subjects (n = 300)

Genotypes and Alleles	Control n (%)	Patient n (%)	p value
G/G	143 (95.3)	60 (40)	<0.0001
G/C	7 (4.7)	5 (3.3)	
C/C	0 (0)	85 (56.7)	
Total	150	150	
Allele G	293 (97.7)	125 (41.7)	<0.0001*
Allele C	7 (3.3)	175 (58.3)	
Total	300	300	
Familial	6 (85)	63 (70)	0.376**
Sporadic	1 (15)	27 (30)	
Total	7	90	

*Odds ratio: 58.6 and relative risk: 18.22; **Odds ratio: 2.57 and relative risk: 2.43.

5' untranslated region (5'UTR) of *MUTYH* gene (Fig. 2) and was named according to Human Genome Variation Society guidelines. Table 5 shows that g.87C>T is significantly associated with colorectal adenoma in comparison to the controls ($p < 0.0001$). Statistical analysis of allele frequencies of both groups showed that the wild-type allele (C) was more common in the control group (96.7%) than the patients group (80%). However, the mutant allele (T) was found in 20% among patients compared to 3.3% among the controls. These data are statistically significant with $p < 0.0001$.

Amplicon 2 contains exon number 2 of *MUTYH* gene. It has been amplified to yield 300 bp PCR products. The PCR products were purified, and DNA cycle sequencing was carried out using the forward primer. However, no variant was found in this exon in the study subjects. Likewise, we carried out a PCR to amplify amplicon number 3, which contains three exons, 3, 4, and 5. This amplicon spans 590 bp as visualized by 2% gel electrophoresis. Analysis of amplicon 3 showed a reported single-nucleotide

polymorphism called c.381A>C in 11 Jordanians with colorectal adenoma. c.381A>C was found to be disease predisposing with Bayes's probability score of 50 (Table 7). Genotype frequency of both mutant (C) and wild type (A) among the study patients were 12 (8%) and 138 (92%), and their allele frequencies were 288 (96%) and 12 (4%), respectively among study patients (Table 6). Furthermore, amplicon 4 in *MUTYH* gene contains exons number 6, 7, 8, and 9. Overall, we identified a reported heterozygous variant, c.724G>A, in seven Jordanians with colorectal adenoma^[20]. c.724G>A was found to be disease causing with a probability score of 61. It was also found probably damaging and deleterious for DNA glycosylase using PolyPhen-2 and PROVEAN pipelines, respectively (Table 7).

Amplicon 5 contains three exons, 10, 11, and 12. The size of PCR products is 690 bp (Fig. 3). Interestingly, the Sanger sequencing revealed a novel variant in 90 (60%) patients with colorectal adenoma from Jordan. As shown in Table 6), c.1264G>C was the highly prevalent variant allele detected in 90 (60%) patients under study. In particular, homozygous allele variant, C/C, was found in 85 (56.7%) patients, while G/C was found in 5 (3.3%) patients. Clearly, a significant association was observed between c.1264G>C variant and colorectal adenoma patients comparable to the controls ($p < 0.0001$). The mutation was mapped to position 1264 of the coding sequence of *MUTYH* (transcript usedNG_008189.1). Besides, c.1264G>C was named according to Human Genome Variation Society guidelines. The mutation was predicted to be disease predisposing variant based on PROVEAN and PolyPhen-2. The normal codon, CAG of glutamine, at position 350 was altered into CAC that codes for histidine. PolyPhen-2 protein prediction analysis showed that the altered amino acid (p.Gln350His) would affect the protein structure and function. C.1264G>C was probably damaging and

Table 6. Genotype and allele frequency of germline variants found in Jordanian patients with colorectal adenoma (n =150)

Amplicon number	Variant identified	Nucleotide (wt:vt)	Genotype frequency			Allele frequency (wt:vt) n (%)
			(wt/wt) n (%)	(wt/vt) n (%)	(vt/vt) n (%)	
1	g.87C>T	C:T	90 (60)	60 (40)	0 (0)	C:240 (80) T:60 (20)
2	-	-	-	-	-	-
3	c. 381A>C	A:C	138 (92)	12 (8)	0 (0)	A:288 (96) C:12 (4)
4	c.724G>A	G:A	143 (95)	7 (5)	0 (0)	G:293 (97.6) A:7 (2.4)
5	c.1264G>C	G:C	60 (40)	5 (3.3)	85 (56.7)	G:125 (41.7) C:175 (58.3)
6	-	-	-	-	-	-
7	-	-	-	-	-	-
8	c.1760C>T	C:T	145 (96.7)	5 (3.3)	0 (0)	C:295 (98.3) T:5 (1.7)

Wt, wild type; vt, variant type




Fig. 2. Mutational analysis of amplicon 1 of *MUTYH* gene. (a) Schematic representation of *MUTYH* gene showing that amplicon 1 contains 5' untranslated region, exon 1, and the intronic boundaries. (b) The reference sequence of amplicon 1 obtained from Gene Bank (NG_008189.1) and the highlighted sequences are orange for TATA box, green for initiator, blue for downstream promoter elements (DPE), and yellow for exon 1. The red box contains the sequence in which g.87C>T variant was identified. The PCR was run using the forward and reverse primers (sequences above and below the block arrows). The expected PCR product was 410 bp. (c) 2% agarose gel electrophoresis of amplicon 1. The amplicon was successfully amplified and 410 bp bands were visualized using ethidium bromide staining. The size was compared to 100 bp ladder (lane 1), and the negative control (lane 2) was included. Lanes from 3-12 and from 13-18 are representative of patients and controls, respectively. (d) Partial chromatogram for forward primer sequencing of amplicon 1 from normal control, showing wild-type sequence (wt/wt). (e) Partial chromatogram for forward primer sequencing from patients illustrating heterozygous, g.87C>T, variant (wt/vt). (f) Partial chromatogram for reverse primer sequencing from normal control indicates wild type (wt/wt). (g) Partial Chromatogram for reverse sequencing from patients, showing heterozygous g.87C>T variant (wt/vt).




Fig. 3. Mutational analysis of the amplicon 5 of *MUTYH* gene. (a) Schematic representation of *MUTYH*, showing that amplicon 5 contains three exons (10, 11, and 12) and the intronic boundaries. (b) The reference sequence of amplicon 5 obtained from Gene Bank (NG_008189.1) and the yellow highlighted sequences are for exons 10, 11, and 12. The PCR was run using the forward and reverse primers (sequences above and below the block arrows). The expected PCR product was 690 bp. The red box shows the position of c.1264G>C mutation. (c) 2% gel agarose gel electrophoresis for amplicon 5. The amplicon was successfully amplified, and 690 bp bands were visualized using ethidium bromide staining. The size was compared to 100 bp ladder (lane 1), and negative control (lane 2) was included. Lanes from 3-12 and 13-20 are representative patients and controls samples, respectively. (d) Partial chromatogram for forward primer sequencing of amplicon 5 from normal control showing wild-type sequence (wt/wt). (e) Partial chromatogram for forward primer sequencing from patients illustrating heterozygous, c.1264G>C, variant (wt/vt). (f) Partial chromatogram for forward primer sequencing from patient indicates homozygous c.1264 G>C mutation (vt/vt). (g) Partial chromatogram for reverse primer sequencing from normal control showing wild-type sequence (wt/wt). (h) Partial chromatogram for reverse primer sequencing from patient with heterozygous, c.1264 C>G, mutation (wt/vt). (i) Partial chromatogram for reverse primer sequence of amplicon 5 from patient with homozygous mutation, c.1264 C>G (vt/vt).

Table 7. Characteristics of germline mutations found in Jordanian patients with colorectal adenoma (n = 150)

Amplicon Number	Variant	dbSNP ID (rs)	MutationTaster Prediction(score)	PolyPhen-2 prediction(score)	PROVEAN prediction(score)
1	g.87C>T	Novel	Polymorphism (12)*	-	-
2	-	-	-	-	-
3	c.381A>C	rs199929178	Disease predisposing (50)*	Benign (0.002)**	-
4	c.724G>A	rs786203212	Disease predisposing (61)*	Probably damaging (1.0)**	Deleterious (-2.863)***
5	c.1264G>C	Novel	Disease predisposing (81)*	Probably damaging (0.58)**	Deleterious (-2.98)***
6	-	-	-	-	-
7	-	-	-	-	-
8	c.1760C>T	rs140118273	Polymorphism (15)*	Benign (0.003)**	Neutral (-1.0)

*Bayes classifier score, which calculates the actual probability that certain alteration causes a disease. The range of result values is between 10 and 100. Therefore, the higher score, the more harmful alteration for human health. **This score indicates the probability that certain mutation is damaging. Hence, variants scores between 0.0 and 0.15 are benign, and variants with scores between 0.15 and 0.40 are possibly damaging. However, variants that score between 0.51 and 1.0 are probably damaging. ***Delta alignment score of PROVEAN: if the score is below or equal to the threshold (-2.5), the alteration is predicted to have deleterious effect. However, if the variant has a score above -2.5, the variant is predicted to be neutral.

deleterious to DNA glycosylase based on data obtained from PolyPhen2 and PROVEAN, which scored 0.58 and -2.98, respectively (Table 7). The high prevalence of c.1264G>C in Jordanian patients with colorectal adenoma stressed the importance of investigating the association between this mutation and the progression of adenoma into carcinoma (Fig. 4). Interestingly, a significant association was observed between c.1264G>C mutation and malignant transformation of colorectal adenoma among patients ($p < 0.001$). This observation brings the necessity to evaluate the relationship between c.1264G>C and colorectal cancer metastasis. Hence, the disease pattern was followed, and the c.1264G>C was found to be significantly associated with CRC metastasis (Fig. 5).

Exon 15 and its intronic boundaries (amplicon 7) were also amplified using the primers listed as 7 in the Table 1. PCR was successfully performed and resulted in amplification of 230-bp products. The products were purified and sequenced using 3130xL Genetic Analyzer (Applied Biosystem, USA). However, no mutation was found in this exon among the study subjects.

Finally, exon 16 and its intronic boundaries (amplicon 8) were amplified using the listed primer 8 (Table 1). PCR successfully amplified 410-bp products. The products were purified and sequenced using 3130XL Genetic Analyzer. A reported variant, rs140118273, was successfully identified in Jordanians with colorectal adenoma. However, the variant was found to be neutral.

DISCUSSION

In this study, five variants were found in Jordanian patients with colorectal adenoma, of which three

(c.381A>C, c.724G>C, and c.1760 C>T) has previously been reported, and two (g.87C>T and c.1264G>C [Q350H]) were novel. These findings are in contrast to Caucasian populations in which Y165C and G382D are the most common mutations associated with multiple polyposis syndrome^[18]. The Y165C and G382D mutations have been shown to impair the MUTYH repair activity and significantly contribute to CRC development^[19]. The missense R245C and the splice site IVS10-2A>G variants have been identified in Japanese and Korean patients^[15,20]. In another studies conducted in Korean and Japanese patients (n = 97), 7.2% were bi-allelic carriers for the germline mutations c.1-18G>T, A359V, and R170G in the *MUTYH* gene^[21,22]. After screening the entire gene, Vandrovcová *et al.*^[23] found no mutation in *MUTYH*




Fig. 4. Association between c.1264G>C and colorectal adenoma malignant transformation. A strong association was found among the study subjects ($p < 0.001$). Of 90 subjects, who harbored the c.1264G>C germline mutation, 82 (91.2%) had progressed to colorectal cancer. In contrast, only 25 wild-type patients had progressed into CRC, comparable to 35 subjects who recovered after polypectomy.




Fig. 5. The association between c.1264G>C and CRC metastasis. A significant ($p < 0.001$) association was observed between c.1264G>C and CRC metastasis. Sixty four subjects included in this study with c.1264G>C germline mutation had CRC metastasis, comparable to 12 who had recovered after colectomy and chemotherapy.

in Singaporean population, though this may be a biased result due to the small sample size ($n = 63$). Furthermore, five unrelated Indians patients with colorectal adenoma were homozygous for the missense mutation E480X^[23]. Other *MUTYH* variants have been found in Italian (A473D), Finnish (p.P391L), Swedish (p.G175E), and Portuguese (E383fsX45)1^[24-27]. In a case-control study and meta-analysis, Win *et al.*^[26] have indicated an association between *MUTYH* gene mutations and the increased CRC risk. Another meta-analysis demonstrated the risk of *MUTYH* in monallelic and biallelic carriers^[28]. *MUTYH* gene loss has been linked to colorectal carcinogenesis due to immunosuppression and altered immune response^[27]. Grasso *et al.*^[28] have shown that *MUTYH* mediates the toxicity of DNA 6-thioguanine and UV radiation. *MUTYH*-OGG1 XRCC1-PARP1-MMP1 is a linear interacting susceptibility locus for CRC^[29].

In Arabian descent populations, four studies have been performed; one in Tunisia that showed high prevalence of c.1227-1228dup^[30], the second in Morocco that introduced three variants (c.494A>G, c.1145G>A, and c.1185_1186dup)^[31], the third in Saudi Arabia that revealed the presence of V22M, Y165C, H324Q, and G382D variants^[32], and the last in Egypt that found a significant association between G396D and Y179C mutations and colorectal carcinogenicity^[33]. These results contradict with those obtained from a study involved 360 Arabic patients^[34]. The best explanation for these discrepancies between data is that these studies only targeted either specific reported mutations, such as G396D and Y179C, or

familial cases. However, in the present study, *MUTYH* gene was sequenced thoroughly for the patients with colorectal adenoma and the controls from the same ethnic group. It has recently been shown that *MUTYH* interacts with *Rad9-Rad1-Hus1* complex (9-1-1 complex) to coordinate the cell cycle checkpoint. The interdomain connector (IDC) of *MUTYH* (residues between 65 and 350) is critical for 9-1-1 complex and *MUTYH* interaction^[11]. The significance of the interaction between *SpMyhII* DC and 9-1-1 has been elucidated by Luncsford *et al.*^[11] who tested *in vivo* the biological effects of V315 and I261 mutations in the impairment of 9-1-1 complex-*MUTYH* interaction. Interestingly, they showed that *MUTYH* IDC is important for *MUTYH* DNA repair function and interaction with 9-1-1 complex. *MUTYH* IDC aspartate at the position number 350 interacts with histidine at the position number 18 of 9-1-1 complex^[35]. Therefore, c.1264G>C missense variant found in Jordanian patients would probably affect *MUTYH*-9-1-1 complex interaction. This finding explains the strong association between c.1264 G>C and colorectal malignant transformation among mutant colorectal adenoma patients. However, this hypothesis requires further functional testing using cell-based cloning and cell culture manipulation.

In conclusion, germline mutations in *MUTYH* gene are highly prevalent among patients with colorectal adenoma from Jordan. In particular, c.1264 G>C is the most common variant associated with malignant transformation of colorectal adenoma into carcinoma. This variant has a potential to be an early biomarker for diagnosis of CRC in Jordan.

ACKNOWLEDGEMENTS

This work was supported by the Deanship of Research at Jordan University of Science and Technology, Jordan (Grant number 22/2016).

CONFLICT OF INTEREST. None declared.

REFERENCES

1. Dulai PS, Sandborn WJ, Gupta S. Colorectal cancer and dysplasia in inflammatory bowel disease: a review of disease epidemiology, pathophysiology, and management. *Cancer prevention research* 2016; **9**(12): 887-894.
2. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002; **420**(6917): 860-867.
3. Levine B, Kroemer G. Autophagy in the pathogenesis of disease. *Cell* 2008; **132**(1): 27-42.
4. Spira A, Yurgelun MB, Alexandrov L, Rao A, Bejar R,

- Polyak K, Giannakis M, Shilatifard A, Finn OJ, Dhodapkar M, Kay NE, Braggio E, Vilar E, Mazzilli SA, Rebbeck TR, Garber JE, Velculescu VE, Disis ML, Wallace DC, Lippman SM. Precancer atlas to drive precision prevention trials. *Cancer research* 2017; **77**(7): 1510-1541.
5. Rubin R, Strayer DS, Rubin E, McDonald JM. Rubin's Pathology: Clinicopathologic Foundations of Medicine (5th ed.). Philadelphia: Lippincott Williams & Wilkins; 2008.
6. Heitman SJ, Ronksley PE, Hilsden RJ, Manns BJ, Rostom A, Hemmelgarn BR. Prevalence of adenomas and colorectal cancer in average risk individuals: a systematic review and meta-analysis. *Clinical gastroenterology and hepatology* 2009; **7**(12): 1272-1278.
7. Kycler W, Korski K, Łoziński C, Teresiak-Mańczak A, Przybyła A, Mackiewicz A, Cybulski Z, Lamperska K. The anti-cancer actions of O6-methylguanine-DNA-methyltransferase in relation to colon polyps. *Pharmacological reports* 2014; **66**(6): 1060-1064.
8. Powell SM, Zilz N, Beazer-Barclay Y, Brayn TM, Hamilton SR, Thibodeau SN, Vogelstein B, Kinzler KW. APC mutations occur early during colorectal tumorigenesis. *Nature* 1992; **359**(6392): 235-237.
9. Redrejo-Rodríguez M, Vigouroux A, Mursalimov A, Grin I, Alili D, Koshenov Z, Akishev Z, Maksimenko A, Bissenbaev AK, Matkarimov BT, Saparbaev M, Ishchenko AA, Moréra S. Structural comparison of AP endonucleases from the exonuclease III family reveals new amino acid residues in human AP endonuclease I that are involved in incision of damaged DNA. *Biochimie* 2016; **129**: 20-33.
10. Out AA, Tops CM, Nielsen M, Weiss MM, van Minderhout IJ, Fokkema IF, Buisine MP, Claes K, Colas C, Fodde R, Fostira F, Franken PF, Gaustadnes M, Heinimann K, Hodgson SV, Hogervorst FB, Holinski-Feder E, Lagerstedt-Robinson K, Olschwang S, van den Ouweland AM, Redeker EJ, Scott RJ, Vankeirsbilck B, Grønlund RV, Wijnen JT, Wikman FP, Aretz S, Sampson JR, Devilee P, den Dunnen JT, Hes FJ. Leiden open variation database of the MUTYH gene. *Human mutation* 2010; **31**(11): 1205-1215.
11. Luncsford PJ, Chang DY, Shi G, Bernstein J, Madabushi A, Patterson DN, Lu AL, Toth EA. A structural hinge in eukaryotic MutY homologues mediates catalytic activity and Rad9-Rad1-Hus1 checkpoint complex interactions. *Journal of molecular biology* 2010; **403**(3): 351-370.
12. Oka S, Nakabeppu Y. DNA glycosylase encoded by MUTYH functions as a molecular switch for programmed cell death under oxidative stress to suppress tumorigenesis. *Cancer science* 2011; **102**(4): 677-682.
13. Waz S, Nakamura T, Hirata K, Koga-Ogawa Y, Chirifu M, Arimori T, Tamada T, Ikemizu S, Nakabeppu Y, Yamagata Y. Structural and kinetics of the human nudix hydrolase MTH1 reveal the mechanism for its broad substrate specificity. *Journal of biological chemistry* 2017; **292**(7): 2785-2794.
14. Nakabeppu Y, Sakumi K, Sakamoto K, Tsuchimoto D, Tsuzuki T, Nakatsu Y. Mutagenesis and carcinogenesis caused by the oxidation of nucleic acids. *Biological chemistry* 2006; **387**(4): 373-379.
15. Miyaki M, Iijima T, Yamaguchi T, Hishima T, Tamura K, Utsunomiya J, Mori T. Germline mutations of the MYH gene in Japanese patients with multiple colorectal adenomas. *Mutation research* 2005; **578**(1-2): 430-433.
16. Mahasneh A, Al-Shaheri F, Jamal E. Molecular biomarkers for an early diagnosis, effective treatment and prognosis of colorectal cancer: Current updates. *Experimental and molecular pathology* 2017; **102**(3): 475-483.
17. Luo HY, Xu RH. Predictive and prognostic biomarkers with therapeutic targets in advanced colorectal cancer. *World journal of gastroenterology* 2014; **20**(14): 3858-3874.
18. Al-Tassan N, Chmiel NH, Maynard J, Fleming N, Livingston AL, Williams GT, Hodges AK, Davies DR, David SS, Sampson JR, Cheadle JP. Inherited variants of MYH associated with somatic G:C->T:A mutations in colorectal tumors. *Nature genetics* 2002; **30**(2): 227-232.
19. Parker AR, Sieber OM, Shi C, Hua L, Takao M, Tomlinson IP, Eshleman JR. Cells with pathogenic biallelic mutations in the human MUTYH gene are defective in DNA damage binding and repair. *Carcinogenesis* 2005; **26**(11): 2010-2018.
20. Lynch, H.T., Tinley, S.T., Lynch, J.F., Attard, T.M., 2004. Challenging pedigrees seen in a hereditary cancer consultation center. *Cancer genetics and cytogenetics* 2004; **153**(2): 91-101.
21. Shinmura K, Goto M, Tao H, Kato H, Suzuki R, Nakamura S, et al. Impaired 8-hydroxyguanine repair activity of MUTYH variant p.Arg109Trp found in a Japanese patient with early-onset colorectal cancer. *Oxidative medicine and cellular longevity* 2014; **2014**: e617351.
22. Kairupan C, Scott RJ. Base excision repair and the role of MUTYH. *Hereditary cancer clinical practice* 2007; **5**(4): 199-209.
23. Vandrovová J, Štekrová J, Kebrdlová V, Kohoutová M. Molecular analysis of the APC and MYH genes in Czech families affected by FAP or multiple adenomas: 13 novel mutations. *Human mutation* 2004; **23**(4): 397.
24. Russell AM, Zhang J, Luz J, Hutter P, Chappuis PO, Berthod CR, Maillet P, Mueller H, Heinimann K. Prevalence of MYH germline mutations in Swiss APC mutation-negative polyposis patients. *International journal of cancer* 2006; **118**(8):1937-1940.
25. Gismondi V, Meta M, Bonelli L, Radice P, Sala P, Bertario L, Viel A, Fornasarig M, Arrigoni A, Gentile M, Ponz de Leon M, Anselmi L, Mareni C, Bruzzi P, Varesco L. Prevalence of the Y165C, G382D and 1395delGGA germline mutations of the MYH gene in Italian patients with adenomatous polyposis coli and colorectal adenomas. *International journal of cancer* 2004; **109**(5): 680-684.
26. Win AK, Hopper JL, Jenkins MA. Association between monoallelic MUTYH mutation and colorectal cancer

- risk: a meta-regression analysis. *Familial cancer* 2011; **10**(1): 1-9.
27. Tenesa A, Campbell H, Barnetson R, Porteous M, Dunlop M, Farrington SM. Association of MUTYH and colorectal cancer. *British journal of cancer* 2006; **95**(2): 239-242.
 28. Grasso F, Ruggieri V, De Luca G, Leopardi, P, Mancuso MT, Casorelli I, Pichierri P, Karran P, Bignami M. MUTYH mediates the toxicity of combined DNA 6-thioguanine and UVA radiation. *Oncotarget* 2015; **6**(10): 7481-7492.
 29. Grasso F, Di Meo S, De Luca G, Pasquini L, Rossi S, Boirivant M, Biffoni M, Bignami M, Di Carlo E. The MUTYH base excision repair gene protects against inflammation-associated colorectal carcinogenesis. *Oncotarget* 2015, **6**(23): 19671-19684.
 30. Abdelmaksoud-Dammak R, Miladi-Abdennadher I, Amouri A, Tahri N, Ayadi L, Khabir A, Frikha F, Gargouri A, Mokdad-Gargouri R. High prevalence of the c.1227_1228dup (p.Glu410GlyfsX43) mutation in Tunisian families affected with MUTYH-associated-polypsis. *Familial cancer* 2012; **11**(3): 503-508.
 31. Laarabi FZ, Cherkaoui Jaouad I, Baert-Desurmont S, Ouldim K, Ibrahimi A, Kanouni N, Frebourg T, Sefiani A. The first mutations in the MYH gene reported in Moroccan colon cancer patients. *Gene* 2012; **496**(1): 55-58.
 32. Adi A, Bin Abbas B, Al Hamed M, Al Tassan N, Bakheet D. Screening for mutations in ABCC8 and KCNJ11 genes in Saudi persistent hyperinsulinemic hypoglycemia of infancy (PHHI) patients. *Genes (Basel)* 2015; **6**(2): 206-215.
 33. Elsaid A, Elshazli R, El-Tarapely F, Darwish H, Abdel-Malak C. Association of monoallelic MUTYH mutation among Egyptian patients with colorectal cancer. *Familial cancer* 2017; **16**(1): 83-90.
 34. Lejbkowitz F, Cohen I, Briness OB, Pinchev M, Poynter J, Gruber SB, Stephen SB, Rennert G. Common MUTYH mutations and colorectal cancer risk in multiethnic populations. *Familial cancer* 2012; **11**(3): 329-335.
 35. Jin J, Hwang BJ, Chang PW, Toth EA, Lu AL. Interaction of apurinic/apyrimidinic endonuclease 2 (Ape2) with Myh1 DNA glycosylase in fission yeast. *DNA repair (Amst)* 2014; **15**: 1-10.