OPEN ACCESS

Gαq Signaling Activates β-Catenin-Dependent Gene Transcription

Sara Ansari, Sedighe Kolivand, Sara Salmanian, Marie Saghaeian Jazi, S Mahmoud A Najafi*

Department of Cell and Molecular Biology, School of Biology, College of Sciences, University of Tehran, P.O.Box 14155-6455, Tehran, Iran

ABSTRACT

Background: The canonical Wnt signal transduction or the Wnt/ β -catenin Received: 21 January 2023 Accepted: 28 May 2023 pathway plays a crucial role in both carcinogenesis and development of Published online: 31 May 2023 animals. Activation of the Gaq class of Ga proteins positively regulates Wnt/ β -catenin pathway, and expression of G α q in HEK293T cells or Xenopus oocytes leads to the inhibition of GSK-3 β and cellular accumulation of β catenin. This study investigated whether Gaq-mediated cellular accumulation of β -catenin could affect the transcriptional activity of this protein. Methods: HEK-293T and HT-29 cells were used for cell culture and transfection. Protein localization and quantification were assessed by using immunofluorescence microscopy, cell fractionation assay, and Western blotting analysis. Gene expression at the transcription level was examined by quantitative reverse transcriptase/real-time PCR method. **Results**: Transcription of two cellular β -catenin target genes (*c*-MYC and CCND1) and the β -catenin/TCF reporter *luciferase* gene (TopFlash plasmid) significantly increased by Gaq activation. The Gaq-mediated increase in the expression level of the β -catenin-target genes was sensitive to the expression of a minigene encoding a specific Gaq blocking peptide. The results of cell fractionation and Western blotting experiments showed that activation of Citation: Gaq signaling increased the intracellular β -catenin protein level, but it Ansari S, Kolivand S, Salmanian S, Saghaeian Jazi M, Najafi SMA. Goq blocked its membrane localization. Signaling Activates β-Catenin-

Conclusion: Our results reveal that the G α q-dependent cellular accumulation of β -catenin can enhance β -catenin transcriptional activity. **DOI**: 10.61186/ibj.3890

Keywords: β-catenin, Wnt signaling, G proteins

Corresponding Authors: S Mahmoud A Najafi

Dependent Gene Transcription.

Iranian biomedical journal 2023;

27(4): 183-190.

Department of Cell and Molecular Biology, School of Biology, College of Sciences, University of Tehran, P.O.Box 14155-6455, Tehran, Iran; Tel.: (+98-21) 61113648; Fax: (+98-71) 66492992; E-mail: smarabnajafi@ut.ac.ir

INTRODUCTION

S pecificity in signal transduction pathway is mediated through many known and unknown mechanisms. One of these mechanisms is the cross-talk between the components of two or more signaling pathways. Both the heterotrimeric G protein and canonical Wnt/ β -catenin signaling pathways play essential roles in various cellular processes and deregulation of these pathways leads to abnormalities in

List of Abbreviations:

APC: adenomatous polyposis coli; AXIN: axis inhibitor; DTT: dithiothreitol; EDTA: ethylenediaminetetraacetic acid; GPCR: G-proteincoupled receptors; GSK-3β: glycogen synthase kinase-3 beta; HEK293: human embryonic kidney 293; IP5: inositol pentakisphosphate; LEF: lymphoid enhancer factor; LRP5: LDL Receptor Related Protein 5; PAR1: protease activated receptor 1; PBS: phosphate-buffered saline; PMSF: phenylmethylsulfonyl fluoride; RT: room temperature; RT-PCR: reverse transcriptase-PCR; TBS: tris-buffered saline; TCF: T-cell factor Gαq Upregulates β-Catenin Nuclear Function

animal development, as well as many human diseases, including cancers^[1-3].

Activation of the canonical Wnt signaling pathway initiates by interacting the Wnt glycoproteins with the Frizzle/LRP5 receptors. While the mechanism of this pathway is still not well understood, its activation results in the inhibition of GSK activity and stabilization of cellular β -catenin protein^[1,2]. Translocation of β -catenin into the nucleus and its interaction with the TCF/LEF transcription factors leads to the transcriptional regulation of the genes involving in diverse cellular processes. In addition, β -catenin has an important role in maintaining epithelial tissues by interacting with the cadherin cell-cell adhesion proteins^[4-5].

Given the structural similarity between GPCRs and Frizzled receptors in terms of possessing a heptahelical transmembrane domain^[5], the possibility of regulating Wnt signaling by heterotrimeric G-proteins has been investigated by several researchers who demonstrated that heterotrimeric G-proteins regulate both canonical and non-canonical Wnt signaling pathways^[6-11]. Formerly, we have also shown that activation of the Gaq class of G proteins in two different cell systems (*Xenopus* oocytes and HEK293T cells) leads to the inhibition of GSK activity and accumulation of β -catenin in the cell membrane^[12,13].

In this study, we have examined whether the Gaqmediated increase in cellular β -catenin protein level affects the β -catenin-dependent gene transcription. Consistent with our previous results, we herein showed that Gaq signaling induces the transcription of the β catenin target genes. Moreover, we provided preliminary results supporting that Gaq signaling increases the intracellular β -catenin level, at least partially, by inhibiting the membrane localization of this protein.

MATERIALS AND METHODS

Cell culture and transfection

HEK293T and HT-29 cells were respectively grown in DMEM and RPMI 1640 medium supplemented with 10% FBS and antibiotics (100 μ g/ml of streptomycin and 100 U/ml of penicillin) in 5% CO₂ at 37 °C. HEK293T cells are responsive to the canonical Wnt signaling and have therefore been used extensively by the researchers in the field of Wnt signaling pathways. At 60% confluency, the medium of HEK-293T cells was replaced with the fresh medium, and then two hours later, the cells were transfected with the expression plasmids. The standard calcium phosphate protocol was used for transfection^[13]. Six hours post transfection, the medium was changed, and 48 hours later, the cells were harvested. The cell pellets were used directly or stored at -70 °C until use. Treatment of cells with carbachol (100 μ M) or thrombin (0.5 U/ml) was performed 3 and 10 hours before the cell harvest, respectively.

Cell fractionation and Western blotting analysis

Isolation of cytoplasmic and membrane proteins from HEK293T cells, as well as measurement of protein concentration, were performed as described before^[13]. Briefly, the cells were suspended in a buffer containing 50 mM of Tris-HCl (pH 7.8), 100 mM of NaCl, 2 mM of EDTA, 1 mM of DTT, 0.5 mM of PMSF, 1.5 µM of pepstatin A and lysed by passing them 25 times through a 27-gauge needle. The lysates were centrifuged at maximum speed in a refrigerated microfuge $(15.339 \times g)$ for 15 min, and the supernatants containing the cytoplasmic proteins were transferred to a new tube and stored at -70 °C until use. The pellet containing membrane proteins was rinsed using the abovementioned buffer and centrifuged, accordingly. The washed pellet was dissolved in 120 µl of 50 mM of Tris-HCl (pH 7.8), 100 mM of NaCl, 1 mM of EDTA, 1 mM of DTT, 2% SDS, 100 mM of β-mecaptoethanol, 10% glycerol, 0.5 mM of PMSF, and 1.5 µM of pepstatin A. The suspension was heated in a 95 °C water bath for 4 min, vortexed and heated again at 95 °C for 3 min. The suspension was centrifuged (15.339 \times g) at RT for 30 min, and the supernatant was preserved as membrane proteins. Protein concentration was measured using a previously described method^[14]. To this end, 20 µg of proteins from each fraction was separated by SDS-PAGE (8%) and transferred to a nitrocellulose membrane. Following electrical transfer, the gel was stained with Coomassie blue overnight to verify transfer efficiency. The membrane was washed using TBS (25 mM of Tris-HCl [pH 7.5], 137 mM of NaCl, and 2.7 mM of KCl) for 15 min and incubated in fat-free milk 5% (in TBS) at RT for 1 hour to block the nonspecific binding. The blot was then incubated with a polyclonal primary antibody against β -catenin at a 1:1000 dilution at RT for 1 hour. The blot was again washed four times (1, 5, 10, and 15 min, respectively) with TBS and incubated for another one hour with horseradish peroxidase-labelled goat anti-rabbit IgG (1:1000 dilution) at RT. The membrane was finally washed in TBS for 15 min, and the antigen-antibody complex was visualized using an ECL detection kit (Pierce, USA).

Indirect immunofluorescence

Indirect immunofluorescence experiments were performed as described previously^[13]. In summary, HEK293T cells grown on glass coverslips were washed in PBS, fixed in 4% paraformaldehyde and permeabilized in 0.2% Triton X-100 in PBS for three

minutes. The coverslips were washed with PBS and then incubated in blocking buffer (3% BSA) for 30 min before being incubated in a 1:500 dilution of β -catenin antibody at RT for 1 h. The coverslips were again rinsed three times (10 min each) in PBS and then incubated with FITC-conjugated secondary antibody (1:250) at RT for 1 h. The coverslips were finally washed and visualized by a fluorescence microscope (Zeiss, Germany).

Quantitative reverse transcriptase and real-time PCR experiments

HEK293T cells were harvested 48 hours post transfection with Trypsin/EDTA (0.53 mM of EDTA and 0.05% [w/v] Trypsin in PBS) and washed twice with cold PBS. Total RNA was extracted from the cell pellets using the RNX-Plus kit (CinnaGen, Tehran, Iran) as described by the supplier. Next, 2 µg of RNA was treated with 1 U of DNase I in a total volume of 10 µl at 37 °C for 30 min. The DNase enzyme was inactivated in 2.5 mM of EDTA at 65 °C for 10 min, and then the reaction was used for reverse transcription by adding 200 U of reverse transcriptase enzyme (Fermentase, USA), 1× RT buffer, 20 U of RiboLock RNase Inhibitor, 1 mM of dNTP mix, and 0.2 µg of random hexamer primer in a total volume of 25 µl. PCR amplification was performed on 1 µl of the reverse transcription reaction using 30 pmol of each primer and 2.5 U of Taq polymerase in a total volume of 25 μ l. The amplification protocol included denaturating at 95 °C for 60 s, annealing (GAPDH and c-MYC at 59 °C, CCND1 at 57 °C, and luciferase at 62 °C) for 60 s, and extension at 72 °C for 60 s. The 30 cycles of PCR were followed by a final extension at 72 °C for 10 min. PCR products were separated on a 1% agarose gel and visualized by ethidium bromide. The results of RT-PCR experiments were quantified by the Image J software. Real time-PCR experiments were performed using the QuantiFast[™] SYBR Green PCR Master Mix kit (Qiagen, Germany) under the following programs: initial denaturation at 95 °C for 5 min, followed by 40 cycles of denaturation at 95 °C for 15 s, annealing at 60 °C for 25 s, and extension at 72 °C for 25 s. For each 10 µl reaction, 1 µl of complementary DNA (100 ng) was used with 1 µl of each primer (1.0 µM) plus 5 µl of SYBR Green I Master Mix on a real-time thermo cycler Rotor Gene 6000 (Qiagen). Analysis was performed using Corbett rotor-gene 6000 software based on the comparative Ct method. The relative amount of the target gene expression was quantified relative to the reference gene (GAPDH) expression. The primers are listed in Table 1.

Statistical analysis

All the experiments were repeated at least three times. The data were expressed as means \pm SE. Statistical data analysis was performed using SPSS 16.0 and Prism 8 software. Unpaired student t-test and one-way ANOVA were used for comparing the groups. To compare differences in gene expression between the groups, nonparametric Kruskal-Wallis H and Mann-Whitney U tests were applied. p values ≤ 0.05 was considered statistically significant.

Primers Gene F: 5' CCA GGT GGT CTC CTC TGA CTT CAA CAG 3' R: 5' AGG GTC TCT CTC TTC TTC CTC TTG TGC TGC 3' GAPDH F: 5' AAG GTG AAG GTC GGA GTC AAC 3' R: 5' GGG GTC ATT GAT GGC AAC AAT A 3' F: 5' TTC CTC TCC AAAATG CCA G 3' R: 5' AGA GAT GGA AGG GGG AAA GA 3' CCND1 F: 5' GAG GGT TGT GCT ACA GAT GA 3' R: 5' CGC CTC CTT TGT GTT AAT GC 3' F: 5' CAC CAA CAG GAA CTA TGA CC 3' R: 5' CGC AGA TGA AAC TCT GGT TC 3' c-MYC F: 5' GGC TCC TGG CAA AAG GTC A 3' R: 5' AGT TGT GCT GAT GTG TGG AGA 3

Table 1. Oligonucleotide primers used for RT-PCR and real-time PCR experiments

F, forward primer; R, reverse primer. When two pairs of primers were designed for gene amplification, the underlined primers were used specifically for real-time PCR experiments.

F: 5' CTC ATA GAA CTG CCT GCG TG 3'

R: 5' GGC GAA GAA GGA GAA TAG GG 3'

luciferase

DOI: 10.61186/ibj.3890]

RESULTS

Activation of Gαq signaling increases the expression of β-catenin target genes

To evaluate β -catenin transcriptional activities, we selected two native cellular genes, CCND1 (the cyclin D1-encoding gene) and c-MYC plus the reporter *luciferase* gene cloned into the TopFlash plasmid^[15]. Although it has been shown that the mentioned genes are responsive to the β -catenin/TCF complex^[15-18], we measured their transcription in the presence and absence of exogenous β -catenin to further verify that the expression of the chosen genes can be induced by β catenin (Fig. 1A). We have previously shown that the expression of Gaq or activation of the endogenous Gaq increases intracellular β -catenin protein level^[12,13]. Therefore, we initially used quantitative reverse transcriptase PCR to test whether Gaq expression affects the transcription of the β -catenin target genes or not. As shown in Figure 1B, the expression of $G\alpha q$ increased the transcription of the target genes (luciferase [p = 0.014], *c-myc* [p = 0.037], and *CCND1* [p = 0.1]) by about two-fold. This increase was blocked by the expression of a Gaq minigene, which encodes a short peptide corresponding to the C-terminal 11 amino acids of Gaq. The specificity of this peptide in blocking Gaq has previously been verified^[13,19,20]. We then examined if endogenous Gαq activation could induce β-catenin target gene expression. In this regard, we treated HEK293T cells with thrombin or carbachol. Thrombin is a known agonist for PAR1, a GPCR that preferentially couples to Gq^[21]. Carbachol is a cholinergic agonist that activates M1/3 muscarinic acetyl choline receptors, the GPCRs that also couple to Gq^[22]. We have previously shown that treatment of HEK293T cells with these two agonists leads to more than two-fold increase in the cellular protein level of β-catenin^[13]. Treatment of HEK293T cells with these agonists (0.5 U/ml thrombin or 100 µM carbachol) did not significantly change the expression of the β -catenin target genes (Fig. 2A). Expression of the reporter luciferase gene was also not affected by the treatment of these cells with carbachol (data not shown). Although there are reports that HEK293T cells endogenously express both PAR1 and muscarinic receptors^[23,24], the expression levels may not induce Gaq-mediated β-catenin be sufficient to transcriptional activity. To verify this assumption, we used HT-29 colon cancer cells, which express higher levels of several Gq-coupled GPCRs including PAR1, as compared to the corresponding normal cells^[25]. As



Fig. 1. (A) HEK293T cells were transfected with empty vector (V) or the β -catenin-encoding plasmid and used for RT-PCR experiments to amplify the *c-MYC*, *CCND1*, *luciferase*, and *GAPDH* genes. The Topflash plasmid, carrying several β -catenin/TCF target elements, was employed to measure the expression of the reporter *luciferase* gene^[14]. The top panel (W) is the result of a Western blot experiment, showing the expression of exogenous β -catenin; (B) Expression of two native β -catenin target genes (*CCND1* and *c-MYC*) and the β -catenin-responsive *luciferase* gene in HEK293T cells transfected with the nonrecombinant (v), *Gaq*, and *mq* plasmids. Fourty eight hours post transfection, RT-PCR assay was performed. *Gaq* and *mq* represent the plasmids encoding Gaq and the Gaq blocking peptide, respectively. The chart shows the average of three independent experiments (* p < 0.05 and † p < 0.1). The result of one of the RT-PCR gels is shown above the chart. V; nonrecombinant vector



Fig. 2. HEK293T (A) or HT-29 (B) cells were treated with 0.5 U/ml of thrombin (10 h) or 100 μ M of carbachol (3 h) and used for real time-PCR experiments to measure the expression of the *c-MYC* and *CCND1* genes. The charts represent an average of three independent experiments (****p < 0.001 for thrombin vs. non-treated cells [NT]).

shown in Figure 2B, while treatment of HT-29 cells with carbachol slightly increased *CCND1* and *c-MYC* gene transcription, the thrombin treatment led to about 2- to 2.5-fold increase in the expression of these two genes (Fig. 2B).

Activation of Gαq signaling appears to block β-catenin cell membrane localization

As shown in Figure 3, in HEK293T cells, β-catenin is mainly present in the cell membrane and the expression of Gaq in these cells has no effect on the membranelocalized β -catenin. However, it was interesting to examine the membrane localization of β -catenin when this protein was co-expressed with Gaq. Therefore, we took advantage of an available β -catenin expression construct producing a Myc-tagged version of the protein. The tagged β -catenin protein had a molecular weight of about 110 kDa, 22 kDa larger than that of the wild type native β -catenin (88 kDa) and, therefore, the two forms of the protein could be observed on the blot. The results of cell fractionation and Western blotting experiments showed that when the Myc-tagged β catenin was expressed in the cells, most of the proteins were localized in the cytoplasm. The protein was also well-localized to the plasma membrane, showing that the 22 kDa Myc tag peptide does not inhibit the membrane localization of β -catenin (Fig. 4A). When Gaq was co-expressed with the Myc-tagged β -catenin, although the total amount of the tagged β -catenin increased in the cell (Fig. 4B), the membrane level of this exogenous β -catenin clearly decreased (Fig. 4C).

DISCUSSION

We have previously reported that activation of the Gaq class of Ga proteins inhibits GSK activity, which in turn increases the intracellular level of β -catenin protein^[13,14]. However, it has not before been examined

whether the Gaq-mediated increase in the β -catenin protein level affects the β -catenin-mediated gene transcription. Therefore, in this study, we attempted to find the answer to this question. For gene expression experiments, two native cellular β -catenin target genes (*c-MYC* and *CCND1*) plus the reporter *luciferase* gene (under control of the TCF/LEF-binding elements, pTopflash) were selected^[15]. As we intended to assess β -catenin transcriptional activity, RT-PCR (quantitative and real-time PCR) experiments were used to measure the *luciferase* gene transcription. Altogether, our results clearly showed that the Gaq signaling activation induced β -catenin transcriptional activity.

In this study, we also found that the G α q activation could inhibit β -catenin membrane localization (Fig. 4). This is a very interesting preliminary observation, which definitely needs further investigation. Upregulation of



Fig. 3. HEK-239T cells transfected with empty vector or $G\alpha q$ encoding plasmid and used for immunofluorescence microscopy experiment using β -catenin antibody.

DOI: 10.61186/ibj.3890]



Fig. 4. HEK293T cells transfected with a Myc-tagged β -catenin-encoding plasmid (A) with or without the Gaq-encoding plasmid (B). Fourty eight hours post transfection, the cells were collected, fractionated and used for Western blotting experiments. (C) HEK293T cells were transfected with β -catenin-encoding plasmid alone or together with the Gaq-encoding plasmid. Total cellular proteins were extracted 48 h post transfection and used for Western blotting analysis. Exo, exogenous; endo, endogenous

 β -catenin through Gaq signaling has also been reported by other researchers. In this regard, it has been shown that Frizzled1-mediated differentiation of F9 teratocarcinoma cells (a process that requires β catenin/TCF-LEF pathway) is dependent on Gaq signaling^[7]. Furthermore, it has been reported that treatment of F9 teratocarcinoma cells with Wnt-3a Gaq-mediated phosphatidylinositol activates а signaling, which results in generating inositol polyphosphates such as IP5^[10]. Evidence has suggested that transient accumulation of IP5 inhibits GSK-3β and, therefore, leads to the accumulation of β -catenin and TCF/LEF-dependent transcription^[10]. Moreover, Gag signaling is required for the Wnt-mediated disruption of Axin/GSK-3 β interaction, which may result in β -catenin cellular stabilization^[8]. β-catenin is a known protooncoprotein, which its upregulation is involved in tumorigenesis of several human cancers including colorectal cancer. It is believed that this protein is a highly potential target for cancer prevention and therapy^[26-28]. In addition to the Gq class of heterotrimeric G-proteins, the involvement of other classes of G-proteins in the regulation of β -catenin expression and function has been reported^[29]. GPCRs are the most diverse class of proteins in mammals involved in many critical cellular processes. Around 900 different GPCRs are encoded by human genome^[3,30],

which are the target of more than 30% of the approved therapeutic drugs^[30,31]. Regulation of β -catenin by heterotrimeric G-proteins raises an interesting question: would it be possible to target G-protein signaling pathways in order to control the expression and function of β -catenin in human cancers? The answer to this question definitely needs much further investigation. Fortunately, the data on the deregulation of G protein signaling pathways in human cancers is growing, and several excellent review articles have already been published, supporting that these important classes of proteins can be considered as potential targets for cancer therapeutics^[3,29-33].

In conclusion, the activation of Gaq signaling not only increases the intracellular β -catenin protein level but also results in the regulation of β -catenin nuclear activity. The present study clearly shows that the expression of the wild-type Gaq or activation of the endogenous Gaq causes transcriptional upregulation of two known β -catenin cellular target genes (*c-MYC* and *CCND1*) and a β -catenin-responsive reporter gene. Since β -catenin deregulation is associated with many human cancers, our results further support this hypothesis that trimeric G-proteins and their receptors are among the potential targets for cancer therapy.

188

DECLARATIONS

Ethical statement

Not applicable.

Data availability

Data supporting this article are included within the article and supplementary file.

Author contributions

SA, SK, SS, and MSJ: contributed to experimental design and also performed experiments; SMAN: contributed to experimental design, supervised the project, and wrote the manuscript.

Conflict of interest

None declared.

Funding/support

This work was supported in part by a grant from Iran National Science Foundation, INSF (93004354) to S Mahmoud A Najafi and also by the University of Tehran Research Department, Tehran, Iran.

REFERENCES

- Moon RT, Kohn AD, De Ferrari GV, Kaykas A. WNT and β-catenin signalling: diseases and therapies. *Nature reviews genetics* 2004; 5(9): 691-701.
- Clevers H. Wnt/β-catenin signaling in development and disease. *Cell* 2006; **127**(3): 469-480.
- 3. Dorsam RT, Gutkind JS. G-protein-coupled receptors and cancer. *Nature reviews cancer* 2007; **7**(2): 79-94.
- 4. McCrea PD, Turck CW, Gumbiner B. A homolog of the armadillo protein in Drosophila (plakoglobin) associated with E-cadherin. *Science* 1991; **254**(5036): 1359-1361.
- Peifer M, Pai LM, Casey M. Phosphorylation of the Drosophila adherens junction protein Armadillo: roles for wingless signal and zeste-white 3 kinase. *Developmental Biology* 1994; 166(2): 543-556.
- Slusarski DC, Corces VG, Moon RT. Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* 1997; **390**(6658): 410-413.
- Liu T, DeCostanzo AJ, Liu X, Wang H y, Hallagan S, Moon RT, Malbon CC. G protein signaling from activated rat frizzled-1 to the β-catenin-lef-Tcf pathway. *Science* 2001; **292**(5522):1718-1722.
- Liu X, Rubin JS, Kimmel AR. Rapid, Wnt-induced changes in GSK3β associations that regulate β-catenin stabilization are mediated by Gα proteins. *Current biology* 2005; 15(22):1989-1997.
- Katanaev VL, Ponzielli R, Sémériva M, Tomlinson A. Trimeric G protein-dependent frizzled signaling in Drosophila. *Cell* 2005; **120**(1): 111-122.
- Gao Y, Wang H y. Inositol pentakisphosphate mediates Wnt/β-catenin signaling. The journal of biological

chemistry 2007; 282(36): 26490-26502.

- 11. Koval A, Katanaev VL. Wnt3a stimulation elicits Gprotein-coupled receptor properties of mammalian Frizzled proteins. *Biochemical journal* 2011; **433**(3): 435-440.
- 12. Najafi SMA. Activators of G proteins inhibit GSK- 3β and stabilize β -catenin in Xenopus oocytes. *Biochemical and biophysical research communications* 2009; **382**(2): 365-369.
- Salmanian S, Najafi SMA, Rafipour M, Arjomand MR, Shahheydari H, Ansari S, Kashkooli L, Rasouli SJ, Saghaeian Jazi M, Minaei T. Regulation of GSK-3β and β-catenin by Gαq in HEK293T cells. *Biochemical and biophysical research communications*. 2010; **395**(4): 577-582.
- 14. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. *The journal of biological chemistry* 1951; **193**(1): 265-275.
- Morin PJ, Sparks AB, Korinek V, Barker N, Clevers H, Vogelstein B, Kinzler KW. Activation of β-catenin-Tcf signaling in colon cancer by mutations in β-catenin or APC. Science 1997; 275(5307): 1787-1790.
- Tetsu O, McCormick F. β-Catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* 1999; 398(6726): 422-426.
- He T C, Sparks AB, Rago C, Hermeking H, Zawel L, Da Costa LT, Morin PJ, Vogelstein B, Kinzler KW. Identification of c-MYC as a target of the APC pathway. *Science* 1998; **281**(5382): 1509-1512.
- Chamorro MN, Schwartz DR, Vonica A, Brivanlou AH, Cho KR, Varmus HE. FGF-20 and DKK1 are transcriptional targets of β-catenin and FGF-20 is implicated in cancer and development. *The EMBO journal* 2005; 24(1): 73-84.
- Gilchrist A, Bünemann M, Li A, Hosey MM, Hamm HE. A dominant-negative strategy for studying roles of G proteins in vivo. *The journal of biological chemistry* 1999; 274(10): 6610-6616.
- Gilchrist A, Vanhauwe JF, Li A, Thomas TO, Voyno Yasenetskaya T, Hamm HE. Gα minigenes expressing Cterminal peptides serve as specific inhibitors of thrombinmediated endothelial activation. *The journal of biological chemistry* 2001; **276**(28): 25672-25679.
- Macfarlane SR, Seatter MJ, Kanke T, Hunter GD, Plevin R. Proteinase-activated receptors. *Pharmacological reviews* 2001; 53(2): 245-282.
- Haraguchi K, Rodbell M. Carbachol-activated muscarinic (M1 and M3) receptors transfected into Chinese hamster ovary cells inhibit trafficking of endosomes. *Proceedings of the national academy of sciences of the United States of America* 1991; **88**(14): 5964-5968.
- 23. Grishina Z, Ostrowska E, Halangk W, Sahin Tóth M, Reiser G. Activity of recombinant trypsin isoforms on human proteinase-activated receptors (PAR): mesotrypsin cannot activate epithelial PAR-1,-2, but weakly activates brain PAR-1. *The British journal of pharmacology* 2005; **146**(7): 990-999.
- 24. Hussmann GP, Yasuda RP, Xiao Y, Wolfe BB, Kellar KJ. Endogenously expressed muscarinic receptors in

DOI: 10.61186/ibj.3890]

[DOR: 20.1001.1.1028852.2023.27.4.4.8]

HEK293 cells augment up-regulation of stably expressed $\alpha 4\beta 2$ nicotinic receptors. *The Journal of biological chemistry* 2011; **286**(46): 39726-39737.

- 25. Darmoul D, Gratio V, Devaud H, Lehy T, Laburthe M. Aberrant expression and activation of the thrombin receptor protease-activated receptor-1 induces cell proliferation and motility in human colon cancer cells. *The American journal of pathology* 2003; **162**(5): 1503-1513.
- Clevers H, Nusse R. Wnt/β-catenin signaling and disease. *Cell* 2012; **149**(6): 1192-1205.
- 27. Nusse R, Clevers H. Wnt/ β -catenin signaling, disease, and emerging therapeutic modalities. *Cell* 2017; **169**(6): 985-999.
- Najafi SMA. Canonical wnt signaling (Wnt/β-Catenin Pathway): A potential target for cancer prevention and therapy. *Iranian biomedical journal* 2020; 24(5): 264-275.
- 29. Shevtsov SP, Haq S, Force T. Activation of β -catenin signaling pathways by classical G-protein-coupled

receptors: mechanisms and consequences in cycling and non-cycling cells. *Cell cycle* 2006; **5**(20): 2295-2300.

- Bar Shavit R, Maoz M, Kancharla A, Nag JK, Agranovich D, Grisaru-Granovsky S, Uziely B. G protein-coupled receptors in cancer. *International journal of molecular sciences* 2016; 17(8): 1320.
- 31. Insel PA, Sriram K, Wiley SZ, Wilderman A, Katakia T, McCann T, Yokouchi H, Zhang L, Corriden R, Liu D, Feigen ME, French RP, Lowy AM, Murray F. GPCRomics: GPCR expression in cancer cells and tumors identifies new, potential biomarkers and therapeutic targets. *Frontiers in pharmacolog* 2018; 9: 431.
- 32. Lappano R, Maggiolini M. GPCRs and cancer. *Acta pharmacologica sinica* 2012; **33**(3): 351-362.
- O'hayre M, Vázquez Prado J, Kufareva I, Stawiski EW, Handel TM, Seshagiri S, Gutkind JS. The emerging mutational landscape of G- proteins and G-proteincoupled receptors in cancer. *Nature reviews cancer* 2013; 13(6): 412-424.