

# Transplantation of Olfactory Mucosa Improve Functional Recovery and Axonal Regeneration Following Sciatic Nerve Repair in Rats

Hamdollah Delaviz\*<sup>1</sup>, Mohammad Taghi Joghataie<sup>2</sup>, Mehdi Mehdizadeh<sup>2</sup>, Mehrdad Bakhtiyari<sup>2</sup>, Maliheh Nobakht<sup>2</sup> and Samideh Khoei<sup>3</sup>

<sup>1</sup>Dept. of Anatomy, Yasuj University of Medical Sciences, Yasuj; <sup>2</sup>Dept. of Anatomy and <sup>3</sup>Cellular and Molecular Research Center, Iran University of Medical Sciences, Tehran, Iran

Received 10 December 2007; revised 6 April 2008; accepted 14 April 2008

## ABSTRACT

**Background:** Olfactory ensheathing glia (OEG) has been shown to have a neuroprotective effect after being transplanted in rats with spinal cord injury. This study was conducted to determine the possible beneficial results of olfactory mucosa transplantation (OMT) which is a source of OEG on functional recovery and axonal regeneration after transection of the sciatic nerve. **Methods:** In this study, 36 adult female Sprague-Dawley rats were used. The sciatic nerve was transected in 24 rats and immediately repaired by sciatic-sciatic anastomosis, and randomly divided equally into two groups. The experimental group received the OMT at the transected site and the control group received the respiratory mucosa transplant. In another twelve rats as sham-operated animals, the sciatic nerve was exposed but no transection was made. DiI retrograde tracing was injected in the gastrocnemius muscle two months after surgery to allow visualization of the extent of axonal regeneration. Functional recovery was also assessed at 15, 30, 45 and 60 days after surgery using walking track analysis and sciatic function index (SFI) calculations. **Results:** The total number of DiI labeled motoneurons in the ventral horn (L4-L6) and the SFI scores were significantly higher in the group of rats that received olfactory mucosa rather than respiratory mucosa. **Conclusions:** The outcome indicates that olfactory mucosa is a useful treatment to improve nerve regeneration in mammals with peripheral nerve injury. *Iran. Biomed. J. 12 (4): 197-202, 2008*

**Keywords:** Olfactory mucosa transplantation (OMT), Retrograde tracing, Olfactory ensheathing glia (OEG), Functional recovery

## INTRODUCTION

Peripheral nerve injuries (PNI) are one of the most challenging problems faced by surgeons [1]. They are estimated to occur to 2.8% of all trauma patients, many of them acquire permanent disabilities and neuropathic pain as a result [2]. Peripheral nerves have the capacity to repair after lesion, but permissive environment and trophic support are required for axonal outgrowth [3]. In order to improve the functional recovery and histological outcome after PNI, the rat-sciatic-nerve model is a mainstay in the evaluation of motor and sensory nerve function [4].

Several strategies have been developed to rebuild the PN defect, including Schwann cell

transplantation [5], peripheral nerve allograft [6], fibroblast growth factor [7], bone marrow stromal cells [8], implantation of neural stem cells [9], and the use of a fibrin sealant containing neurotrophic factors [2]. Transplantation of olfactory ensheathing glia (OEG) is another existing strategy used in response to nerve injury [10]. These cells have the unique property of ensheathing the entire axonal path from olfactory mucosa in the peripheral nervous system (PNS) to the outer layer of olfactory bulb (OB) in the central nervous system (CNS), thus preventing exposure of olfactory axon to inhibitory molecules [11,12]. They share properties with both Schwann cells of (PNS) and astrocytes of the CNS [13]. Extracting these cells from OB in humans presents major difficulty, whereas the olfactory

\*Corresponding Author; Tel. & Fax: (+98-21) 88058689; E-mail: nilohamdi@yahoo.com. **Abbreviations:** OEG, olfactory ensheathing glia; OMT, olfactory mucosa transplantation; RMT, respiratory mucosa transplant; SFI, sciatic function index; PNI, peripheral nerve injuries; PNS, peripheral nervous system; CNS, central nervous system; OB, olfactory bulb

mucosa is a source of these cells by a simple biopsy through the external nares [14]. OMT induces a sustained expression of trophic factors at the lesion site in PNI [15]. The survival of some neurons that depends on the retrograde transport of trophic molecules, because such neurons die when this transport is interrupted [16]. Trophic factors of olfactory mucosa support neuron production and survival [15] and this may be the OEG work optimally in concert with connective tissue element [17]. However, PNI complete functional recovery does not occur in most cases, despite optimal surgical treatment [18]. In the present study, our purpose is to evaluate the effect of the OMT in the functional recovery and axonal regeneration of the sciatic nerve following transection.

## MATERIALS AND METHODS

**Animals.** All animal experiments were performed according to the guidelines of the Iranian Council for the Use and Care of Animals Guidelines and were approved by the Animal Research Ethical Committee of Iran Medical University (Tehran, Iran). Adult female Sprague-Dawley rats ( $n = 36$  and 200-250 g) were randomized and divided equally into three groups: OMT, respiratory mucosa transplant (RMT) and a sham group. The rats were maintained on a 12 hours light/dark cycle with free access to food and water.

**Surgery procedures and transplantation.** Rats were anesthetized with an intraperitoneal injection consisting of a combination of ketamine (80 mg/kg) and xylazine (10 mg/kg). The left sciatic nerve was exposed and transected by means of sharp microscissors near the obturator tendon at mid thigh [19]. The proximal stump was then immediately microsurgically reconnected to the distal stump with two 11-0 atraumatic sutures (Ethicon EH 7438G, Ethicon, Norderstedt, Germany). The olfactory mucosa and RM were provided as described previously [14] from same old adult Sprague-Dawley rats. Either the olfactory mucosa or RM was gently laid over the sutured epineurium in experimental and control groups, respectively. The muscles were then sutured in layers and the skin was closed. The rats were returned to their cages with access to water and food ad libitum. In the sham-operated animals, the sciatic nerve was exposed by separating the surrounding muscles in the same manner as performed in the grafted animals, but the sciatic

nerve was not transected.


**Footprint recording and analysis.** Motor functional assessment was performed 15, 30, 45 and 60 days after injury using walking track analysis [20]. The hind feet of the rats were dipped in dilute China ink and the animals were allowed to walk down an  $11 \times 45$  cm corridor into a darkened box. The floor of the corridor was covered with a sheet of paper to record the footprints. The print length (PL), toe spread from the first to the fifth toe (TS), and intermediary toe spread (IT) from the second to the fourth toe were measured on the experimental (EPL, ETS, and EIT) and normal sides (NPL, NTS, and NIT). The sciatic functional index (SFI) was calculated according to the following formula [20]:

$$\text{SFI} = -38.3 \left( \frac{\text{EPL} - \text{NPL}}{\text{NPL}} \right) + 109.5 \left( \frac{\text{ETS} - \text{NTS}}{\text{NTS}} \right) + 13.3 \left( \frac{\text{EIT} - \text{NIT}}{\text{NIT}} \right) - 8.8.$$

A SFI of zero indicates normal nerve function and -100 represents complete dysfunction.

### Retrograde tracing and histological procedures.

Two months after sciatic nerve transection and behavioral assessment, six rats from each group were used for retrograde tracing with 1, 1'-diiododecyl-3, 3', 3''-tetramethylindocarbocyanin perchlorate (DiI) from Molecular Probes (Leiden, The Netherlands; cat. No. D-282). The animals were anesthetized and the gastrocnemius muscles on the left side were exposed by an incision through the overlying skin. DiI (8-9  $\mu\text{l}$ , in 170 mg/ml DMSO) was diluted 1:10 in saline and injected into 5 locations on the body of the muscle [21] using a 10- $\mu\text{l}$  Hamilton syringe. The skin was sutured and the rats were allowed to recover. Two weeks after injection [22], all animals were transcardially perfused with 0.9% NaCl in distilled water followed by fixation with 4% paraformaldehyde (0.1 M phosphate buffer, pH 7.4) under deep anesthesia. The embedded spinal cord (L4-L6) was dissected out and cryoprotected in 30% sucrose overnight. Serial 20  $\mu\text{m}$  -thick transverse sections of the segment were made on a freezing microtome (Leica cryostat). Every second section was mounted onto gelatin-coated glass slides, cover slipped and searched for labeled neurons using fluorescent microscopy (Olympus Ax70). As previously described [2] in each spinal segment the number of labeled motoneurons of each section was summed together to give the total number of motoneurons for each rat.



**Fig. 1.** Fluorescence photomicrographs of retrogradely DiI labeled motoneurons in the left ventral horn at the level of L4-L6. The presence of the retrograde tracing in the motoneurons is indicative of regeneration of the some transected axons into OMT (**B**) and the respiratory mucosa transplant (**C**) compared to the sham group (**A**). The number of DiI labeled motoneurons in OMT rats was significantly higher than RMT animals.  $n = 6$ ,  $P < 0.05$  and scale bar, 50  $\mu\text{m}$ .

**Data analysis.** Significant differences among groups were determined by two-way ANOVA. Data are presented as the mean  $\pm$  S.E.M.  $P < 0.05$  between any two groups was considered significant according to the Bonferroni procedure.

## RESULTS


**DiI-labelled motoneurons in ventral horn.** Two months after sciatic nerve transection, the axon exhibited regeneration past the transected site, and retrograde tracing DiI had been transported to the motoneurons in the ventral horn. The average number of counted retrogradely labeled motoneurons in the left ventral horn (operated side) in the sham group that had not received transection of the sciatic nerve was (mean  $\pm$  SEM)  $132.07 \pm 4.2$  (Figs.1 and 2). This number was less in the OMT and RMT groups:  $84.76 \pm 4.5$  and  $53.23 \pm 5.61$ , respectively at the corresponding level (L4-L6). Furthermore, a Two-way ANOVA test followed by Bonferroni's test showed that there was a significant difference in the ratio of the number of labeled motoneurons in OMT rats compared to the RMT animals,  $n = 6$  and  $P < 0.05$ . This result shows that OMT to the transected sciatic nerve improves nerve regeneration.

**Functional assessment.** In sham operated animals, the hind foot toes completely spread with normal gait, with a mean SFI ( $-7.36 \pm .8$ ) at 15 days and ( $-6.22 \pm .7$ ) at 60 days. The OMT and RMT groups showed an adduction of the toes and foot drop, and were unable to bear weight. The mean SFI for the OMT rats was  $-86 \pm 2.3$  and for RMT group was  $-89 \pm 1.7$  at 15 days. At 15, 30, and 45 days, there were no statistically significant differences between the OMT and RMT groups (Fig. 3). The

mean SFI scores increased at 60 days in the OMT group when ( $-69 \pm 0.4$ ) compared with a mean of ( $-80 \pm 0.8$ ) for the RMT group (Table. 1). At this time Statistical analysis by SFI showed that there was significant differences ( $P < 0.05$ ).

## DISCUSSION

In the present work, the mature olfactory mucosa engraftment enhances axonal regeneration after sciatic nerve transection. The increase of SFI scores and DiI labeled motoneurons in L4-L6 spinal level in the OMT rats compared to control group (the RMT rats) demonstrates that OMT to the transected sciatic nerve enhances nerve regeneration and



**Fig. 2.** The total number of DiI labeled motoneurons in each group. Single asterisks indicate a significant difference between OMT and RMT, double asterisks indicate a significant difference between the sham group and both transplanted group.  $n = 6$  and  $P < 0.05$ .

**Table 1.** SFI scores for sham group, olfactory mucosa transplantation and respiratory mucosa transplantation.

Group	15 d	30 d	45 d	60 d
Sham	-9.46 ± 4.04*	-8.72 ± 2.81*	-8.61 ± 5.11*	-7.34 ± 6.12*
OMT	-98.21 ± 7.11	-86.41 ± 10.71	-81.64 ± 11.62	-79.34 ± 8.14**
RMT	-96.51 ± 9.42	-83.23 ± 6.52	-74.14 ± 4.91	-65.41 ± 11.17


\* $P < 0.05$ , sham group compared to OMT and RMT groups; \*\* $P < 0.05$ , OMT group compared to RMT group.

improves motor performance. Autograft is the most common strategy of peripheral nerve gap [23] and olfactory mucosa is a readily accessible source of OEG for autologous grafting from the olfactory system and is not an additional burden to the immune system [14, 17]. In this study, olfactory mucosa provided an appropriate microenvironment for induction and enhancement of nerve regeneration. Axonal regeneration is probably influenced by the growth factors released by OEG in the olfactory mucosa [15]. Neurotrophic factors are regulatory proteins that modulate neuronal survival, axonal growth, synaptic plasticity and neurotransmission [24]. The olfactory neuro-epithelium undergoes lifelong repair by progenitor cells, which are capable of replacing both neuronal and supporting cells [25]. Extracellular matrix proteins unable to reduce the number of axonal branches on the facial nerve injury of the rats [26], whereas OMT minimizes axonal branching after facial nerve repair in rats [15]. The improved motor recovery seen after 8 weeks post-transplantation in our study correlated with improved motor axon regeneration exemplified by the retrograde tracing in the OMT rats. Immunostaining for trophic factors at the lesion site of olfactory mucosa -transplanted rats in the transected facial nerve showed increased expression of NGF, BDNF, and FGF-2 [15].

Thus, the trophic factors within the graft act as a nerve guide and direct the outgrowing nerve fibers towards the distal nerve stump. Growth factors such as NGF and NT3 retrogradely transported in the sciatic nerve [27] and NT3 plays a role in the conveyance of trophic signals from organs to neurons such as motoneurons and proprioceptive sensory neurons in adult rats [27]. The increase of the SFI scores at 60 day after operation in the OMT rats in the present study shows that the olfactory

mucosa may provide a weaker but long-lasting secretion of neurotrophins and bFGF at the lesion site [15].

In the olfactory system, OEG can promote axonal regeneration across the PNS: CNS boundary and perhaps may become a prime candidate for cell-mediated repair following different CNS lesions [28, 29]. OEG have differential expression in the PNS and CNS and they are comprised of a heterogeneous population of cells [10]. OEG in the Lamina propria of the olfactory mucosa are responsible for histological outcome and promotion of functional recovery. Also, they exhibit a higher mitotic rate, migratory and reduced cavity and lesion site compared with OEG from the OB in the lesion spinal cord [30]. Although axonal regeneration is influenced by the growth factors that are provided by reactive Schwann cells and macrophages, the number of these cells is low in both olfactory



**Fig. 3.** Walking track analysis is showing the SFI in each group at 15, 30, 45 and 60 days after surgery. The SFI scores of OMT was significant compare to the RMT at 60 days. \* $P < 0.05$ .

mucosa and RM [14]. It is likely that not only OEG but also neuroprogenitor cells of the olfactory neuroepithelium are responsible for axonal regeneration and promotion of recovery. Multipotent progenitor cells in the adult olfactory epithelium could give rise to neurons and non-neural cells [31]. These cells in the adult human olfactory epithelium can assist spinal cord regeneration and promote functional recovery [32]. A pilot clinical study has shown that autograft of olfactory mucosa is fairly safe and feasible and may possibly promote functional recovery in chronic, severe spinal cord injury in humans [33]. Thus, OMT, including its lamina propria and olfactory neuroepithelium, is a feasible means of achieving functional recovery of peripheral nerves via axonal regeneration.

### ACKNOWLEDGEMENTS

This research study was financially supported by a grant from Iran University of Medical Sciences (Tehran). We thank Dr. Rozbehei for his excellent technical support and Hamoun Delaviz for assistance in revision from the University of Toledo College of Medicine (Ohio, USA).

### REFERENCES

- Hadlock, T., Elisseeff, J., Langer, R., Vacanti, J. and Cheny, M. (1998) A tissue-engineered conduit for peripheral nerve repair. *Arch. Otolaryngol. Head Neck Surg.* 124: 1081-1086.
- Jubran, M. and Widenfalk, J. (2003) Repair of peripheral nerve transections with fibrin sealant containing neurotrophic factors. *Exp. Neurol.* 181: 204-212.
- Yin, Q., Kemp, G.J., Yu, L.G., Wagstaj, S.C. and Frostick, S.P. (2001) Expression of Schwann cell-specific proteins and low-molecular-weight neurofilament protein during regeneration of sciatic nerve treated with neurotrophin-4. *Neuroscience* 105 (3): 779-783.
- Madorsky, S.J., Swett, J.E. and Crumley, R.L. (1998) Motor versus sensory neuron regeneration through collagen tubules. *Plast. Reconstr. Surg.* 102: 430-436.
- Bryan, D.J., Wang, K.K. and Chakalis-Haley, D.P. (1996) Effect of Schwann cells in the enhancement of peripheral nerve regeneration. *J. Reconstr. Microsurg.* 12: 439-446.
- Pollard, J.D. and Fitzpatrick, L. (1973) A comparison of the effects of irradiation and immunosuppressive agents on regeneration through peripheral nerve allografts: an ultrastructural study. *Acta Neuropathol.* 23: 166-180.
- Eckenstein, F.P. (1994) Fibroblast growth factors in the nervous system. *J. Neurobiol.* 25: 1467-1480.
- Chen, C.J., Ou, Y.C., Liao, S.L., Chen, W.Y., Chen, S.Y., Wu, C.W., Wang, C.C., Wang, W.Y., Huang, Y.S. and Hsu, S.H. (2007) Transplantation of bone marrow stromal cells for peripheral nerve repair. *Exp. Neurol.* 204: 443-453.
- Heine, W., Conant, K., Griffin, J.W. and Hoke, A. (2004) Transplanted neural stem cells promote axonal regeneration through chronically degenerated peripheral nerves. *Exp. Neurol.* 189: 231-240.
- Boyd, J.G., Skihar, V., Kawaja, M. and Doucette, R. (2003) Olfactory ensheathing cells: historical perspective and therapeutic potential. *Anat. Rec (New Anat).* 271B: 49-60.
- Doucette, R. (1991) PNS-CNS transitional zone of the first cranial nerve. *J. Comp. Neurol.* 312: 451-466.
- Pasterkamp, R.J., De Winter, F., Holtmaat, A.J. and Verhaagen, J. (1998) Evidence for a role of the chemorepellent Semaphorin III and its receptor Neuropilin-1 in the regeneration of primary olfactory axons. *J. Neurosci.* 18: 9962-9976.
- Chuah, M.I. and Au, C. (1994) Olfactory cell cultures on ensheathing cell monolayers. *Chem. Senses* 19: 25-34.
- Lu, J., Feron, F., Ho, S.M., Mackay-Sim, A. and Waite, P.M. (2001) Transplantation of nasal olfactory tissue promotes partial recovery in paraplegic adult rats. *Brain Res.* 889: 344-357.
- Guntinas-Lichius, O., Wewetzer, K., Tomov, T.L., Azzolin, N., Kazemi, S., Streppel, M., Neiss, W.F. and Angelov, D.N. (2002) Transplantation of olfactory mucosa minimizes axonal branching and promotes the recovery of vibrissae motor performance after facial nerve repair in rats. *J. Neurosci.* 22 (16): 7121-7131.
- Linden, R. (1994) The survival of developing neurons: a review of affect control. *Neuroscience* 58: 671-682.
- Franklin, R.J. (2002) Obtaining olfactory ensheathing cells from extra-cranial sources a step closer to clinical transplant-mediated repair of the CNS? *Brain* 125: 2-3.
- Schenker, M., Kraftsik, R., Glauser, L., Kuntzer, T., Bogousslavsky, J. and Barakat-Walter, I. (2003) Thyroid hormone reduces the loss of axotomized sensory neurons in dorsal root ganglia after sciatic nerve transection in adult rat. *Exp. Neurol.* 184: 225-236.

19. Chen, Z.Y., Chai, Y.F., Cao, L., Lu, L. and He, C. (2001) Glial cell line-derived neurotrophic factor enhances axonal regeneration following sciatic nerve transection in adult rats. *Brain Res.* 902: 272-276.
20. Bain, J.R., Mackinnon, S.E. and Hunter, D.A. (1989) Functional evaluation of complete sciatic, peroneal and posterior tibial nerve lesions in the rat. *Plast. Reconstr. Surg.* 83 (1): 129-138.
21. Harriott, A.M., Dessem, D. and Gold, M.S. (2006) Inflammation increase the excitability of masseter muscle afferents. *Neuroscience* 141: 433-442.
22. Tsai, E.C., van Bendegem, R.L., Hwang, S.W. and Tator, C.H. (2001) A novel method for simultaneous anterograde and retrograde labeling of spinal cord motor tracts in the same animal. *J. Histochem. Cytochem.* 49: 1111-1122.
23. Evans, P.J., Mackinnon, S.E., Best, T.J., Wade, J.A., Awerbuck, D.C., Makino, A.P., Hunter, D.A. and Midha, R. (1995) Regeneration across preserved peripheral nerve grafts. *Muscle Nerve* 18 (10): 1128-1138.
24. Jones, L.L., Oudega, M., Bunge, M.B. and Tuszynski, M.H. (2001) Neurotrophic factors, cellular bridges and gene therapy for spinal cord injury. *J. Physiol.* 533 (1): 83-89.
25. Calof, A.L. and Chikaraishi, D.M. (1989) Analysis of neurogenesis in a mammalian neuroepithelium: proliferation and differentiation of an olfactory neuron precursor *in vitro*. *Neuron* 3: 115-127.
26. Dohm, S., Streppel, M., Guntinas-Lichius, O., Pesheva, P., Probstmeier, R., Walther, M., Neiss, W.F., Stennert, E. and Angelov, D.N. (2000) Local application of extracellular matrix proteins fail to reduce the number of axonal branches after varying reconstructive surgery on rat facial nerve. *Restor. Neurol. Neurosci.* 16: 117-126.
27. Nitta, A., Ohmiya, M., Jin-Nouchi, T., Sometani, A., Asami, T., Kinukawa, H., Fukumitsu, H., Nomoto, H. and Furukawa, S. (1999) Endogenous neurotrophin-3 is retrogradely transported in the rat sciatic nerve. *Neuroscience* 88 (3): 679-685.
28. Bunge, M.B. (2002) Bridging the transected or contused adult rat spinal cord with Schwann cell and olfactory ensheathing glia transplants. *Prog. Brain Res.* 137: 275-282.
29. Mackay-Sim, A. (2005) Olfactory ensheathing cells and spinal cord repair. *Keio J. Med.* 54 (1): 8-14.
30. Richter, M.W., Fletcher, P.A., Liu, J., Tetzlaff, W. and Roskams, A.J. (2005) Lamina propria and olfactory bulb ensheathing cells exhibit differential integration and migration and promote differential axon sprouting in the lesioned spinal cord. *J. Neurosci.* 25 (46): 10700-10711.
31. Huard, J.M., Youngentob, S.L., Goldstein, B.J., Luskin, M.B. and Schwob J.E. (1998) Adult olfactory epithelium contains multipotent progenitor's cells that give rise to neurons and non-neural cells. *J. Comp. Neurol.* 400: 469-486.
32. Xiao, M., Klueber, K.M., Lu, C., Guo, Z., Marshalla, C.T., Wang, H. and Roisen, F.J. (2005) Human adult olfactory neural progenitors rescue axotomized rodent rubrospinal neurons and promote functional recovery. *Exp. Neurol.* 194: 12-30.
33. Lima, C., Pratas-Vital, J., Escada, P., Hasse-ferreira, A., capucho, C. and Peduzzi, J. (2006) Olfactory mucosa autograft in human spinal cord injury: A pilot clinical study. *JSCM* 29: 191-203.