



Chest Pain As a Sign of Cardiac Cystic Echinococcosis: A Systematic Review and Meta-Analysis

Pouriya Golmohammadi Khankandi¹, Erfan Banisefid¹, Kosar Baghernezhad¹, Rasa Beheshti¹, Sina Hamzezadeh¹, Soheil Nemati², Zahra Samadifar², Hamid Owaysee³, Elnaz Javanshir^{2*}, Amirreza Naseri¹

¹Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

²Cardiovascular Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

³Department of Infectious Disease, Tabriz University of Medical Sciences, Tabriz, Iran

OPEN ACCESS

*Corresponding Author:

Cardiovascular Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

Citation:

Golmohammadi Khankandi P, Banisefid E, Baghernezhad K, Beheshti R, Hamzezadeh S, Nemati S, Samadifar Z, Owaysee H, Javanshir E, Naseri A. Chest Pain As a Sign of Cardiac Cystic Echinococcosis: A Systematic Review and Meta-Analysis. *Iranian biomedical journal. Supplementary* (12-2024): 124.

ABSTRACT

Introduction: Human echinococcosis (CE) is a prevailing health issue in underdeveloped and developing nations. Cardiac involvement is a moderately infrequent manifestation of Echinococcus infection. The present study aimed to determine the rate of chest pain as a symptom of cardiac CE.

Search Strategy: This study included clinical studies reporting the proportion of cardiac EC patients with chest pain. Reviews, letters, preclinical studies, conference abstracts, and non-English papers were not included. A systematic search was performed in the PubMed and Scopus databases, and the risk of bias in the included studies was assessed using the JBI Case Series checklist. Meta-analysis was performed using the Comprehensive Meta-Analysis Third Edition (CMA3).

Results: The rate of chest pain in patients with cardiac CE was reported in 23 out of 3,985 studies. The rates were 41.5% (95% CI: 36.1%-47.0%) and 43.2% (95% CI: 35.1%-51.7%) based on the finite element method and the rapid-eye-movement analysis, respectively. Heterogeneity was borderline ($I^2: 49.07\%$; p value for heterogeneity = 0.004). Therefore, the results of both models were presented.

Conclusion and Discussion: Cardiac involvement of CE can be presented with chest pain, and it should be considered in suspected patients.

Keywords: Chest pain, Echinococcosis, Systematic review