

Histologic and Histomorphometric Evaluation of the Effects of Particle Size and Different Types of Allografts on Bone Regeneration in a Rat Calvarium Model

Niloufar Golafshan¹, Mohammad Mohammadi², Mohadeseh Arab Solghar², Mahsa Kalantari³, Sina Kakooei⁴, Sajjad Farrokhi^{5*}

¹Student Research Committee, Urmia University of Medical Sciences, Urmia, Iran

²Oral and Dental Diseases Research Center and Department of Periodontics, Kerman Dental School, Kerman University of Medical Sciences, Kerman, Iran

³Department of Oral and Maxillofacial Pathology, Oral and Dental Diseases Research Center, Kerman Dental School, Kerman University of Medical Sciences, Kerman, Iran

⁴Endodontology Research Center, Kerman University of Medical Sciences, Kerman, Iran

⁵Department of Periodontics, Faculty of Dentistry, Urmia University of Medical Sciences, Urmia, Iran

OPEN ACCESS

*Corresponding Author:

Dept. of Periodontics, Faculty of Dentistry, Urmia University of Medical Sciences, Urmia, Iran

Citation:

Golafshan N, Mohammadi M, Arab Solghar M, Kalantari M, Kakooei S, Farrokhi S. Histologic and Histomorphometric Evaluation of the Effects of Particle Size and Different Types of Allografts on Bone Regeneration in a Rat Calvarium Model. *Iranian biomedical journal*. Supplementary (12-2024): 176.

ABSTRACT

Introduction: Many studies have explored the influence of allograft size and type, but their findings require clarification. This study aimed to conduct a comparative histological and histomorphometric evaluation of the effect of allograft type and particle size differences on bone regeneration in a rat model, an area that has not been addressed in previous research.

Methods and Materials: Seventy male Wistar rats were randomly divided into seven groups: (1) no material, (2) 150–500 μm of freeze-dried bone allograft (FDBA), (3) 150–1000 μm of FDBA, (4) 1000–2000 μm of FDBA, (5) allograft cancellous bone block, (6) putty form allograft, and (7) particulate autogenous bone. A full-thickness flap was created, and a 7-mm defect was prepared. After filling the defects, the rats were monitored in separate cages. Light microscopy histological evaluation was carried out in the eighth week.

Results: Based on the findings, autogenous group exhibited the highest average in bone formation ($p = 0.05$). The FDBA (1000-2000 μm and 150-1000 μm) groups also displayed higher new bone formation averages than the other allograft groups ($p = 0.05$). The block and FDBA (1000-2000 μm) groups had the highest residual bone average ($p = 0.05$). The control group demonstrated the highest connective tissue average ($p = 0.05$).

Conclusion and Discussion: This study highlights the substantial impact of allograft type and particle size on bone regeneration. The autogenous bone graft consistently demonstrated the highest levels of new bone formation, indicating its superior performance within our experimental framework.

Keywords: Allografts, Bone regeneration, Dental implants