



# Effect of *Malva neglecta* Total Extract on Bleomycin-Induced Pulmonary Fibrosis in Rats

Amir Albabay<sup>1</sup>, Ali Hosseini Sharifabad<sup>1\*</sup>, Afsaneh Yegdaneh<sup>2</sup>, Mahdi Tavakolizadeh<sup>3</sup>, Hamide Dabaghydost<sup>1</sup>

<sup>1</sup>School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Science, Esfahan, Iran

<sup>2</sup>Department of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Science, Esfahan, Iran

<sup>3</sup>Department of Pharmacognosy, Faculty of Pharmacy, Zanjan University of Medical Sciences, Zanjan, Iran

## OPEN ACCESS

### \*Corresponding Author:

School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Science, Esfahan, Iran

### Citation:

Albabay A, Hosseini Sharifabad A, Yegdaneh A, Tavakolizadeh M, Dabaghydost H. Effect of *Malva neglecta* Total Extract on Bleomycin-Induced Pulmonary Fibrosis in Rats. *Iranian biomedical journal. Supplementary* (12-2024): 236.

## ABSTRACT

**Introduction:** Idiopathic pulmonary fibrosis is a chronic and progressive respiratory disease. *Malva neglecta* possesses significant anti-inflammatory and antioxidant activities. This study investigated the therapeutic effects of hydroalcoholic extract of this plant species on pulmonary fibrosis induced by bleomycin (BLM).

**Methods and Materials:** Male Wistar rats (n = 36; 180-200 g) were randomly divided into six groups (n = 6). The control group (Group I) received normal saline intratracheally (single dose) on day one. Other groups received a single dose of BLM (7.5 IU/Kg) intratracheally on the first day. The animals were then fed daily for 28 days as follows: normal saline (Group II), pirfenidone (Group III), 300, 600, and 900 mg/kg *Malva neglecta* extract (Groups IV, V, and VI, respectively). The rats were finally euthanized, and the lung tissues were removed for histological analysis and biochemical assessments.

**Result:** Intratracheal administration of BLM significantly increased the lung tissue levels of hydroxyproline, malondialdehyde, and free radicals compared to the control group ( $p = 0.001$ ). *Malva neglecta* at the dose of 900 mg/kg significantly prevented the increase of these factors compared to the BLM group ( $p = 0.001$ ). At the same dosage, the plant significantly decreased the aforementioned factors in comparison to its lower dosage ( $p = 0.01$ ). Lung tissues in BLM-treated groups showed severe tissue damage. In addition, hydroalcoholic extracts of *Malva neglecta* prevented the pathological damage of BLM on the lung tissue.

**Conclusion and Discussion:** Our findings indicate that the total extract of *Malva neglecta* may be effective in treating pulmonary fibrosis induced by BLM. This effect is likely associated with anti-inflammatory and antioxidant properties of the *Malva neglecta* extract.

**Keywords:** Bleomycin, Fibrosis, Rats

