



Using Machine Learning to Differentiate Between Radiation Necrosis and Tumor Progression in Brain Tumor Patients: A Systematic Review and Meta-Analysis

Mohammad Amin Habibi¹, Saeed Kargar Soleimanabad², Sara Faaliat², Sajjad Sibvei³, Mohammad Reza Ahmadi^{4*}, Mohammad Sina Mirjani⁴, Zahra Saadat⁵, Sajjad Rezvan⁶

¹Department of Neurosurgery, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran

²Student Research Committee, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

³Student Research Committee, Faculty of Medicine, Tehran University of Medical Science, Tehran, Iran

⁴Student Research Committee, Faculty of Medicine, Qom University of Medical Science, Qom, Iran

⁵Cardiovascular Research Centre, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

⁶Department of Radiology, Qom University of Medical Sciences, Qom, Iran

OPEN ACCESS

*Corresponding Author:

Student Research Committee,
Faculty of Medicine, Qom
University of Medical Science,
Qom, Iran

Citation:

Habibi MA, Kargar Soleimanabad S, Faaliat S, Sibvei S, Ahmadi M, Mirjani MS, Saadat Z, Rezvan S. Using Machine Learning to Differentiate Between Radiation Necrosis and Tumor Progression in Brain Tumor Patients: A Systematic Review and Meta-Analysis. *Iranian biomedical journal. Supplementary* (12-2024): 360.

ABSTRACT

Introduction: This Research has demonstrated that distinguishing between radiation necrosis (RN) and actual progression (TP) is essential for making informed clinical decisions. Multiple studies have investigated the use of machine learning (ML) to differentiate between RN and TP in patients with brain tumors. This specific study evaluated the effectiveness of ML in identifying RN and TP in brain tumor patients who have undergone radiation therapy.

Methods and Materials: This systematic review and meta-analysis followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. We searched PubMed, Embase, Scopus, and Web of Science databases from their inception until May 17, 2023, to identify relevant studies. The sensitivity and specificity of studies that used ML algorithms to distinguish between RN and TP in patients with brain tumors were pooled using a random-effects model. Statistical analysis was conducted using STATA v.17.

Results: Sixteen studies met the eligibility criteria. The pooled sensitivity was 0.84 (95% CI: 0.76-0.89), and the specificity was 0.87 (95% CI: 0.82-0.91). Specificity showed no significant heterogeneity, while sensitivity exhibited considerable heterogeneity. The pooled area under receiver operating characteristics curve was 0.89 (95% CI: 0.86-0.91). The positive and negative likelihood ratio were 6.47 (95% CI: 4.64-9.03) and 0.19, respectively. The pooled diagnostic score was 3.53 (95% CI: 2.91-4.15), and the diagnostic odds ratio was 34.19 (95% CI: 18.36-63.64).

Conclusion and Discussion: The findings indicate that ML can effectively differentiate between RN and TP. However, additional studies are required to validate these results. ML-driven imaging analysis has the potential to enhance diagnostic accuracy and patient management.

Keywords: Brain neoplasms, Radiation, Radiosurgery, Radiotherapy

