



Thyroiditis Incidence and COVID-19 Epidemic in Sanandaj, Iran: A Time-Series Analysis

Maziar Afshar¹, Abbas Aghaei², Bijan Nuri², Bahar Moasses Ghafari^{3*}

¹Student Research Committee, Kurdistan University of Medical Sciences, Sanandaj, Iran

²Department of Epidemiology and Biostatistics, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

³Department of Nuclear Medicine, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

OPEN ACCESS

*Corresponding Author:

Dept. of Nuclear Medicine,
Faculty of Medicine, Kurdistan
University of Medical Sciences,
Sanandaj, Iran

Citation:

Afshar M, Aghaei A, Nuri B, Moasses Ghafari B. Thyroiditis Incidence and COVID-19 Epidemic in Sanandaj, Iran: A Time-Series Analysis. *Iranian biomedical journal. Supplementary* (12-2024): 429.

ABSTRACT

Introduction: This study aimed to investigate the potential correlation between reported cases of thyroiditis and the COVID-19 pandemic in Sanandaj, Iran, using time-series analysis techniques. The data was collected from the two nuclear medicine centers in Kurdistan province, which are the only centers and the main referral facilities for nuclear medicine.

Methods and Materials: Based on thyroid scan findings from these two centers in 2019 and 2020, individuals diagnosed with thyroiditis were included. Time-series analysis, cross-correlation analysis, and analysis of lagged effects examined the temporal patterns and potential correlations between thyroiditis cases and COVID-19 data.

Results: The time-series analysis employed an autoregressive integrated moving average (ARIMA) model with a lag length of 3. While the cross-correlation analysis showed positive and significant correlations at lags 1 to 5, with the highest cross-correlation coefficient of 0.4674 ($p = 0.05$) at lag 3, suggesting a potential 3-month optimal lag, this lag was not meaningful. The analysis of lagged effects did not reveal a statistically significant relationship between the two variables ($F(6, 27) = 1.18, p = 0.3444$), indicating that the trends of thyroiditis and COVID-19 cases are not related and COVID-19 cannot predict subsequent thyroiditis cases.

Conclusion and Discussion: Despite clinical and observational reasoning suggesting a potential association between COVID-19 and thyroiditis, this study did not find strong evidence for a significant correlation between these two incidents in the population studied.

Keywords: COVID-19, Nuclear medicine, Thyroiditis