

Alterations in the Expression of *miR-148a-5p*, *TGF-β1*, and *TGF-βR2* in Skin Samples Exposed to Sulfur Mustard

Atiyeh Tavakoli^{1,2}, Saeed Heidari Keshel³, Mohades Valizadeh^{3,4*}

¹Molecular Genetics Researcher, Science and Technology Park of Tarbiat Modares University, Tehran, Iran; ²Molecular Genetics Researcher, Pars Genome Company, Science and Technology Park of Tarbiat Modares University, Tehran, Iran; ³Hematopoietic Stem Cell Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran; ⁴Department of Tissue Engineering and Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

OPEN ACCESS

Article type: Research Article

Received: June 30, 2025

Revised: August 18, 2025

Accepted: August 26, 2025

Published online: August 27, 2025

How to cite:

Tavakoli A, Heidari Keshel S, Valizadeh M. Alterations in the Expression of *miR-148a-5p*, *TGF-β1*, and *TGF-βR2* in Skin Samples Exposed to Sulfur Mustard. *Iran. Biomed. J.* 2025; 29(5): 344-351.



This article is licensed under a Creative Commons Attribution-NonDerivatives 4.0 International License.

ABSTRACT

Background: Sulfur mustard exposure causes chronic cutaneous injuries characterized by inflammation, fibrosis, and delayed wound healing. MiRNAs, such as *miR-148a-5p*, have been implicated in regulating the TGF-β signaling pathways involved in these processes. This study aimed to evaluate whether alterations in the expression of *miR-148a-5p*, *TGF-β1*, and *TGF-βR2* are associated with long-term SM-induced skin damage.

Methods: Skin biopsy samples were collected from 20 SM-exposed veterans and 20 healthy controls. Total RNA was extracted, and quantitative real-time PCR was performed to assess the expression levels of *miR-148a-5p*, *TGF-β1*, and *TGF-βR2*. Group differences were analyzed using a t-test, and ROC curves were generated to evaluate diagnostic performance.

Results: Expression levels of *miR-148a-5p*, *TGF-β1*, and *TGF-βR2* were significantly lower in SM-exposed skin compared with controls (*miR-148a-5p*: $p = 0.0010$; *TGF-β1*: $p < 0.0001$; *TGF-βR2*: $p < 0.0001$). Based on ROC analysis, *miR-148a-5p* and *TGF-βR2* indicated promising discriminative potential, whereas *TGF-β1* did not reach statistical significance (AUC = 0.65; $p = 0.0877$).

Conclusion: Our findings suggest that reduced expression of *miR-148a-5p* and *TGF-βR2* may contribute to SM-related skin injury. Both markers demonstrated potential diagnostic utility and could aid in risk stratification and monitoring in SM-induced skin disease, pending further validation in larger cohorts. **DOI: 10.61882/ibj.5198**

Keywords: Biomarkers, Gene expression, Mustard gas, Skin, Transforming growth factor beta

Corresponding Author: Mohades Valizadeh

Department of Tissue Engineering and Applied Cell Sciences, School of Advanced Technologies in Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran, Tel.: (+98-21) 44838970; E-mail: mohadesevali@sbmu.ac.ir, mohadeseh_vali@yahoo.com; ORCID ID: 0000-0001-5112-2698

INTRODUCTION

Sulfur mustard is a vesicant chemical warfare agent that has been utilized in many conflicts, most prominently during World War I (1914-1918) and the Iran-Iraq War (1980-1988). Its first recorded use as a chemical weapon occurred on July 12, 1917, when German forces deployed it against French troops in Belgium, resulting in 15,000 injuries within three weeks.

During World War I, approximately 77% of British chemical warfare attacks involved SM^[1]. In the Iran-Iraq War, nearly 100,000 Iranian troops and civilians were exposed to SM, with many continuing to suffer long-term effects decades later^[2].

The severity of injuries among surviving veterans varies based on multiple factors, such as the time and dose of exposure, the presence of other chemical agents,

List of Abbreviations:

AUC: area under the curve; **cDNA:** complementary DNA; **miRNA:** microRNA; **ROC:** receiver operating characteristic; **R-Smads:** receptor-regulated Smads; **SD:** standard deviation; **SM:** sulfur mustard

and individual characteristics such as age and gender. These factors influence the severity of tissue damage, the onset of symptoms, treatment requirements, and the duration of hospitalization.

SM causes extensive cellular damage through interactions with DNA and other cellular components, leading to apoptosis, necrosis, inflammation, and impaired wound healing^[3-5]. Alkylation and cross-linking of DNA significantly disrupt vital cellular functions, likely resulting in cell death or cell growth inhibition^[4,6,7]. The severity of skin damage induced by SM exposure depends on several factors, including the dose of exposure, duration of exposure, and the area of skin involved^[8-10]. SM exposure typically causes skin toxicity characterized by initial inflammation, blistering, and vesicle formation. These blisters mostly rupture, giving rise to necrotic tissue and subsequent scar formation^[7,11]. Healing from SM-induced cutaneous injury is a prolonged process and often results in residual pigmentation changes and scar tissue development^[12]. Skin lesions caused by SM are generally characterized by erythema, blistering, necrosis, and fibrosis^[13].

A major clinical challenge in SM-exposed veterans is the impaired wound healing. Beyond the direct alkylating injury caused by SM, the specific molecular drivers of this impairment remain incompletely defined^[14]. Research has highlighted a key role for miRNAs in the wound-healing process^[15]. These small non-coding RNAs are transcribed as precursors and processed in the nucleus and cytoplasm to generate mature miRNAs that regulate diverse cellular pathways^[16,17]. Notably, the miR-148 family—including *miR-148a-5p*—has been linked to the modulation of the TGF- β /Smad signaling pathway. When activated, TGF- β receptors phosphorylate R-Smads, which translocate to the nucleus and influence gene expression^[18-21]. Studies have indicated that *miR-148a-5p* regulates TGF- β signaling and is implicated in fibrotic and inflammatory processes across multiple tissues, including skin^[22,23]. *miR-148a-5p* has been reported to modulate profibrotic factors (e.g., collagen) and influence inflammatory cytokines, thereby contributing to fibrosis and inflammation^[24]. Moreover, TGF- β is a central mediator of cutaneous fibrosis and inflammation, and dysregulation of this pathway has been associated with chronic skin disease as well as SM-induced cutaneous lesions^[20,25,26]. These literature references provide biological context only and do not imply measurement of these pathways beyond the markers examined in the present study.

Currently, no specific antidotal therapy exists for SM exposure; available treatments are mainly symptom-directed, including anti-inflammatory approaches. Advancing our understanding of the molecular effects

of SM may enable the development of gene-based and targeted therapies. In this study, we examined the skin expression of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* in individuals exposed to SM. We hypothesize that chronic cutaneous lesions in SM-exposed veterans may involve dysregulation of *miR-148a-5p* and TGF- β signaling, contributing to impaired wound healing, excessive fibrosis, and persistent inflammation. This hypothesis is supported by prior literature linking altered miRNA expression—particularly *miR-148a-5p*—to the TGF- β /Smad pathway in chronic skin conditions.

By identifying the molecular mechanisms underlying compromised wound healing following SM exposure, this study aimed to identify possible biomarkers and therapeutic targets that could enhance treatment strategies for affected individuals. Furthermore, understanding the interactions between these miRNAs and TGF- β signaling pathways may provide insights into the chronic impacts of SM on skin regeneration, contributing to the design of novel gene-based therapeutic strategies.

MATERIALS AND METHODS

Study population

Twenty skin biopsy specimens were obtained from the chemical warfare-exposed veterans hospitalized in Baqiyatallah Hospital in Tehran, along with 20 control skin tissue samples sourced from the biobank of the hospital (Table 1). Biopsy samples were collected from active lesion sites characterized by erythematous plaques and pruritus, including both the epidermal and dermal layers. All the specimens were stored at -80 °C before further processing.

Site selection

Biopsies were performed only on active skin lesions, characterized by erythema (erythematous plaques) and pruritus. Lesions with necrosis or scarring (i.e., healed wounds) were not sampled. As patients presented with multiple skin lesions in different regions of the body (e.g., trunk, forearms, and legs), the biopsy site varied among individuals. In each patient, a visible, accessible, and active lesion was selected. Standardization was achieved by sampling clinically similar lesions (active and inflammatory) across all patients to ensure consistency in clinical status. A 3-mm punch biopsy was used to obtain tissue from both the epidermal and dermal layers.

Timing of biopsy

Biopsies were performed during the first clinical visit, immediately after the suitability of the lesion for sampling was confirmed by clinical examination. At the time of biopsy, no new treatments had been

Table 1. Clinical specifications of samples

Normal volunteers		SM-exposed patients	
Type	Age	Type	Age
N1	53	C1	57
N2	56	C2	53
N3	60	C3	60
N4	62	C4	62
N5	55	C5	58
N6	58	C6	63
N7	58	C7	59
N8	51	C8	58
N9	49	C9	60
N10	61	C10	49
N11	64	C11	60
N12	58	C12	61
N13	55	C13	48
N14	58	C14	55
N15	62	C15	58
N16	58	C16	60
N17	61	C17	64
N18	49	C18	61
N19	58	C19	56
N20	60	C20	54

administered, and the lesions were in their active inflammatory phase. The time between the initial SM exposure and biopsy collection spanned several years, indicating the chronic nature of the lesions. Thus, all biopsies were obtained during the chronic active phase of the disease, excluding those taken during the acute phase.

Lesion stage

All biopsies were collected from clinically active lesions, characterized as erythematous, pruritic plaques with persistent inflammation. These represented chronic active lesions—long-standing, non-healed sites—demonstrating clear inflammatory activity in the absence of necrosis, blistering, or scarring. Healed or inactive lesions were not included. Inclusion criteria for participants required a confirmed history of SM exposure, an age range similar to that of the study group, and male participants. The control group was age-matched to the exposed group, with no history of military service and residence in urban areas affected by war or chemical attacks. Exclusion criteria included individuals with occupational or environmental exposure to other chemical substances or those with pre-existing skin conditions before SM exposure.

RNA extraction and cDNA synthesis

A proteinase K treatment step was performed to eliminate the inhibitory effects of proteins in skin samples. Materials and tools required for this experiment included skin samples, diethyl pyrocarbonate-treated microtubes containing proteinase K at a concentration of 20 mg/ml (Takara, Japan), and a

thermal block set at 56 °C for one hour. RNA extraction was performed using TRIzol (Invitrogen Corp., Carlsbad, CA) following the manufacturer's protocol. The concentration of RNA was measured with a NanoDrop ND-1000 instrument, and its quality was assessed through gel electrophoresis using an ultraviolet-visible spectrophotometer. The extracted RNA was then treated with DNase I enzyme to remove residual genomic DNA. Subsequently, cDNA was synthesized using 20 ng of purified total RNA treated - DNase I and the Universal cDNA Synthesis Kit (Exiqon, Denmark). The incubation was carried out at 42 °C for one hour, followed by inactivation at 95 °C for 5 minutes to deactivate the reverse transcriptase enzyme. For the selected genes, specific primers were used, including custom primers for *miR-148a-5p* and 5S *rRNA* (Exiqon). Unfortunately, the exact catalog numbers for these primers were not provided. The primers for *TGF- β 1*, *TGF- β 2*, and *GAPDH* were synthesized by Gene Fanavaran Teb Azma (Tehran, Iran). The sequences for the primers included *TGF- β 1* forward: TGGCGATACCTCAGCAAC, *TGF- β 1* reverse: ACCCGTTGATGTCCACTTG; *TGF- β 1* receptor 2 forward: CCTATGAGCCAAGCCTATTG; *TGF- β 1* receptor 2 reverse: AATAAGTTCTGGGAAG GGAAG, *GAPDH* forward: GTGAACCATGAGAA GTATGACAAC, and *GAPDH* reverse: CATGAGTC CTTCCACGATACC.

Real-time PCR

For normalization of miRNA and mRNA expression, 5S *rRNA* and *GAPDH* were used as internal reference genes, respectively. These genes were selected based on previous validation studies in skin and wound-healing models^[27-31]. Real-time PCR was performed using SYBR Green Master Mix (Takara) on an ABI 7500 instrument (Applied Biosystems, Foster City, CA, USA) under the following conditions: initial denaturation at 95 °C for 10 minutes, followed by 45 cycles consisting of denaturation at 95 °C for 10 seconds, annealing, and extension at 60 °C for 60 seconds. Primer efficiency was assessed using LinRegPCR software (2021.2; AMC, Amsterdam, The Netherlands, <http://LinRegPCR.nl>; Table 2). Standard curve analysis was performed using serial dilutions of the template to assess the amplification efficiency of each assay. The expression levels of the candidate genes were calculated using the CT^{ΔΔ2-} method, normalized to reference genes: 5S *rRNA* for *miR-148a-5p* and *GAPDH* for *TGF- β 1* and *TGF- β 2*. All experiments were conducted in triplicate. The data were presented as mean $\pm$ SD and analyzed using a t-test. ROC curves and AUC were used to evaluate the sensitivity and specificity of the selected miRNA in distinguishing patients exposed to SM from normal controls.

Table 2. Primer amplification efficiency values (E) calculated by both serial dilution standard curve and LinRegPCR software.

Genes	Gene Efficiency based on serial dilution (E)	Efficiency determined by LinRegPCR (E)
<i>miR-148a-5p</i>	0.923	1.923
<i>5s rRNA</i>	0.912	1.912
<i>TGF-β1</i>	0.934	1.934
<i>TGF-β2</i>	0.956	1.956
<i>GAPDH</i>	0.947	1.947

An efficiency value of E = 2.0 represents 100% amplification efficiency.

Statistical analysis

All statistical analyses were performed using GraphPad Prism v.6.07 (GraphPad Software Inc., CA, USA). The normality of data was assessed using Kolmogorov-Smirnov tests, and an independent t-test was used for comparisons among the groups. A *p* value less than 0.05 was considered statistically significant.

RESULTS

Expression of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* in SM-exposed samples

Given the use of fluorescence in this study, melt curve analysis was performed using the StepOnePlus™ Real-Time PCR System to confirm the correct primer binding, ensure the amplification of specific fragments and check for the absence of non-specific products and primer dimers in the PCR product (Fig. 1). Serial dilutions were used to generate standard curves for the assessment of primer efficiency, as summarized in Table 2. The expression levels of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* were assessed in the patients exposed to SM and then compared to healthy controls using real-time PCR. The reference genes used for normalization in this analysis were *5S rRNA* and *GAPDH* (Fig. 2). Quantitative analysis revealed significant differences in the expression of these markers between the SM-exposed patients and control skin samples (*p* < 0.0001). Specifically, the expression levels of *miR-148a-5p* (Fig. 2A; *p* = 0.0010), *TGF- β 1* (Fig. 2B; *p* < 0.0001), and *TGF- β 2* (Fig. 2C; *p* < 0.0001) significantly reduced in the exposed samples compared to the controls. This reduction in expression was further analyzed using the real-time PCR analysis program (ABI; Fig. 3). Real-time PCR analysis confirmed the downregulation of these biomarkers in the SM-exposed group.

ROC curve analysis of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2*

ROC curve analysis was conducted to evaluate the diagnostic potential of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* in distinguishing SM-exposed patients from healthy controls. For *miR-148a-5p*, the ROC analysis showed a sensitivity of 100%, specificity of 65%, AUC of 0.81, and *p* = 0.0008. For *TGF- β 1*, the sensitivity was 90%, specificity 60%, AUC 0.65, and *p* = 0.0877. For *TGF- β 2*, the sensitivity was 80%, specificity 80%, AUC 0.87, and *p* = 0.0001 (Fig. 4). These findings indicate that *miR-148a-5p* and *TGF- β 2* could have a strong diagnostic potential and serve as reliable biomarkers for distinguishing SM-exposed patients from healthy ones. In contrast, *TGF- β 1* did not reach statistical significance in ROC analysis (*p* = 0.0877), suggesting limited diagnostic utility in this context. Supplementary Figure S1 represents a conceptual diagram based on the findings from a previous study^[25], in which they propose a molecular pathway that *TGF- β* induces the expression of *miR-21* and *miR-20a*. These miRNAs subsequently target and inhibit *SMAD7* expression. Overall, *miR-148a-5p* and *TGF- β 2* demonstrated strong diagnostic performance, whereas *TGF- β 1* showed limited discriminatory value.

DISCUSSION

This study demonstrates a significant reduction in the expression of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* in skin biopsy samples from individuals exposed to SM compared to healthy controls. These findings suggest a disruption of the cutaneous TGF- β signaling axis and provide a plausible molecular context for the chronic inflammatory phenotype and delayed wound healing observed in SM-related skin lesions. Interpretation of these results was confined to the markers directly measured in this study; therefore, we limited our discussion to the potential relationship between these expression changes and impaired inflammatory resolution and matrix remodeling in the skin. Our analysis focused specifically on the expression of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2*. Other molecules mentioned, such as *miR-21*, *miR-20a*, and *SMAD7*, are cited only to provide biological context from previous literature and were not evaluated in our dataset^[19,20,21].

In our cohort of SM-exposed veterans, expression levels of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* were quantified in skin biopsies, and a significant downregulation was identified relative to healthy controls. These findings indicate a reduction in the expression of key components of the TGF- β signaling




Fig 1. Melt curve analysis of (A) *miR-148a-5p*, (B) *TGF- β 1*, and *TGF- β 2* in skin tissue samples from patients with chemical injuries and healthy controls. Each curve represents a melting profile automatically generated by the real-time PCR software, where different colors and letter codes (A-F) correspond to individual wells/samples rather than distinct genes or conditions. All reactions were performed using SYBR Green, and each target displayed a single, sharp peak, confirming amplification specificity. The injury group refers to skin biopsies obtained from MS-exposed veterans, whereas the control group includes healthy individuals with no history of chemical exposure.

pathway, a mechanism that could contribute to persistent inflammation and delayed wound repair in this population. DNA damage and additional cytokine pathways were not addressed; therefore, our interpretations were limited to the observed expression changes in these three markers and their established roles in cutaneous homeostasis and healing.

Studies have linked SM exposure to broad inflammatory responses in the skin^[32,33]. Because our

data are restricted to gene expression measurements, we did not draw conclusions about pathways or molecular mechanisms that were not directly assessed in this study. Anti-inflammatory roles for *TGF- β* and IL-10 in skin biology and SM-related contexts have been indicated in earlier studies^[34,35]. In our study, reduced expression of *TGF- β 1* and *TGF- β 2* was identified in skin tissue. This reduction may contribute to incomplete inflammatory resolution, which is consistent with prior reports describing impaired TGF- β -mediated regulation in chronic cutaneous injury. Findings from transgenic




Fig. 2. Expression analysis of (A) *miR-148a-5p*, (B) *TGF- β 1*, and (C) *TGF- β 2* in skin tissues from patients with chemical injuries and healthy controls.




Fig. 3. Expression levels of (A) *miR-148a-5p*, (B) *TGF- β 1*, and (C) *TGF- β 2* in skin tissue samples from SM-exposed patients, assessed by relative quantification. Each bar represents an individual patient sample. Control samples are not shown in the Figure, as comparisons with healthy controls were performed statistically and are reported in the Results Section. Numbers below the bars indicate sample identifiers.

mouse models have demonstrated that disruption of TGF- β signaling can lead to inflammatory skin lesions^[36,37], though direct causal inferences in humans require functional validation. The central role of TGF- β signaling pathway in maintaining cutaneous homeostasis and facilitating wound repair has been well established. Accordingly, the reduced expression of *TGF- β 1* and *TGF- β 2* suggests a disruption this pathway, potentially contributing to persistent

inflammation and delayed healing in this cohort^[15,24].

This study included a modest sample size ($n = 40$; 20 patients and 20 controls), which may limit statistical power.. The cohort consisted of a highly specific population—Iranian veterans with chronic SM-related skin lesions from defined exposure regions (e.g., Oramanat: Paveh, Javanroud, Thalab Babajani, Ravansar) during the Iran-Iraq war. Therefore, the findings may not generalize to broader populations. Although this specificity provides rare and clinically valuable insight into SM-associated skin pathology, the findings should be interpreted with caution. Larger and independent studies are needed to confirm these observations and evaluate their reproducibility.

The use of *5S rRNA* and *GAPDH* as reference genes represents a methodological consideration in this study. Both genes have been validated in previous skin-related studies^[27-31]. While the stability of housekeeping genes can differ between tissues, the identical processing of all samples in this study minimizes potential bias. Future studies will include validation of multiple reference genes using the geNorm and NormFinder algorithms to ensure optimal normalization accuracy.

CONCLUSION

This study shows a significant reduction in the expression levels of *miR-148a-5p*, *TGF- β 1*, and *TGF- β 2* in the skin tissues of patients exposed to SM compared to healthy controls. Our findings underscore the critical role of TGF- β signaling pathways in regulating inflammation and wound-healing processes associated with chemical burn injuries. Dysregulation of these pathways may contribute to the chronic inflammatory state and impaired healing typically observed in dermatological conditions resulting from SM exposure. Further investigation into these pathways could provide valuable insights for developing therapeutic strategies targeting TGF- β signaling, with the potential to alleviate the long-term effects of SM exposure on skin health.

DECLARATIONS

Acknowledgments

We would like to express our sincere gratitude to Dr. Mojtaba Amiri, a dermatology specialist at Baqiyatallah Hospital, for his collaboration in sample diagnosis and his support throughout the course of this study. No artificial intelligence-assisted tools were used in the production of the submitted work.




Fig. 4. ROC curve analysis of (A) *miR-148a-5p*, (B) *TGF- β 1*, and (C) *TGF- β R2* in skin tissue samples from chemical injury patients and healthy controls. The analysis demonstrates the diagnostic potential of these biomarkers in distinguishing SM-exposed patients from healthy controls.

Ethical approval

The study was approved by the Ethics Committee of the Islamic Azad University, Science and Research Branch, Tehran, Iran (ethical code: IR.SBMU.RETECH.REC.1404.139).

Consent to participate

All participants voluntarily agreed to take part in this study. Written informed consent was obtained from each individual enrolled in the research.

Consent for publication

All authors reviewed the results and approved the final version of the manuscript.

Authors' contributions

AT: performed experiments and collected data; SHK: conducted data analysis and contributed to scientific editing; MV: designed the study, supervised the research, and drafted the manuscript.

Data availability

All relevant data can be found within the manuscript.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Supplementary information

The online version contains supplementary material.

REFERENCES

1. Rosenblatt DH, Small MJ, Kimmell TA, Anderson AW. Background chemistry for chemical warfare agents and decontamination processes in support of delisting waste streams at the US army dugway proving ground. Environmental Assessment Division. Tech Rep. 1996.
2. Khateri S, Ghanei M, Keshavarz S, Soroush M, Haines D. Incidence of lung, eye, and skin lesions as late complications in 34,000 Iranians with wartime exposure to mustard agent. *J Occup Environ Med*. 2003;45(11):1136-43.
3. Ghasemi BM, Karami GR, Pourfarzam SH, Emadi S, Ghasemi H. Late concurrent ophthalmic, respiratory, coetaneous and psychiatric complications of chemical weapons exposure in 479 war patients. *Daneshvar Med*. 2007;70(4):81-92.
4. Kavehie B, Faghihzadeh S, Eskandari F, Kazemnejad A, Ghazanfari T, Soroosh MR. Studying the surrogate validity of respiratory indexes in predicting the respiratory illnesses in wounded people exposed to sulfur mustard. *Arak Uni Med Sci*. 2011;13(4):75-82.
5. Kehe K, Szinicz L. Medical aspects of sulphur mustard poisoning. *Toxicology*. 2005;214(3):198-209.
6. Jowsey PA, Williams FM, Blain PG. DNA damage responses in cells exposed to sulphur mustard. *Toxicol lett*. 2012;209(1):1-10.
7. Kehe K, Thiermann H. Clinical picture of sulfur mustard poisoning. *Toxicol*. 2009;263(1):1.
8. Graham JS, Schoneboom BA. Historical perspective on effects and treatment of sulfur mustard injuries. *Chem Biol Interac*. 2013;206(3):512-22.
9. Shakarjian MP, Heck DE, Gray JP, Sinko PJ, Gordon MK, Casillas RP, et al. Mechanisms mediating the vesicant actions of sulfur mustard after cutaneous exposure. *Toxicol Sci*. 2010;114(1):5-19.
10. Sepehr A, Aghamohammad S, Ghanavati R, Talebi M, Pourshafie MR, Rohani M. Role of native probiotic *Lactobacillus* species via TGF- β signaling pathway modulation in CRC. *Iran Biomed J*. 2023;28(4):168-178.

11. Salamati P, Razavi SM. The late clinical and forensic symptoms and signs of sulfur mustard. *Burns*. 2015;41(6):1370-1.
12. Poursaleh Z, Ghanei M, Babamahmoodi F, Izadi M, Harandi AA, Emadi SE, et al. Pathogenesis and treatment of skin lesions caused by sulfur mustard. *Cutan Ocul Toxicol*. 2012;31(3):241-9.
13. Feister AJ. Medical defense against mustard gas toxicity: Mechanisms, pharmacology, implications. CRC-Press; 1991.
14. Tahmasbpour E, Emami SR, Ghanei M, Panahi Y. Role of oxidative stress in sulfur mustard-induced pulmonary injury and antioxidant protection. *Inhal Toxicol*. 2015;27(13):659-72.
15. Mann J, Mann DA. Epigenetic regulation of wound healing and fibrosis. *Curr Opin Rheumatol*. 2013;25(1):101-7.
16. Ambros V. The functions of animal microRNAs. *Nature*. 2004;431(7006):350-5.
17. Cullen BR. Transcription and processing of human microRNA precursors. *Mol Cell*. 2004;16(6):861-5.
18. Li J, Song Y, Wang Y, Luo J, Yu W. MicroRNA-148a suppresses epithelial-to-mesenchymal transition by targeting ROCK1 in non-small cell lung cancer cells. *Mol Cell Biochem*. 2013;380(1-2):277-82.
19. Valizadeh M, Soleimani M, Irani S, Tavallaei M. Evaluation of changes in miR-20a expression changes in war wounded exposed to mustard gas. *J Anim Biol*. 2023;15(3):281-91.
20. Valizade M, Irani S, Tavallaei M, Soleimani M. Differential expression of miR-21-5p, miR-20a-5p, TGF- β 1, and TGF- β receptor 2 in skin, serum, and lung samples exposed to sulfur mustard. *Iran J Allergy Asthma Immunol*. 2023;22(4):366-78.
21. Wang H, Pan J-Q, Luo L, Ning X-J, Ye Z-P, Yu Z, et al. NF- κ B induces miR-148a to sustain TGF- β /Smad signaling activation in glioblastoma. *Mol Cancer*. 2015;14:2.
22. Miscianinov V, Martello A, Rose L, Parish CL, Brown C, Watson A, et al. MicroRNA-148b targets the TGF- β pathway to regulate fibrosis. *Matrix Biol*. 2018;68(69):47-60.
23. Babalola O, Mamalis A, Lev-Tov H, Jagdeo J. The role of microRNAs in skin fibrosis. *Arch Dermatol Res*. 2013;305(9):763-76.
24. Deng Z, Fan T, Xiao C, Tian H, Zheng Y, Li C, et al. TGF- β signaling in health, disease and therapeutics. *Signal Transduct Target Ther*. 2024;9:61.
25. Zhu L, Han X, Gao M, et al. Exosomal miR-148a-3p suppresses TGF- β /Smad4-mediated fibrosis and inflammation. *J Transl Med*. 2025;23:540.
26. Valizadeh M, Soleimani M, Irani S, Tavallaei M. Decreased expression of TGF- β 1 and TGF- β 2 in skin lesions induced by sulfur mustard in Iranian war veterans. *Iran J Allergy Asthma Immunol*. 2015;14(4):332-6.
27. Suo L, Cheng J, Yuan H, Jiang Z, Tash D, Wang L, et al. miR-26a/30d/152 are reliable reference genes for miRNA quantification in skin wound age estimation. *Forensic Sci Res*. 2023;8(3):230-40.
28. Etich J, Bergmeier V, Pitzler L, Brachvogel B. Identification of a reference gene for the quantification of mRNA and miRNA expression during skin wound healing. *Connect Tissue Res*. 2017;58(2):196-207.
29. Luo M, Wang L, Zhu W, Fu J, Song F, Fang M, et al. Identification and characterization of skin color microRNAs in koi carp (*Cyprinus carpio* L.) by Illumina sequencing. *BMC Genomics*. 2018;19(1):779.
30. Barber RD, Harmer DW, Coleman RA, Clark BJ. GAPDH as a housekeeping gene: Analysis of GAPDH mRNA expression in a panel of 72 human tissues. *Physiol Genomics*. 2005;11;21(3):389-95.
31. Arai K, Okabe M, Kobashi D, Ichimura K, Fathy M, Oba J, et al. Importance of housekeeping gene optimization for the analysis of mRNA expression during wound healing in a third-degree burn injury model. *J Burn Care Res*. 2023;44(1):146-57.
32. Renshaw B. Mechanisms in production of cutaneous injuries by sulfur and nitrogen mustard. Chemical warfare agents and related chemical problems. 1946;4(23):479-518.
33. Smith KJ, Hurst CG, Moeller RB, Skelton HG, Sidell FR. Sulfur mustard: Its continuing threat as a chemical warfare agent, the cutaneous lesions induced, progress in understanding its mechanism of action, its long-term health effects, and new developments for protection and therapy. *J Am Acad Dermatol*. 1995;32(5):765-76.
34. Qabar A, Nelson M, Guzman J, Corun C, Hwang B-J, Steinberg M. Modulation of sulfur mustard induced cell death in human epidermal keratinocytes using IL-10 and TNF- α . *J Biochem Mol Toxicol*. 2005;19(4):213-25.
35. Saki S, Monjezi S, Ghaffari F, Orak G, Bavarsad SS, Khedri A, et al. Unraveling the exosome-miR-133a axis: targeting TGF- β signaling via WJ-MSC-derived exosomes for anti-fibrotic therapy in liver fibrosis. *Iran Biomed J*. 2024;28(5 and 6):235-244.
36. Kulkarni AB, Huh CG, Becker D, Geiser A, Lyght M, Flanders KC, et al. Transforming growth factor beta 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA*. 1993;90(2):770-4.
37. Shojaei Jeshvaghani Z, Soleimani M, Asgharpour S, Arefian E. Latency-associated transcript-derived MicroRNAs in herpes simplex virus type 1 target SMAD3 and SMAD4 in TGF- β /smad signaling pathway. *Iran Biomed J*. 2021;25(3):169-179.