RNA Expression Analysis of Mycobacterial Methyltransferases Genes in Different Resistant Strains of *Mycobacterium tuberculosis*

Samira Tarashi^{1,2}, Mohammad Saber Zamani³, Golnaz Bahramali^{4,5}, Andrea Fuso⁶, Farzam Vaziri^{1,2}, Morteza Karimipoor⁷, Abolfazl Fateh^{1,2*}, Seyed Davar Siadat^{1,2*}

Department of Mycobacteriology and Pulmonary Research, Pasteur Institute of Iran, Tehran, Iran;
 Microbiology Research Center, Pasteur Institute of Iran, Tehran, Iran;
 Immunoregulation Research Center, Shahed University, Tehran, Iran;
 Department of Hepatitis and AIDS, Pasteur Institute of Iran, Tehran, Iran;
 Viral Vaccine Research Center, Pasteur Institute of Iran, Tehran, Iran;
 Medicine, Sapienza University of Rome, Italy;
 Molecular Medicine Department, Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran

OPEN ACCESS ABSTRACT

Received: 15 July 2021 Accepted: 3 August 2021 Published online: 12 February 2022

Background: Tuberculosis infection still represents a global health issue affecting patients worldwide. Strategies for its control may be not as effective as it should be, specifically in case of resistant strains of *M.tb*. In this regard, the role of mycobacterial MTases in TB infection can be fundamental, though it has not been broadly deciphered.

Methods: Five resistant isolates of *M.tb* were obtained. *M.tb* H37Rv (ATCC 27249) was used as a reference strain. Seven putative mycobacterial MTase genes (Rv0645c, Rv2966c, Rv1988, Rv1694, Rv3919c, Rv2756c, and Rv3263) and Rv1392 as SAM synthase were selected for analysis. PCR-sequencing and qRT-PCR were performed to compare mutations and expression levels of MTases in different strains. The $2^{-\Delta\Delta Ct}$ method was employed to calculate the relative expression levels of these genes.

Results: Only two mutations were found in INH^R strain for *Rv3919c* (T to G in codon 341) and *Rv1392* (G to A in codon 97) genes. Overexpression of *Rv0645c*, *Rv2756c*, *Rv3263*, and *Rv2966c* was detected in all sensitive and resistant isolates. However, *Rv1988* and *Rv3919c* decreased and *Rv1694* increased in the sensitive strains. The *Rv1392* expression level also decreased in INH^R isolate.

Conclusion: We found a correlation between mycobacterial MTases expression and resistance to antibiotics in *M.tb* strains. Some MTases undeniably are virulence factors that specifically hijack the host defense mechanism. Further evaluations are needed to explore the complete impact of mycobacterial MTases within specific strains of *M.tb* to introduce novel diagnosis and treatment strategies. *DOI:* 10.52547/ibj.26.3.240

Citation:

Tarashi S, Zamani MS, Bahramali G, Fuso A, Vaziri F, Karimipoor M, Fateh A, Siadat SD. RNA Expression Analysis of Mycobacterial Methyltransferases Genes in Different Resistant Strains of Mycobacterium tuberculosis. Iranian biomedical journal 2022; 26(3): 240-251.

Keywords: Drug resistance, Methyltransferases, Mycobacterium tuberculosis, S-adenosylmethionine

Corresponding Authors:

Abolfazl Fateh

Pasteur Institute of Iran, No. 358, 12th Farvardin Ave., Jomhhoori St., Tehran 1316943551, Iran; E-mail: afateh2@gmail.com

Seyed Davar Siadat

Pasteur Institute of Iran, No. 358, 12th Farvardin Ave., Jomhhoori St, Tehran 1316943551, Iran; E-mail: d.siadat@gmail.com

List of Abbreviations:

DST, drug susceptibility testing; **INH**, Isoniazid; **INH**^R, isoniazid resistance; *M.tb*, *Mycobacterium tuberculosis*; **MDR**, multidrugresistant; **MTases**, methyltransferases; **qRT-PCR**, quantitative reverse transcription-polymerase chain reaction; **RIF**, rifampin; **RIF**^R, rifampin resistance; **SAM**, S-adenosylmethionine; **TB**, tuberculosis; **WHO**, World Health Organization; **XDR**, extensively drugresistant

INTRODUCTION

uberculosis is an infectious disease mostly Taffecting the lungs and is caused by M.tb. Historically, TB is defined as one of the most devastating epidemiological concerns in human health. According to the latest WHO report in 2019, 10 million new TB cases and 1.2 million TBrelated deaths were estimated^[1]. The importance of this infection in healthcare settings increases with the appearance of resistant strains, specifically, MDR-TB (resistant to INH and RIF), and XDR-TB (resistant to INH and RIF plus fluoroquinolones and at least one injectable agent, including amikacin, kanamycin, and capreomycin) strains^[2]. Resistance to a range of antibiotics is an important feature of M.tb due to the fact that these pathogens have in-built complex regulation machinery. In 2019, 465,000 individuals were infected with drug-resistant TB^[1]. Early diagnosis and accurate treatment are two major objectives in TB infection control^[3]. Therefore, delay in the correct diagnosis of M.tb strains and proper treatment are the most significant causes of TB mortality^[4]. Despite the intense research, the control of TB is still largely suboptimal, and the mechanisms of its progression cannot completely be explained. The methylation process is one of the most prominent regulatory mechanisms that has attracted enormous interests in TB researches^[5].

Some of the microorganisms induce methylation in the genome of the host cells or even their own genome to exacerbate virulence and decrease immune responses^[6,7]. Although mycobacterial methylation has been addressed, little information is available in drugresistant M.tb strains^[8]. The process of methylation is catalyzed by different MTases^[9]. About 121 different MTases have been coded by the genome of M.tb, which employs several substrates, including DNA, RNA, protein, intermediates of mycolic acid biosynthesis, and other fatty acids^[10]. Among them, methylation of DNA and proteins (histones) are known as two most important epigenetic modifications, which notably regulate gene function without any direct modification in the DNA sequence^[11]. Recently, M.tb has been recognized as an epigenome regulator controlling many cellular processes involved in host immune system^[12]. The inter-talk between epigenetic modifications and M.tb is also very remarkable in the pathogenicity^[7]. About 65% of the *M.tb* genome consists of GC-rich regions; therefore, DNA methylation introduced as a highly efficient mechanism to control gene expression^[10].

Overall, different mycobacterial MTases exert their regulatory function on the host cell (mostly genes

involved in immune response) or their own genome to finally exacerbate infection by increasing virulence and decreasing immune reactions^[5]. Lately, it has been indicated that the methylation at different stages of TB infection, as a potential biomarker, requires further consideration^[7], and assessment of mycobacterial MTases may revolutionize the current view on TB infection. It is expected the evaluation of the possible between different mycobacterial expressions and TB resistance can be helpful for developing some strategies to accurately prevent, treat, and control TB. Accordingly, the aim of the current study was to assess the expression level of mycobacterial MTases in different resistance strains of M.tb (sensitive, INH^R, Rif^R, MDR, and XDR). The presented data may shed new light on host susceptibility and TB severity.

MATERIALS AND METHODS

Mycobacterial strains and DST

Five phenotypically characterized resistant isolates of M.tb were obtained from the Mycobacteriology and Pulmonary Research Department of Pasteur Institute of Iran, Tehran. These strains included sensitive, INH^R, Rif^R, MDR, and XDR. H37Rv strain (ATCC 27249) was used as a reference strain. All isolates were grown to mid log phase in Middlebrook 7H9 medium supplemented with 0.05% Tween 80, 10% (v/v) oleic acid, albumin, dextrose, catalase^[13]. The proportion method was performed using the Lowenstein-Jensen medium^[14]. The critical concentrations of DST were $0.2 \mu g/mL \text{ (INH)}, 40 \mu g/mL \text{ (RIF)}, 2.0 \mu g/mL$ (ethambutol), 4.0 μg/mL (streptomycin), 40 μg/mL (kanamycin), 30 µg/mL (amikacin), 40 µg/mL (capreomycin), and $4.0~\mu g/mL$ (ofloxacin) for phenotypic confirmation^[15]. All experiments were conducted in accordance with the guidelines approved by the Centers for Disease Control and Prevention $(USA)^{[16]}$.

Selection of mycobacterial MTase genes

Putative mycobacterial MTase genes were selected based on searching from the TubercuList database (http://genolist.pasteur.fr/TubercuList/). The important mycobacterial MTase genes and their related functions are presented in Table 1. After review of MTase genes, we selected seven putative genes (*Rv0645c*, *Rv2966c*, *Rv1988*, *Rv1694*, *Rv3919c*, *Rv2756c*, and *Rv3263* for analysis. All the selected genes were SAM-dependent mycobacterial MTases. In addition, *Rv1392* as SAM synthase was selected for the evaluation of SAM biosynthesis in different *M.tb* strains^[17].

Table 1. Mycobacterial MTases

Mycolic acid MTases			Protein MTases	Other MTases
Rv3392c (CmaA1)	Rv1317c (AdaA-AlkA) (Neutralization of mutagenic effect of alkylation stress by demethylation of methylated bacterial DNA)	Rv2118c (tRNA MTases)	Rv1988 (Methylation of H3 in the host cell)	Rv0447c (UfaA) (tuberculostearic acid MTases)
Rv0503c (CmaA2)	Rv1316c (AdaB/OGT) (Neutralization of mutagenic effect of alkylation stress by demethylation of methylated bacterial DNA)	Rv2966c (Methylation of mycobacterial 16SrRNA) (Rv2666c is also a DNA MTase which often methylate the non- CpG regions of host genome) (Rv2966c is a histone MTase in the genome of host cells)	Rv2966c (Methylation of H4 and H3 in the host cell)	Rv2952 (Phthiotriol/phenolp hthiotriol dimycocerosates MTases)
Rv0645c (MmaA1)	Rv3263 (MamA) (Protection of <i>M.</i> tuberculosis in hypoxic status)	Rv2372c (16SrRNA MTase)	Rv1198 (Methylation of H3 in the host cell)	Rv2954c (methylation of phenolic glycolipids)
Rv0644c (MmaA2)	MamB	Rv1988 (23SrRNA MTase) (Increasing of Macrolide- Lincosamides-Streptomycin resistance)	SET8 monomethylase) (<u>H4</u>	Rv2955c (methylation of phenolic glycolipids)
Rv0643c (MmaA3)	Rv2756c (HsdM)	Rv1694 (23S/16SrRNA MTase) (Increasing of Capreomycin resistance)	SUV39H1 (Methylation of mycobacterial Histone like proteins HupB)	Rv2956 (methylation of phenolic glycolipids)
Rv0642c (MmaA4)	-	Rv3919c (16SrRNA MTase) (Increasing of Streptomycin resistance)	-	-
Rv0470c (PcaA/UmaA2)	-	-	-	-
Rv0469 (UmaA)	-	-	-	-

Multifunctional mycobacterial MTases are shown in bold, and some enzymes affecting the genome of host cells are marked by underline. The other enzymes modify the methylation pattern of mycobacterial genome.

Evaluation of mycobacterial MTase genes by PCR sequencing

To confirm the presence of these mycobacterial MTase genes in selected strains, we extracted genomic DNA from the mycobacterial culture in 7H9 media using DNA Technology kit (DNA Technology Co., Moscow, Russia) according to the manufacturer's instructions. The extracted DNA was dissolved in 30 μ l

of diethyl pyrocarbonate treated water, and a 10-µl aliquot of total DNA was used for a 25-µl PCR mixture. Table 2 lists the primers employed to amplify the selected regions of mycobacterial DNA in each studied strain. The amplification parameters are shown in the supplementary data (Table S1). The PCR products were analyzed by agarose gel electrophoresis (Fig. S1). Positive and negative controls with or

Table 2. Primers used for PCR and qRT-PCR in this study

Gene	Sequence (5'-3')	Tm	Product length (bp)
Rv0645c	F-ACCATATTACGAAGAGTCACA	54.12	195
	R-ACGTCGAGTAGCGTCATC	56.62	
Rv3263	F-AATTGCTTCAAGTCACCTA	51.65	194
	R-GTTCAGATCGTTCGCATC	53.75	
Rv2756c	F-AGGATGTCTCGATCTATG	49.67	190
	R-CAGTCTTTGATGTTGAAC	48.62	
D 20//	F-TCGCTATTCAACATCGTGACT	57.5	192
Rv2966c	R-CGCACCGGAGAGACCTAG	58.89	
D 1000	F-CGATTCCCTGGCATTACC	55.17	192
Rv1988	R-AGAAGCGAATTTACATACGA	52.48	
Rv1694	F-ATATCGTTCCACTGGTGA	52.38	196
KV1094	R-AGGAAGTACTCGACATTG	50.87	
D 2010	F-CAGGTAGTTCTCCTAGAA	48.88	188
Rv3919c	R-CTCCATTTCGTCAACTTG	51.07	
D 1202	F-GCTGTTTACCAGTGAGTC	52.65	198
Rv1392	R-TTGGTGATGTCGGCAAAC	55.91	
G 4	F-CACTACGAGGTCGATCTA	52.13	180
SecA	R-GACGATGTAGTCCTTGTC	51.83	

without the genomic DNA of *M.tb* complex were included in each run. Afterward, all the PCR products were purified by PCR purification kit (Thermo ScientificTM, USA) and sequenced to evaluate the possible mutations. Raw data was analyzed by MEGA 5.2 software.

Evaluation of mycobacterial MTase genes expression

The total mycobacterial RNA was isolated using Trizol kit (Plus RNA; Invitrogen, USA) and according to a method described previously^[18]. Reverse transcription of complementary DNAs and qRT-PCR were performed simultaneously using a QuantiTect SYBR Green RT-PCR Kit (Qiagen, Germany) and a Light Cycler 2.0 instrument (Roche, Switzerland). Each sample was performed in triplicate in a 10-μL capillary tube containing 5 μL of 2× QuantiTect SYBR Green RT-PCR Master Mix, 0.5 μL of a 0.5-μM solution of each primer, 3 μL of RNase-free water, and 1 μL of template RNA (100 ng/μL). The primers used for qRT-PCR are listed in Table 2. The assay conditions were represented in supplementary data

(Table S2). The secA gene was used for normalization as a housekeeping gene. The $2^{-\Delta\Delta Ct}$ method was applied to calculate the relative expression levels of mycobacterial MTase genes^[17].

Statistical analysis

GraphPad Prism 8 (GraphPad, La Jolla, CA, USA) was used for statistical analyses and drawing graphs. Data were expressed as mean \pm SEM. Differences between the two groups were assessed using the nonparametric test, Mann–Whitney U test. A p value ≤ 0.05 was considered statistically significant.

RESULTS

The susceptibility of all five *M.tb* isolates was confirmed by DST. No growth of sensitive and H37Rv isolates was observed in the Lowenstein-Jensen medium combined with the mentioned antibiotics, i.e. INH, RIF, ethambutol, streptomycin, kanamycin, amikacin, capreomycin, and ofloxacin. The resistance to streptomycin was detected in RIF^R isolate, as well as in MDR and XDR isolates.

The important mycobacterial MTase genes are categorized in Table 1 considering their substrate. In the current study, the selection of mycobacterial MTase genes for analysis was based on their category and dependent on SAM. The relative expression level of each evaluated gene compared to H37Rv has been presented in Figure 1. Rv0645c was selected as a SAM-dependent mycolic acid MTase, which increased in all sensitive (p = 0.006), INH^R (p = 0.00), RIF^R (p = 0.007), MDR (p = 0.0005), and XDR (p < 0.0001) isolates.


The overexpression of a multifunctional mycobacterial MTase gene), Rv2966c, foundin sensitive (p = 0.02), INH^R (p = 0.002), RIF^R (p =0.002), MDR (p = 0.005), and XDR (p = 0.0002) isolates. Besides, three genes, including Rv1988, Rv1694, and Rv3919c, were selected from RNA MTase category to be evaluated in antibiotic resistance. The Rv1988 has been identified as a histone MTase in host cells during TB infection^[19]. Its downregulation was shown in sensitive (p = 0.02) isolate, and its upregulation was found in INH^{R} (p = 0.0008), RIF^{R} (p = 0.0008) = 0.0005), MDR (p = 0.0007), and XDR (p = 0.001) isolates. Rv1694 (tlyA) is a 16S rRNA/23S MTase inducing capreomaycin sensitivity, which upregulated only in sensitive (p = 0.02) isolate and downregulated in XDR (p = 0.002), INH^R (p = 0.02), RIF^R (p = 0.00) 0.009), and MDR (p = 0.0006). Rv3919c (gidB) gene, a mycobacterial 16S rRNA MTase, is involved in the induction of streptomycin resistance in M.tb. We observed a mutation in INH^R strain for Rv3919c (T to G in codon 341). Its expression level showed downregulation only in sensitive (p = 0.0003) and upregulation in INH^R (p = 0.0002), RIF^R (p = 0.0005), MDR (p = 0.0005), and XDR (p = 0.001) isolates. Rv3263 and Rv2756c genes were selected for the evaluation of DNA MTases. The activity of Rv3263 (MamA) has been confirmed in the hypoxic conditions^[20]. Its upregulation was detected in sensitive (p < 0.0001), INH^R (p = 0.002), RIF^R (p = 0.0008), MDR (p = 0.0007), and XDR (p = 0.0006). The expression level of Rv2756c (HsdM) increased in all sensitive (p = 0.03), INH^R (p < 0.0001), RIF^R (p < 0.0001) 0.0001), MDR (p = 0.0002), and XDR (p = 0.0005) isolates.

Finally, in addition to mycobacterial MTase genes, Rv1392 was selected for comparing the SAM biosynthesis in different M.tb strains. Rv1392, a SAM synthase, is known as a fundamental member of SAM synthases family^[17]. A mutation (G to A in codon 97) was detected in INH^R strain for this gene. The Rv1392 expression level decreased only in INH^R (p = 0.2) and increased in sensitive (p = 0.01), RIF^R (p = 0.0009), MDR (p = 0.001), and XDR (p = 0.0002) isolates.

DISCUSSION

In this study, we evaluated the relationship between mycobacterial MTases expression in different resistant strains of M.tb. Also, we compared four categories of mycobacterial MTase genes in five sensitive and resistant strains of M.tb. The mycobacterial MTases can be divided into DNA MTases, RNA MTases, protein MTases, mycolic acid MTases, and other MTases^[5]. These enzymes transfer a methyl (-CH3) group from methyl donors such as SAM to nucleic acids, proteins, lipids, and secondary metabolites. Detection of methylated cytosine in H37Rv and its absence in H37Ra may indicate the possible role of mycobacterial MTases in the pathogenesis^[21]. More than 1% of human genes have been predicted to encode MTases, while this number has increased to 3% in M.tb, in spite of its genome reduction^[5]. Sharma et al. [22] detected 17 putative genes coding for proteins with methylase activity in M.tb. The identification of these genes is based on the sequence analysis, SAM binding site prediction, and gene binding domains.

We reported the overexpression of Rv0645c (methoxy mycolic acid synthase 1) as a mycolic acid MTase gene with no mutation in all sensitive and resistant M.tb isolates. The upregulation of Rv0645c was significantly higher in XDR than other isolates. A study compared differences in the cell wall thickness of sensitive (15.6 \pm 1.3 nm), MDR (17.1 \pm 1.03 nm) and XDR (20.2 ± 1.5 nm) isolates $(p < 0.05)^{[23]}$. This result suggests the possible reason for significant overexpression of Rv0645c in XDR in comparison to other strains. The suitable structure of mycolic acid is important in both the permeability of the cell envelope and resistance to antimycobacterial agents. Although various mycolic acid MTases are involved formation of different mycolic acid structures^[24], the exact role of these methylases in the pathogenicity of *M.tb* is unclear yet. Nevertheless, many of the mycolic acid MTase genes are conserved, and no mutation has been detected. In this sense, they may play a critical function in the *M.tb* pathogenicity and have a potential to introduce as a new target in the development of anti-mycobacterial drugs^[25]. The upregulation of Rv0645chas been confirmed during primary infection of macrophages^[26]. This result suggests that the intracellular M.tb actively senses its environment and repairs or synthesizes its cell wall and other components. In addition, a mouse model study showed that the deletion of mycolic acid MTase genes leads to the disruption of the mycolic acid structure, thereby resulting in sensitivity to detergents and causing the general loss of mycobacterial acid-fast feature^[27].






Fig. 1. Expression level of mycobacterial MTases and Rv1392 in different resistant strains of M.tb; Mann–Whitney U test.

Expression Level Fold Change

Our investigation explored the overexpression of Rv2756c and Rv3263, mycobacterial DNA MTases, in all sensitive and resistant M.tb isolates. DNA methylation epigenetically affects the tolerance, virulence, antibiotic resistance, survival during stress conditions, and intensity of pathogenicity^[8,28]. An evaluation on DNA methylation diversity in M.tb has indicated some mutations in certain mycobacterial DNA MTases such as Rv2756c and Rv3263, affecting their activity in lineages of $M.tb^{[29]}$. The activity of Rv3263 was detected in all analyzed isolates, except Beijing strains that had high potency to cause active TB and drug resistance. Nevertheless, this fact should not be ignored that 70% of mycobacterial MTases often belongs to a multigene family that may influence the expression level of each other^[10]. As an example, the Rv3263 site shares overlapping -10 sigma factor binding site, which is directly bound by the RNA polymerase during the initiation of transcription process and regulates gene expression in response to numerous environmental conditions such as stress and antibiotic resistance^[20]. However, this document could not establish a direct association between mutation. multigene, and lack of mycobacterial MTase activity, and DNA MTase efficiency has been proved to be particularly variable, despite the presence of their genes^[29]. A point to the role of different MTases has been highlighted by Grover *et al.*^[10] who observed Rv2756c only in the pathogenic mycobacteria, reflecting the importance of mycobacterial MTases in the pathogenesis. In general, there is a very limited evidence to clearly explain the cause of such expression modifications of DNA MTases in different resistant and sensitive strains of M.tb, and more evaluations are needed to explore their impact.

Among various groups of mycobacterial MTases, some are multifunctional. Rv2966c has been introduced as one of the most important MTases that has been categorized as DNA MTase, protein MTase, and also RNA MTase^[22,30,31]. We reported the upregulation of Rv2966c in the sensitive and drug-resistant mycobacterial strains. Upregulation of this gene appears to be associated with a greater need for the suppression of host gene expression during infection by such mycobacterial strains. Indeed, Rv2966c is known as a specific mycobacterial DNA MTase, which is capable of host genes' nethylation, particularly in non-CpG sites^[30]. Notably, Rv2966c can also interact with histone proteins in the host cells^[22] and is a highly specific RNA MTase that methylates 16S rRNA of M.tb ribosome^[31]. In addition to Rv2966c, other mycobacterial RNA MTases, such as Rv3919c, Rv1694, and Rv1988, are another group of mycobacterial MTases responsible for the methylation

of certain nucleotides of rRNA. In turn, they affect various mechanisms in particular resistance to antibiotics^[32]. Most antibiotics target the ribosomes of bacterial cells, and changes in these sites induce antibiotic resistance^[33]. We reported the significant upregulation of Rv3919c in INHR, RIFR, MDR, and XDR strains, as well as a mutation (T to G) in codon 341 in INH^R strain. The streptomycin resistance in M.tb has been demonstrated to be associated with mutations reported in the Rv3919c (gidB) gene, a mycobacterial 16S rRNA MTase. Deletion of the Rv3919c gene in M.tb pathogens alters rRNA methylation and disrupts the binding of streptomycin to 16S rRNA, resulting in streptomycin resistance^[34]. We confirmed resistance to streptomycin in RIFR, MDR, and XDR strains that indicated Rv3919c overexpression. Also, a possible reason for upregulation in INH^R strain may be its detected mutation in our study.

In this study, we detected no mutation in Rv1694 and Rv1988 genes. The upregulation of Rv1694 and downregulation of Rv1988 was observed only in sensitive strains. Rv1694 (tlyA) induces capreomycin sensitivity, and mutation in the Rv1694 gene is associated with the induction of capreomycin resistance^[35]. *Rv1988* is a macrolide-lincosamidestreptomycin resistance gene belonging to the erythromycin resistance rRNA methylase family^[36]. Similar to Rv2966c, Rv1988 is multifunctional that can epigenetically affect host genes and target immune against $M.tb^{[37]}$. defense genes involved in Surprisingly, in this study, we presented the overexpression of Rv1988 only in the drug resistant isolates. Deletion of this gene has been shown to be associated with reduced intracellular survival and is considered as an important virulence factor in $M.tb^{[19]}$. This report confirmed that some MTases (such as Rv2966c and Rv1988) were certainly virulence factors that are involved in hijacking of the host transcriptional machinery and dampening its defense during initial of TB infection^[37]. The variation of virulence may be along with the increasing antibiotic resistance. Thus, in controlling the spread of antibiotic resistance, controlling the spread of virulence is essential^[38].

The findings of the current study suggested the overexpression of Rv1392 in sensitive, RIF^R, MDR, and XDR strains, interestingly highlighting the role of MTases and methyl transfer in the resistance and pathogenesis of $M.tb^{[37]}$. A significant upregulation of Rv1392 has also been reported in M.tb in comparison to mutant mycobacterial strain^[39,40]. We detected a mutation (G to A) in INH^R strain that may be the cause of its downregulation.

In this study, we reported the difference of MTases expression between resistant and sensitive strains

of *M.tb*. Overexpression of Rv0645c, Rv2756c, Rv3263, and Rv2966c was detected in all sensitive and resistant isolates. In the sensitive strain, Rv1988 and Rv3919c decreased but Rv1694 increased. Rv1392 expression level also decreased in INH isolate. Our results can be extended to this finding that mycobacterial MTases and SAM synthetase gene expression are more often reported in sensitive strains. A better understanding of the complete effects of mycobacterial MTase within a particular *M.tb* strain requires more detailed subsequent studies to integrate the data and develop new diagnostic and therapeutic strategies.

DECLARATIONS

Acknowledgments

The authors are grateful to the personnel of Mycobacteriology and Pulmonary Research Department, Pasteur Institute of Iran (Tehran) for their assistance in this project. We tender our apologies to those authors whose deserving research was not cited in this manuscript.

Ethical statement

Not applicable.

Data availability

Data supporting this article are included within the article and supplementary file.

Author contributions

ST and MSZ: performed all the experiments; ST, AF, and GB: analyzed and interpreted data; AF, FV, and MK: read and approved the final manuscript; SDS and AF: designed and supervised clinical study, interpreted data, read and approved the manuscript.

Conflict of interest

None declared.

Funding/support

The current project was supported by a grant (no. B-9426) from the Pasteur Institute of Iran, Tehran, Iran. The authors have no other relevant financial involvement with any organization or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

REFERENCES

1. Chakaya J, Khan M, Ntoumi F, Aklillu E, Fatima R, Mwaba P, Kapata N, Mfinanga S, Hasnain E, Katoto

- PDMC, Bulabula ANH, Sam-Agudu N A, Nachega JB, Tiberi S, McHugh TD, Abubakar I, Zumla A. Global Tuberculosis report 2020–reflections on the global TB burden, treatment and prevention efforts. *International journal of infectious diseases* 2021; **113**(Suppl1): S7-S12
- 2. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C, Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M, Cox EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N, WHO Pathogens Priority List Working Group. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *The lancet. infectious diseases* 2018; **18**(3): 318-327.
- 3. Tarashi S, Ahmadi Badi S, Moshiri A, Nasehi M, Fateh A, Vaziri F, Siadat SD. The human microbiota in pulmonary tuberculosis: Not so innocent bystanders. *Tuberculosis* 2018; **113**: 215-221.
- 4. Vesga JF, Hallett TB, JA Reid M, Singh Sachdeva K, Rao R, Khaparde S, Dave P, Rade k, Kamene M, Omesa E, Masini E, Omale N, Onyango E, Owiti P, Karanja M, Kiplimo R, Alexandru S, Vilc V, Crudu V, Bivol S, Celan C, ArinaminpathyN. Assessing tuberculosis control priorities in high-burden settings: a modelling approach. *The lancet. Global health* 2019; 7(5): e585-e595.
- 5. Grover S, Gangwar R, Jamal S, Ali S, Nisaa K, Ehtesham NZ, Hasnain SE. Mycobacterial methyltransferases: significance in pathogenesis and virulence. *Mycobacterium tuberculosis: molecular infection biology, pathogenesis, diagnostics and new interventions* 2019; 103-122.
- 6. Ding W, Smulan LJ, Hou NS, Taubert S, Watts JL, Walker AK. s-Adenosylmethionine levels govern innate immunity through distinct methylation-dependent pathways. *Cell metabolism* 2015; **22**(4): 633-645.
- 7. Tarashi S, Ahmadi Badi S, Moshiri A, Ebrahimzadeh N, Fateh A, Vaziri F, Aazami H, Siadat SD, Fuso A. The inter-talk between Mycobacterium tuberculosis and the epigenetic mechanisms. *Epigenomics* 2020; **12**(5): 455-469.
- 8. Gong Z, Wang G, Zeng J, Stojkoska A, Huang H, Xie J. Differential DNA methylomes of clinical MDR, XDR and XXDR Mycobacterium tuberculosis isolates revealed by using single-molecule real-time sequencing. *Journal of drug targeting* 2021; **29**(1): 69-77.
- 9. Edwards J.R., Yarychkivska O, Boulard M, Bestor TH. DNA methylation and DNA methyltransferases. *Epigenetics and chromatin* 2017; **10**(1): 1-10.
- Grover S, Gupta P, Kahlon PS, Goyal S, Grover A, Dalal K, Sabeeha, Ehtesham NZ, Hasnain SE. Analyses of methyltransferases across the pathogenicity spectrum of different mycobacterial species point to an extremophile connection. *Molecular biosystems* 2016; 12(5): 1615-1625.
- 11. Miro-Blanch J, Yanes O. Epigenetic regulation at the interplay between gut microbiota and host metabolism. *Frontiers in genetics* 2019; **10**: 638.

- 12. Kathirvel M, Mahadevan S. The role of epigenetics in tuberculosis infection. *Epigenomics* 2016; **8**(4): 537-549
- 13. Caceres N, Vilaplana C, Prats C, Marzo E, Llopis I, Valls J, Lopez D, Cardona P-J. Evolution and role of corded cell aggregation in Mycobacterium tuberculosis cultures. *Tuberculosis* 2013; **93**(6): 690-698.
- van Klingeren B, Dessens-Kroon M, van der Laan T, Kremer K, Soolingen DV. Drug susceptibility testing of Mycobacterium tuberculosis complex by use of a highthroughput, reproducible, absolute concentration method. *Journal of clinical microbiology* 2007; 45(8): 2662-2668.
- 15. Richter E, Rüsch-Gerdes S, Hillemann D. Drugsusceptibility testing in TB: current status and future prospects. *Expert review of respiratory medicine* 2009. **3**(5): 497-510.
- 16. American Thoracic Society. Diagnostic standards and classification of tuberculosis. *The American review of respiratory disease* 1990; **142**(3); 725-735.
- 17. Kozbia PZ, Mushegian AR. Natural history of Sadenosylmethionine-binding proteins. *BMC structural biology* 2005. **5**(1): 1-26.
- Rustad TR, Roberts DM, Liao RP, Sherman DR. Isolation of mycobacterial RNA, in Mycobacteria protocols. *Methods in molecular biology* 2009; 465: 13-22.
- 19. Yaseen I, Kaur P, Nandicoori VK, Khosla S. Mycobacteria modulate host epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3. *Nature communications* 2015; **6**(1): 1-13.
- Shell S.S, Prestwich EG, Baek S-H, Shah RR, Sassetti CM, Dedon PC, Fortune SM. DNA methylation impacts gene expression and ensures hypoxic survival of Mycobacterium tuberculosis. *Plos pathogens* 2013; 9(7): e1003419.
- 21. Srivastava R, Gopinathan K, Ramakrishnan T. Deoxyribonucleic acid methylation in mycobacteria. *Journal of bacteriology* 1981; **148**(2): 716-719.
- Sharma G, Upadhyay S, Srilalitha M, Nandicoori VK, Khosla S. The interaction of mycobacterial protein Rv2966c with host chromatin is mediated through non-CpG methylation and histone H3/H4 binding. *Nucleic* acids research 2015; 43(8): 3922-3937.
- 23. Velayati A.A, Farnia P, Ibrahim TA, Haroun RZ, Kuan HO, Ghanavi J, Farnia P, Naghee Kabarei A, Tabarsi P, Omar AR, Varahram M, Masjedi MR. Differences in cell wall thickness between resistant and nonresistant strains of Mycobacterium tuberculosis: using transmission electron microscopy. *Chemotherapy* 2009; 55(5): 303-307.
- 24. Barkan D, Liu Z, Sacchettini JC, Glickman MS. *My*colic acid cyclopropanation is essential for viability, drug resistance, and cell wall integrity of Mycobacterium tuberculosis. *Chemistry and biology* 2009; **16**(5): 499-509.
- 25. Barry CE, Crick DC, McNeil MR. Targeting the formation of the cell wall core of M. tuberculosis. *Infectious disorders drug targets* 2007; **7**(2): 182-202.

- 26. Cappelli G, Volpe E, Grassi M, Liseo B, Colizzi V, Mariani F. Profiling of Mycobacterium tuberculosis gene expression during human macrophage infection: upregulation of the alternative sigma factor G, a group of transcriptional regulators, and proteins with unknown function. *Research in microbiology* 2006; **157**(5): 445-455.
- Barkan D, Hedhli D, Yan HG, Huygen K, Glickman MS. Mycobacterium tuberculosis lacking all mycolic acid cyclopropanation is viable but highly attenuated and hyperinflammatory in mice. *Infection and immunity* 2012. 80(6): 1958-1968.
- 28. DiNardo A.R, Rajapakshe K, Nishiguchi T, Grimm SL, Mtetwa G, Dlamini Q, Kahari J, Mahapatra S, Kay A, Maphalala G, Mace EM, Makedonas G, Cirillo JD, Netea MG, R Crevel RV, Coarfa C, Mandalakas AM. DNA hypermethylation during tuberculosis dampens host immune responsiveness. *The Journal of clinical investigation* 2020; **130**(6): 3113-3123.
- 29. Ndhlovu V, Kiran A, Sloan D, Mandala W, Nliwasa M, Everett DB, Mwapasa M, Kontogianni K, Kamdolozi M, Corbett EL, Caws M, Davies G. Understanding the diversity of DNA methylation in Mycobacterium tuberculosis. *The preprent server for biology* 2020.
- 30. Sharma G, Sowpati DT, Singh P, Zahoor Khan M, Ganji R, Upadhyay S, Banerjee Sh, Kumar Nandicoori V, Khosla S. Genome-wide non-CpG methylation of the host genome during M. tuberculosis infection. *Scientific reports* 2016; **6**(1): 1-15.
- 31. Khan S.H, Bijpuria S, Maurya A, Taneja B. Structural and thermodynamic characterization of a highly stable conformation of Rv2966c, a 16S rRNA methyltransferase, at low pH. *International journal of biological macromolecules* 2020; **164**: 3909-3921.
- 32. Osterman I, Dontsova O, Sergiev PV. rRNA methylation and antibiotic resistance. *Biochemistry*. *Biokhimiia* 2020; **85**(11): 1335-1349.
- 33. Poehlsgaard J, Douthwaite S. The bacterial ribosome as a target for antibiotics. *Nature reviews. Microbiology* 2005; **3**(11): 870-881.
- 34. Wong S.Y, JavidB, Addepalli B, Piszczek G, Strader MB, Limbach PA, Barry CA. Functional role of methylation of G518 of the 16S rRNA 530 loop by GidB in Mycobacterium tuberculosis. *Antimicrobial agents and chemotherapy* 2013; **57**(12): 6311-6318.
- Witek M.A, Kuiper EG, Minten E, Crispell EK, Conn GA. A novel motif for S-adenosyl-L-methionine binding by the ribosomal RNA methyltransferase TlyA from Mycobacterium tuberculosis. *The Journal of biological chemistry* 2017; 292(5): 1977-1987.
- 36. Buriánková K, Doucet-Populaire F, Dorson O, Gondran A, Ghnassia JC, Weiser J, Pernodet JL. Molecular basis of intrinsic macrolide resistance in the Mycobacterium tuberculosis complex. *Antimicrobial agents and chemotherapy* 2004; **48**(1): 143-150.
- 37. Khosla S, Sharma G, Yaseen I. Learning epigenetic regulation from mycobacteria. *Microbial cell* 2016; **3**(2): 92.

[DOR: 20.1001.1.1028852.2022.26.3.6.1]


- 38. Zhan L, Wang J, Wang L, Qin C. The correlation of drug resistance and virulence in Mycobacterium tuberculosis. *Biosafety and health* 2020; **2**(01): 18-24.
- 39. Mehta M, Rajmani R.S, Singh A. Mycobacterium tuberculosis WhiB3 responds to vacuolar pH-induced changes in mycothiol redox potential to modulate phagosomal maturation and virulence. *The Journal of biological chemistry* 2016; **291**(6): 2888-2903.
- 40. Nixon M.R, Saionz KW, Koo MS, Szymonifka MJ, Jung H, Roberts JP, Nandakumar M, Kumar A, Liao R, Rustad T, Sacchettini JC, Rhee KY, Freundlich JS, Sherman DR. Folate pathway disruption leads to critical disruption of methionine derivatives in Mycobacterium tuberculosis. *Chemistry and biology* 2014; 21(7): 819-830.


Supplementary Table 1. Amplification parameters of PCR reaction

Step	Time	Temperature °C	Cycle
Initial Denaturing	3 min	94	1
Denaturing	45 s	94	
Annealing	45 s	54	35
Extension	45 s	72	
Melting	10 min	72	1

Supplementary Table 2. Amplification parameters of real-time PCR reaction

Holding	Cycling (40 cycle)	Melting
95 °C/60 s	95 °C/5 min	95 °C/5 s
93 C/00 S	50 °C/30 s	60 °C/60 s
	72 °C/30 sec	95°C/1 s





Supplementary Fig. 1. The PCR products of mycobacterial MTases and *Rv1392* genes electrophoresed on 2% (w/v) agarose gel. The related fragments was observed and verified by sequencing. DNA ladder (100 bp); NegCont, negative control.